arXiv:1003.1012v2 [math.QA] 14 Sep 2010

ELLIPTIC ASSOCIATORS

BENJAMIN ENRIQUEZ

Abstract. We study the action of the arithmetic fundamental group of the moduli space of elliptic curves on completions of braid groups in genus 1. This action factors through an explicit profinite group $\widehat{\mathrm{GT}}_{ell}$, which is an analogue of the Grothendieck–Teichmüller group and may be viewed as the group of universal automorphisms of elliptic structures over braided monoidal categories. We relate the action of $\widehat{\mathrm{GT}}_{ell}$ on braid groups in genus 1 to that of a prounipotent version $\mathrm{GT}_{ell}(-)$ of $\widehat{\mathrm{GT}}_{ell}$. We prove that the Lie algebra of $\mathrm{GT}_{ell}(-)$ is isomorphic to a graded Lie algebra, using a scheme of elliptic associators. This enables us to compute the Zariski closure of the image of the mapping class group in genus one B_3 in the automorphism groups of the prounipotent completions of the braid groups in genus 1. We study algebraic relations between multiple zeta values (MZVs) and iterated Mellin transforms of Eisenstein series, derived from the action of B_3 on a particular elliptic associator.

Introduction

In this paper, we develop an elliptic analogue of the theory of associators and the Grothendieck–Teichmüller group. We first review some results in this theory (Section 1). We then state our main results (Section 2): the construction and properties of a group scheme related to representations of the arithmetic fundamental group of the moduli space of elliptic curves (Theorem 7); the computation of the Zariski closure of the image of the mapping class group in genus one in natural representations (Theorem 8); algebraic relations between iterated Mellin transforms of Eisenstein series and multizeta values (MZVs).

We then develop the theory leading to the proof of these results. We introduce the notion of an elliptic structure over a braided monoidal category (BMC) (Section 3); such a structure gives rise to representations of braid groups in genus 1. The universal endomorphism semigroup of elliptic structures is an analogue $\underline{\mathrm{GT}}_{ell}$ of the Grothendieck–Teichmüller semigroup $\underline{\mathrm{GT}}$ (Section 4), which we identify with an extension $\widetilde{\mathrm{GL}_2\,\mathbb{Z}}$ of $\mathrm{GL}_2(\mathbb{Z})$ by $\mathbb{Z}$; $\underline{\mathrm{GT}}_{ell}$ admits a prounipotent version $\mathrm{GT}_{ell}(\mathbf{k})$, which decomposes as a semidirect product $\mathrm{GT}_{ell}(\mathbf{k}) \simeq R_{ell}(\mathbf{k}) \rtimes \mathrm{GT}(\mathbf{k})$. We construct a scheme $Ell(\mathbf{k})$ of elliptic associators (Section 5); elements of $Ell(\mathbf{k})$ give rise to elliptic structures over particular BMCs, and yield formality isomorphisms of braid groups in genus 1. $Ell(\mathbf{k})$ is a torsor over $\mathrm{GT}_{ell}(\mathbf{k})$; we have a natural torsor morphism $Ell(\mathbf{k}) \to M(\mathbf{k})$ to the scheme of associators, which admits a section $M(\mathbf{k}) \xrightarrow{\sigma} Ell(\mathbf{k})$. We study the group $\mathrm{GRT}_{ell}(\mathbf{k})$ of $\mathrm{GT}_{ell}(\mathbf{k})$-automorphisms of $Ell(\mathbf{k})$ (Section 6). We have again a semidirect product decomposition $\mathrm{GRT}_{ell}(\mathbf{k}) \simeq R^{gr}_{ell}(\mathbf{k}) \rtimes \mathrm{GRT}(\mathbf{k})$, compatible with the torsor morphisms; moreover, $\mathrm{Lie}\,\mathrm{GRT}_{ell}$ is graded. The nonemptiness of $Ell(\mathbf{k})$ then implies that $\mathrm{Lie}\,\mathrm{GT}_{ell}$ is isomorphic to a graded Lie algebra. We exhibit an explicit graded Lie subalgebra $\mathfrak{b}_3 \subset \mathrm{Lie}\, R^{gr}_{ell}$; this Lie algebra arose in the study of a universal version of the Knizhnik–Zamolodchikov–Bernard (KZB) connection in [CEE] from the explicit comparison of 'infinitesimally close' formality isomorphisms.

The analogue of the KZ associator is a map $\mathfrak{H} \xrightarrow{e} Ell(\mathbb{C})$ (where $\mathfrak{H}$ is the Poincaré half-plane); the $e(\tau)$, $\tau \in \mathfrak{H}$, give rise to the formality isomorphisms from [CEE]. We relate $\sigma(\Phi_{KZ})$ to a renormalization of $e(\tau)$ (Φ_{KZ} is the KZ associator), and we study the behavior of the map $\tau \mapsto e(\tau)$ under the action of $B_3 \simeq \widetilde{\mathrm{SL}_2\,\mathbb{Z}} \subset \underline{\mathrm{GT}}_{ell}$ (Section 7).

These results enable us to compute the morphism $B_3 \overset{\text{can}}{\hookrightarrow} R_{ell}(\mathbb{C}) \to R^{gr}_{ell}(\mathbb{C})$ associated with $\sigma(\Phi_{KZ})$, ; it gives rise to an isomorphism $\mathrm{Lie}\langle B_3\rangle^{\mathbb{C}} \simeq \hat{\mathfrak{b}}_3^{\mathbb{C}}$ of the Lie algebra of the Zariski closure of the canonical morphism $B_3 \to R_{ell}(\mathbb{Q})$ with the degree completion of $\mathfrak{b}_3$ (Section 8).

We gather some conjectures in Section 9. We conjecture that the inclusion $\mathfrak{b}_3 \subset \mathrm{Lie}\, R^{gr}_{ell}$ is actually an equality (Generation Conjecture). We relate this conjecture to the question of the Zariski density of a canonical morphism $B_3 \to R_{ell}(\mathbf{k})$ and to a (very bold) Transcendence Conjecture on the coefficients of Φ_{KZ}.

We then prove the results of Section 2. The results on representations of arithmetic fundamental groups and on Zariski closures are proved in Section 10. The theory of iterated Mellin transforms of modular forms is developed in Section 11; there we also prove the results on Mellin transforms of Eisenstein series announced in Section 2.

Acknowledgements. The author would like to thank P. Etingof and D. Calaque for discussions and their collaboration in [CEE], and J. Barge, H. Furusho and D. Harari for discussions.

1. The theory of associators

1.1. **Profinite Galois representations.**

Let $n \geq 3$ and $M^{\mathbb{Q}}_{0,n}$ be the moduli stack over $\mathbb{Q}$ of genus 0 curves with n marked points. It is equipped with a collection of rational tangential base points corresponding to maximally degenerate curves, which we may view a sections of the morphism $M^{\mathbb{Q}}_{0,n} \to \mathrm{Spec}\,\mathbb{Q}$.

We denote by $T_{0,n}$ the fundamental groupoid of $M^{\mathbb{Q}}_{0,n}$ with respect to these base points, and by $\widehat{T}_{0,n}$ its profinite completion. We then have a group morphism $G_{\mathbb{Q}} = \mathrm{Gal}(\bar{\mathbb{Q}}/\mathbb{Q}) \to \mathrm{Aut}(\widehat{T}_{0,n})$ (see [G1, Dr]).

Theorem 1. ([Dr, Sch]) *This morphism factors as $G_{\mathbb{Q}} \to \widehat{\mathrm{GT}} \to \mathrm{Aut}(\widehat{T}_{0,n})$, where $\widehat{\mathrm{GT}}$ is an 'explicit' profinite group.*

Here 'explicit' means that $\widehat{\mathrm{GT}}$ is defined by a collection of equations in a product of profinite groups.

1.2. **Pro-l and prounipotent completions.**

If π is a group and l is a prime number, we denote by π_l and $\pi(-)$ be its pro-l and $\mathbb{Q}$-prounipotent completions; $\pi(-)$ is a $\mathbb{Q}$-group scheme, i.e., a functor $\{\mathbb{Q}\text{-rings}\} \to \{\text{groups}\}$. If π is finitely generated, then we have a morphism $\pi_l \to \pi(\mathbb{Q}_l)$ (see [HM], Lemma A.7).

Recall that if $G(-)$ is a $\mathbb{Q}$-group scheme, then its Lie algebra is $\mathrm{Lie}(G) := \mathrm{Ker}(G(\mathbb{Q}[\varepsilon]/(\varepsilon^2)) \to G(\mathbb{Q}))$. Then $\mathfrak{g} := \mathrm{Lie}\,\pi(-)$ is a pronilpotent $\mathbb{Q}$-Lie algebra. For $\mathbf{k}$ a $\mathbb{Q}$-ring, we set $\mathfrak{g}_{\mathbf{k}} := \lim_{\leftarrow}(\mathfrak{g}/\mathfrak{g}_n) \otimes \mathbf{k}$ (where $\mathfrak{g}_0 = \mathfrak{g}$, $\mathfrak{g}_{n+1} = [\mathfrak{g}, \mathfrak{g}_n]$).

We then define a $\mathbb{Q}$-group scheme $\underline{\mathrm{Aut}\,\pi}(-)$ by $\underline{\mathrm{Aut}\,\pi}(\mathbf{k}) := \mathrm{Aut}_{\mathrm{Lie}}(\mathfrak{g}_{\mathbf{k}})$. We then have a morphism $\underline{\mathrm{Aut}\,\pi}(-) \to \mathrm{GL}(\mathfrak{g}^{\mathrm{ab}})$ with prounipotent kernel (here $\mathfrak{g}^{\mathrm{ab}} = \mathfrak{g}/\mathfrak{g}_1$). As $\pi(\mathbf{k}) = \exp(\mathfrak{g}_{\mathbf{k}})$, we have a group morphism $\underline{\mathrm{Aut}\,\pi}(\mathbf{k}) \to \mathrm{Aut}(\pi(\mathbf{k}))$ for any $\mathbb{Q}$-ring $\mathbf{k}$, denoted $\theta \mapsto \theta_*$. Note that $\mathrm{Lie}\,\underline{\mathrm{Aut}\,\pi} = \mathrm{Der}\,\mathrm{Lie}\,\pi$.

We then define a group $\mathrm{Aut}(\pi(\mathbb{Q}_l), \pi_l) := \{(\theta, \theta_l) \in \underline{\mathrm{Aut}\,\pi}(\mathbb{Q}_l) \times \mathrm{Aut}(\pi_l) | \theta_* i = i\theta_l\}$, where $i : \pi_l \to \pi(\mathbb{Q}_l)$ is as above. We have morphisms

$$\mathrm{Aut}(\pi_l) \leftarrow \mathrm{Aut}(\pi(\mathbb{Q}_l), \pi_l) \to \underline{\mathrm{Aut}\,\pi}(\mathbb{Q}_l). \tag{1}$$

Let now $\mathcal{G} \rightrightarrows B$ be a groupoid where for any $b \in B$, $\mathcal{G}(b)$ is finitely generated. We denote by $\mathcal{G}_l$, $\mathcal{G}(-)$ its pro-l and prounipotent completions, given by $\mathcal{G}_l(b,c) := \mathcal{G}(b)_l \times_{\mathcal{G}(b)} \mathcal{G}(b,c)$ and $\mathcal{G}(\mathbf{k})(b,c) := \mathcal{G}(b)(\mathbf{k}) \times_{\mathcal{G}(b)} \mathcal{G}(b,c)$. We have a morphism $\mathcal{G}_l \to \mathcal{G}(\mathbb{Q}_l)$. $\mathcal{G}(-)$ is a prounipotent groupoid scheme, i.e., a functor $\{\mathbb{Q}\text{-rings}\} \to \{\text{groupoids}\}$, such that for each $b \in B$, $\mathcal{G}(-)(b)$ is prounipotent.

Assume that $\mathcal{G}$ is connected (i.e., for any $b, c \in B$, $\mathcal{G}(b,c) \neq \emptyset$). Then the choice of $b \in B$ and $g_{bc} \in \mathcal{G}(b,c)$ for any $c \in B - \{b\}$ gives rise to an isomorphism $\mathrm{Aut}(\mathcal{G}) \simeq \pi^{B-\{b\}} \rtimes \mathrm{Aut}(\pi)$ where $\pi = \mathcal{G}(b)$; the inverse map takes $(\theta, (g_c)_{c\in B-\{b\}})$ to the automorphism whose restriction to $\mathcal{G}(b)$ is θ and such that $g_{bc} \mapsto g_{bc}g_c$. We then define a $\mathbb{Q}$-group scheme $\underline{\mathrm{Aut}\,\mathcal{G}}(-)$ by $\underline{\mathrm{Aut}\,\mathcal{G}}(\mathbf{k}) := \pi(\mathbf{k})^{B-\{b\}} \rtimes \underline{\mathrm{Aut}\,\pi}(\mathbf{k})$; we have a group morphism $\underline{\mathrm{Aut}\,\mathcal{G}}(\mathbf{k}) \to \mathrm{Aut}(\mathcal{G}(\mathbf{k}))$ for any $\mathbb{Q}$-ring $\mathbf{k}$. We define as above a group $\mathrm{Aut}(\mathcal{G}(\mathbb{Q}_l), \mathcal{G}_l)$, equipped with morphisms

$$\mathrm{Aut}(\mathcal{G}_l) \leftarrow \mathrm{Aut}(\mathcal{G}(\mathbb{Q}_l), \mathcal{G}_l) \to \underline{\mathrm{Aut}\,\mathcal{G}}(\mathbb{Q}_l).$$

1.3. **Pro-l Galois representations and the group scheme** $\mathrm{GT}(-)$**.** We now study the composite morphism $G_{\mathbb{Q}} \to \mathrm{Aut}(\widehat{T}_{0,n}) \to \mathrm{Aut}(T^l_{0,n})$. The following theorem can be derived from [Dr, Sch].

Theorem 2. *1) This morphism admits a factorization* $G_{\mathbb{Q}} \to \mathrm{Aut}(T_{0,n}(\mathbb{Q}_l), T^l_{0,n}) \to \mathrm{Aut}(T^l_{0,n})$.

2) There exists a $\mathbb{Q}$*-group scheme* $\mathrm{GT}(-)$ *with the following properties:*

(a) we have an epimorphism $\mathrm{GT}(-) \to \mathbb{G}_m$ *with a prounipotent kernel;*

(b) for any l*, we have a morphism* $G_{\mathbb{Q}} \to \mathrm{GT}(\mathbb{Q}_l)$*, fitting in a diagram*

$$\begin{tikzcd} G_{\mathbb{Q}} \arrow[r] \arrow[d] & \widehat{\mathbb{Z}}^{\times} \arrow[d] \\ \mathrm{GT}(\mathbb{Q}_l) \arrow[r] & \mathbb{Q}_l^{\times} \end{tikzcd}$$

(c) we have a morphism $\mathrm{GT}(-) \to \underline{\mathrm{Aut}\,T_{0,n}}(-)$*, such that the diagram*

$$\begin{tikzcd} G_{\mathbb{Q}} \arrow[r] \arrow[dr] \arrow[dd] & \mathrm{Aut}(T^l_{0,n}) \\ & \mathrm{Aut}(T_{0,n}(\mathbb{Q}_l), T^l_{0,n}) \arrow[u] \arrow[d] \\ \mathrm{GT}(\mathbb{Q}_l) \arrow[r] & \underline{\mathrm{Aut}\,T_{0,n}}(\mathbb{Q}_l) \end{tikzcd}$$

commutes.

Let us say that a $\mathbb{Q}$-group scheme $\pi(-)$ is graded if $\mathfrak{g} := \mathrm{Lie}\,\pi$ is equipped with a decomposition $\mathfrak{g} = \hat{\oplus}_{k\geq 0}\mathfrak{a}_k$, such that $[\mathfrak{a}_k, \mathfrak{a}_l] \subset \mathfrak{a}_{k+l}$. A morphism $\pi(-) \to \pi'(-)$ is called graded iff the map $\mathfrak{g} \to \mathfrak{g}'$ preserves the gradings. If $\pi(-)$ is graded and prounipotent, then $\mathfrak{a}_0 = 0$ and $\underline{\mathrm{Aut}\,\pi}(-)$ is a graded $\mathbb{Q}$-group scheme.

A connected $\mathbb{Q}$-groupoid $\mathcal{G}(-)$ is called graded, prounipotent, etc., if $\mathcal{G}(-)(b)$ has the same properties for some (equivalently, any) $b \in B$. If $\mathcal{G}^{gr}(-)$ is a graded prounipotent group scheme, then $\underline{\mathrm{Aut}\,\mathcal{G}^{gr}}(-)$ is a graded group scheme. The following theorem may be derived from the theory of associators.

Theorem 3. *(*[Dr, E]*) There exists a graded group scheme* $\mathrm{GRT}(-)$*, a prounipotent graded groupoid scheme* $T^{gr}_n(-)$*, a graded morphism* $\mathrm{GRT}(-) \to \underline{\mathrm{Aut}\,T^{gr}_n}(-)$ *and isomorphisms* $\mathrm{GT}(-) \to \mathrm{GRT}(-)$*,* $T_n(-) \to T^{gr}_n(-)$*, such that the diagram*

$$\begin{tikzcd} \mathrm{GT}(-) \arrow[r] \arrow[d] & \underline{\mathrm{Aut}\,T_n}(-) \arrow[d] \\ \mathrm{GRT}(-) \arrow[r] & \underline{\mathrm{Aut}\,T^{gr}_n}(-) \end{tikzcd}$$

commutes.

1.4. **Multiple zeta values.** The main ingredient in the proof of Theorem 3 is the theory of associators. This theory also applies to the theory of relations between multiple zeta values (MZVs). These numbers are defined by

$$\zeta(k_1,\ldots,k_m) := \sum_{0<n_1<\ldots<n_m} \frac{1}{n_1^{k_1}\cdots n_m^{k_m}}$$

for $m, k_1, \ldots, k_m \in \mathbb{Z}_{>0}$ and $k_m > 1$.

Define then the noncommutative formal power series

$$\Phi_{KZ} := 1 + \sum_{m>0} \sum_{k_1,\ldots,k_{m-1}>0, k_m>1} \zeta(k_1,\ldots,k_m)\psi(A^{k_1-1}B\cdots A^{k_m-1}B),$$

where for $w \in \mathbb{C}\langle\langle A,B\rangle\rangle$, $\psi(w) = \sum_{k,l\geq 0}(-1)^{k+l}B^l\partial_A^k\partial_B^l(w)A^k$, and ∂_A, ∂_B are the commuting derivations such that $\partial_A : (A,B) \mapsto (1,0)$ and $\partial_B : (A,B) \mapsto (0,1)$.

Theorem 4. *([Dr]) Φ_{KZ} is group-like in the structure for which A, B are primitive. It satisfies the duality, hexagon and pentagon relations*

$$\Phi_{KZ}(B,A) = \Phi_{KZ}(A,B)^{-1}, \quad \Phi_{KZ}(A,B)e^{\mathrm{i}\,\pi A}\Phi_{KZ}(C,A)e^{\mathrm{i}\,\pi C}\Phi_{KZ}(B,C)e^{\mathrm{i}\,\pi B} = 1,$$

$$\Phi_{KZ}(t_{23},t_{34})\Phi_{KZ}(t_{12}+t_{13},t_{24}+t_{34})\Phi_{KZ}(t_{12},t_{23}) = \Phi_{KZ}(t_{12},t_{23}+t_{24})\Phi_{KZ}(t_{13}+t_{23},t_{34})$$

where the t_{ij} are subject to $t_{ji} = t_{ij}$, $[t_{ij}, t_{ik}+t_{jk}] = 0$, $[t_{ij}, t_{kl}] = 0$ for i,j,k,l distinct, and $A+B+C = 0$.

Other families of relations between MZVs are known, like the stuffle relations; however, it has been shown ([F2]) that these relations are implied by the pentagon relation. It has also been shown ([F1]) that the pentagon relation implies the hexagon and duality ones.

2. The main results

This theory is based on the action of the algebraic fundamental group $\pi_1(M_{1,1^2}^{\mathbb{Q}})$ on completions of elliptic braid groups, based on the fibration $M_{1,n}^{\mathbb{Q}} \to M_{1,1}^{\mathbb{Q}}$, as studied in [G2, O].

2.1. **Profinite Galois representations.** Let $M_{1,1^2}^{\mathbb{Q}}$ (resp., $M_{1,n}^{\mathbb{Q}}$) be the moduli space of triples $(E, p, \vec{v})$ of an elliptic curve, a marked point and a nonzero tangent vector at this point (resp., of pairs $(E, \underline{p})$ of an elliptic curve and an injection $\underline{p} : \{1,\ldots,n\} \to E$). A rational tangential base point ξ on $M_{1,1^2}^{\mathbb{Q}}$ is provided by the Tate elliptic curve $\mathbb{G}_m/q^{\mathbb{Z}}$, with marked point $(q^{\mathbb{Z}}, z\frac{d}{dz})$, at the formal neighborhood of $q = 0$. We have a fibration $M_{1,n}^{\mathbb{Q}} \to M_{1,1}^{\mathbb{Q}}$ and if we set $M_{1;1^2,n-1}^{\mathbb{Q}} := M_{1,n^{\mathbb{Q}}}^{\mathbb{Q}} \times_{M_{1,1}^{\mathbb{Q}}} M_{1,1^2}^{\mathbb{Q}}$, we derive a fibration $M_{1;1^2,n-1}^{\mathbb{Q}} \to M_{1,1^2}^{\mathbb{Q}}$.

We also have a collection $(\sigma_T)_{T\in\{trees\}}$ of sections of this fibration, indexed by rooted trivalent planar trees, with leaves indexed by $\{1,\ldots,n\}$ (the root is not a leaf), modulo 'mirror symmetry'. Let $T_{ell,n}$ be the fundamental groupoid of the fiber of ξ with respect to the points $\sigma_T(\xi)$.

As in [G2, O], we get a morphism

$$\pi_1(M_{1,1^2}^{\mathbb{Q}}, \xi) \to \mathrm{Aut}(\widehat{T}_n^{ell}). \tag{2}$$

Note that we also have an exact sequence $1 \to \widehat{B}_3 \to \pi_1(M_{1,1^2}^{\mathbb{Q}}, \xi) \to G_{\mathbb{Q}} \to 1$; a section is provided by the base point ξ, it gives rise to an isomorphism

$$\pi_1(M_{1,1^2}^{\mathbb{Q}}) \simeq \widehat{B}_3 \rtimes G_{\mathbb{Q}}.$$

The morphism $G_{\mathbb{Q}} \to \mathrm{Aut}(\widehat{B}_3)$ has been computed in [Na], Cor. 4.15 (we recall it in Subsection 10.1).

Proposition 5. *We have a commutative diagram*

$$\begin{tikzcd} \pi_1(M^{\mathbb{Q}}_{1,1^2},\xi) \arrow[r, "\curvearrowleft"] \arrow[d] & G_{\mathbb{Q}} \arrow[d] \\ \mathrm{GL}_2(\widehat{\mathbb{Z}}) \arrow[r, "\curvearrowleft"] & \widehat{\mathbb{Z}}^{\times} \end{tikzcd}$$

where bottom morphisms are det *and* $\lambda \mapsto \left(\begin{smallmatrix} \lambda & 0 \\ 0 & 1 \end{smallmatrix}\right)$ *and the right map is the cyclotomic character.*

Theorem 6. *The morphism (2) factors as* $\pi_1(M^{\mathbb{Q}}_{1,1^2}) \to \widehat{\mathrm{GT}}_{ell} \to \mathrm{Aut}(\widehat{T}^{ell}_n)$*, where* $\widehat{\mathrm{GT}}_{ell}$ *is an 'explicit' profinite group. Moreover, we have a semidirect product decomposition* $\widehat{\mathrm{GT}}_{ell} \simeq \widehat{R}_{ell} \rtimes \widehat{\mathrm{GT}}$*, compatible with the semidirect product decomposition of* $\pi_1(M^{\mathbb{Q}}_{1,1^2})$.

Here 'explicit' has the same sense as above.

2.2. **Pro-l Galois representations.** We now study the composite map $\pi_1(M^{\mathbb{Q}}_{1,1^2}) \to \mathrm{Aut}(\widehat{T}_{ell,n}) \to \mathrm{Aut}(T^l_{ell,n})$. Its restriction to $G_{\mathbb{Q}}$ is computed in [Na] for $n = 1, 2$.

Theorem 7. *1) This morphism admits a factorization* $\pi_1(M^{\mathbb{Q}}_{1,1^2}) \to \mathrm{Aut}(T_{ell,n}(\mathbb{Q}_l), T^l_{ell,n}) \to \mathrm{Aut}(T^l_{ell,n})$.

2) There exists a group scheme $\mathrm{GT}_{ell}(-)$ *with a semidirect product structure* $\mathrm{GT}_{ell}(-) \simeq R_{ell}(-) \rtimes \mathrm{GT}(-)$ *and the following properties:*

(a) we have a diagram

$$\begin{tikzcd} \mathrm{GT}_{ell}(-) \arrow[r, "\curvearrowleft"] \arrow[d] & \mathrm{GT}(-) \arrow[d] \\ \mathrm{GL}_2 \arrow[r, "\curvearrowleft"] & \mathbb{G}_m \end{tikzcd}$$

where the vertical morphisms have prounipotent kernels;

(b) for any l, we have a morphism $\pi_1(M^{\mathbb{Q}}_{1,1^2}) \to \mathrm{GT}_{ell}(\mathbb{Q}_l)$*, which extends to a morphism from*

$$\begin{tikzcd} \pi_1(M^{\mathbb{Q}}_{1,1^2}) \arrow[r, "\curvearrowleft"] \arrow[d] & G_{\mathbb{Q}} \arrow[d] \\ \mathrm{GL}_2(\widehat{\mathbb{Z}}) \arrow[r, "\curvearrowleft"] & \widehat{\mathbb{Z}}^{\times} \end{tikzcd}$$

to

$$\begin{tikzcd} \mathrm{GT}_{ell}(\mathbb{Q}_l) \arrow[r, "\curvearrowleft"] \arrow[d] & \mathrm{GT}(\mathbb{Q}_l) \arrow[d] \\ \mathrm{GL}_2(\mathbb{Q}_l) \arrow[r, "\curvearrowleft"] & \mathbb{Q}_l^{\times} \end{tikzcd}$$

(c) there is a morphism $\mathrm{GT}_{ell}(-) \to \underline{\mathrm{Aut}\,T_{ell,n}}(-)$*, such that*

$$\begin{tikzcd} \pi_1(M^{\mathbb{Q}}_{1,1^2}) \arrow[r] \arrow[dr] \arrow[dd] & \mathrm{Aut}(T^l_{ell,n}) \\ & \mathrm{Aut}(T_{ell,n}(\mathbb{Q}_l), T^l_{ell,n}) \arrow[u] \arrow[d] \\ \mathrm{GT}_{ell}(\mathbb{Q}_l) \arrow[r] & \underline{\mathrm{Aut}\,T_{ell,n}}(\mathbb{Q}_l) \end{tikzcd}$$

commutes.

3) There exists a graded group scheme $\mathrm{GRT}_{ell}(-)$*, a prounipotent group scheme* $T^{gr}_{ell,n}(-)$ *and isomorphisms* $\mathrm{GT}_{ell}(-) \to \mathrm{GRT}_{ell}(-)$, $T_{ell,n}(-) \to T^{gr}_{ell,n}(-)$*, such that the diagram*

$$\begin{tikzcd} \mathrm{GT}_{ell}(-) \arrow[r] \arrow[d] & \underline{\mathrm{Aut}\,T_{ell,n}}(-) \arrow[d] \\ \mathrm{GRT}_{ell}(-) \arrow[r] & \underline{\mathrm{Aut}\,T^{gr}_{ell,n}}(-) \end{tikzcd}$$

commutes.

The third statement follows from the theory of elliptic associators developed in this paper. Using the morphism $B_3 \to R_{ell}(\mathbb{Q})$, one derives from it the following statements.

2.3. **Computation of Zariski closures.** Let $P_{1,n}$ be the pure braid group of n points on a curve of genus 1. We have an identification $B_3 \simeq \Gamma_{1,1^2}$ between the the Artin braid group in 3 strands and the mapping class group of a surface of type (1,1) with marked boundary. Therefore we have morphisms $B_3 \to \operatorname{Aut} P_{1,n} \to \underline{\operatorname{Aut} P_{1,n}}(\mathbb{Q})$. Let $\langle B_3\rangle(-) \hookrightarrow \underline{\operatorname{Aut} P_{1,n}}(-)$ be the Zariski closure of B_3.

On the other hand, let $\mathfrak{t}_{1,2}$ be the $\mathbb{Q}$-Lie algebra with generators $x^\pm, t$ and relation $[x^+, x^-] + t = 0$. $\operatorname{Der}_*(\mathfrak{t}_{1,2}) := \{\delta \in \operatorname{Der} \mathfrak{t}_{1,2} | \delta(t) = 0\}$ is graded by $\deg x^\pm = 1$. There are uniquely defined derivations e_+, e_-, δ_{2k} $(k \geq 0)$ in $\operatorname{Der}_*(\mathfrak{t}_{1,2})$, such that e_+, e_- extend the actions of $\left(\begin{smallmatrix} 0 & 1 \\ 0 & 0 \end{smallmatrix}\right), \left(\begin{smallmatrix} 0 & 0 \\ 1 & 0 \end{smallmatrix}\right) \in \mathfrak{sl}_2$ on $\operatorname{Span}(x_+, x_-)$, δ_{2k} has degree $2k+2$ and $\delta_{2k}(x_+) = \operatorname{ad}(x_+)^{2k+2}(x_-)$. We denote by $\mathfrak{b}_3 := \langle \mathfrak{sl}_2, \delta_{2k}, k \geq 0\rangle$ the Lie subalgebra of $\operatorname{Der}_*(\mathfrak{t}_{1,2})$ generated by these elements; it inherits the grading of $\operatorname{Der}_*(\mathfrak{t}_{1,2})$.

Theorem 8. *1) The Lie algebra* $\operatorname{Lie}\langle B_3\rangle$ *is isomorphic to the degree completion of* $\mathfrak{b}_3$.

2) There exists a positively graded Lie algebra $\mathfrak{t}_{1,n}$*, an isomorphism* $\operatorname{Lie} P_{1,n} \simeq \hat{\mathfrak{t}}_{1,n}$*, and a graded morphism* $\mathfrak{b}_3 \to \operatorname{Der} \mathfrak{t}_{1,n}$*, such that the diagram*

$$\begin{tikzcd}
\operatorname{Lie}\langle B_3\rangle \arrow[r] \arrow[d, "\sim"'] & \operatorname{Der}\operatorname{Lie} P_{1,n} \arrow[d, "\sim"] \\
\hat{\mathfrak{b}}_3 \arrow[r] & \operatorname{Der} \hat{\mathfrak{t}}_{1,n}
\end{tikzcd}$$

commutes.

An explicit presentation of $\mathfrak{t}_{1,n}$ is by generators $x_i^\pm$, $i \in \{1, \ldots, n\}$ and relations $\sum_i x_i^\pm = 0$, $[x_i^\pm, x_j^\pm] = 0$, $[x_j^+, x_k^-] = [x_k^+, x_j^-](= t_{jk})$ for $k \neq j$, $[x_i^\pm, t_{jk}] = 0$ for i, j, k distinct. The action of $\mathfrak{b}_3$ on $\mathfrak{t}_{1,n}$ is then $e_\pm \cdot x_i^\mp = x_i^\pm$, $e_\pm \cdot x_i^\pm = 0$, $\delta_{2m} \cdot x_i^+ = \operatorname{ad}(x_i^+)^{2m+2}(x_i^-)$, $\delta_{2m} \cdot x_i^- = \frac{1}{2}\sum_{p+q=2k+1}(-1)^p[(\operatorname{ad} x_i^+)^p(x_i^-), (\operatorname{ad} x_i^+)^q(x_i^-)]$.

2.4. **Iterated Mellin transforms of Eisenstein series and MZVs.** In Section 11, we introduce the iterated Mellin transform $L^*_{f_1,\ldots,f_n}(s_1, \ldots, s_n)$ of a family of modular forms $f_1, \ldots, f_n$. We set $L^*_{l_1,\ldots,l_n} := L^*_{f_1,\ldots,f_n}$ for $f_i = E_{2l_i+2}$, where E_{2l+2} is the Eisenstein series of weight $2l+2$ (equal to 1 at $\mathrm{i}\,\infty$). We will give relations satisfied by the numbers

$$L^\sharp_{l_1,\ldots,l_n}(k_1{+}1, \ldots, k_n{+}1) := (2\pi\,\mathrm{i})^{2(l_1+\cdots+l_n)+n}\,\mathrm{i}^{k_1+\cdots+k_n+n}\,L^*_{l_1,\ldots,l_n}(k_1{+}1, \ldots, k_n{+}1), \quad k_i = 0, \ldots, 2k_i$$

and between these numbers and the MZVs.

Let $\tilde{\mathfrak{b}}_3$ be the complex Lie algebra with generators $e_\pm, \tilde\delta_{2m}$, $m \geq 0$, and relations $[[e_+, e_-], e_\pm] = \pm 2e_\pm$, $[e_-, \tilde\delta_{2m}] = (\operatorname{ad} e_+)^{2m+1}(\tilde\delta_{2m}) = 0$. It is graded by $\deg e_\pm = 0$, $\deg \tilde\delta_{2m} = 2m+2$. The action of $\mathfrak{sl}_2 = \langle e_+, e_-\rangle$ on the positive degree part $\tilde{\mathfrak{b}}_3^+$ of $\tilde{\mathfrak{b}}_3$ extends to an action of $\mathrm{PSL}_2(\mathbb{C})$.

We have a Lie algebra morphism $\tilde{\mathfrak{b}}_3 \to \mathfrak{b}_3^{\mathbb{C}}$, $e_\pm \mapsto e_\pm$, $\tilde\delta_{2m} \mapsto \delta_{2m}$, which prolongates into $\tilde{\mathfrak{b}}_3 \twoheadrightarrow \mathfrak{b}_3^{\mathbb{C}} \hookrightarrow \operatorname{Der}(\mathfrak{t}_{1,2}^{\mathbb{C}})$; the latter morphism extends to an algebra morphism $U(\tilde{\mathfrak{b}}_3) \to \operatorname{End}_{Vect}(U\mathfrak{t}_{1,2}^{\mathbb{C}})$, compatible with the morphism $\mathrm{SL}_2(\mathbb{C}) \to \operatorname{Aut}(U\mathfrak{t}_{1,2}^{\mathbb{C}})$ induced by $\left(\begin{smallmatrix} a & b \\ c & d \end{smallmatrix}\right)\left(\begin{smallmatrix} x_i^+ \\ x_i^- \end{smallmatrix}\right) = \left(\begin{smallmatrix} ax_i^+ + bx_i^- \\ cx_i^+ + dx_i^- \end{smallmatrix}\right)$. The matrix realization of $\mathfrak{sl}_2 = \langle e_+, e_-\rangle$ is then $e_- \leftrightarrow \left(\begin{smallmatrix} 0 & 1 \\ 0 & 0 \end{smallmatrix}\right)$, $e_+ \leftrightarrow \left(\begin{smallmatrix} 0 & 0 \\ 1 & 0 \end{smallmatrix}\right)$.

Define $(a_{2l})_{l\geq 0}$ by $1 + \sum_{l\geq 0} a_{2l}x^{2l+2} = (\frac{x}{e^{x/2} - e^{-x/2}})^2$.

Theorem 9. *Let*

$$\tilde\theta := \sum_{n\geq 0}\sum_{l_1,\ldots,l_n>0}\sum_{k_1=0}^{2l_1}\cdots\sum_{k_n=0}^{2l_n}(2\pi\,\mathrm{i})^n L^\sharp_{l_1\ldots,l_n}(k_1{+}1, \ldots, k_n{+}1)\cdot a_{2l_1}\frac{(\operatorname{ad} e_-)^{k_1}(\tilde\delta_{2l_1})}{k_1!}\cdots a_{2l_n}\frac{(\operatorname{ad} e_-)^{k_n}(\tilde\delta_{2l_n})}{k_n!}.$$

Then:

(1) $\tilde\theta \in \widehat{U(\tilde{\mathfrak{b}}_3^+)}$ *is group-like (the hat means degree completion);*

(2) let

$$\tilde\psi := \sum_{n\geq 0}\sum_{l_1,\dots,l_n>0}\sum_{k_1=0}^{2l_1}\cdots\sum_{k_n=0}^{2l_n} \frac{1}{(k_1+1)\cdots(k_1+\cdots+k_n+n)}(2\pi\,\mathrm{i})^{2(l_1+\cdots+l_n+n)}\cdot$$
$$\cdot a_{2l_1}\frac{(\mathrm{ad}\,e_-)^{k_1}(\tilde\delta_{2l_1})}{k_1!}\cdots a_{2l_n}\frac{(\mathrm{ad}\,e_-)^{k_n}(\tilde\delta_{2l_n})}{k_n!}.$$

then

$$\tilde\theta\cdot\alpha(\tilde\theta) = (\tilde\theta\tilde\psi)\cdot\beta(\tilde\theta\tilde\psi)\cdot\beta^2(\tilde\theta\tilde\psi) = 1, \tag{3}$$

where $\alpha := \left(\begin{smallmatrix}0 & -1\\ 1 & 0\end{smallmatrix}\right)$, $\beta = \left(\begin{smallmatrix}1 & -1\\ 1 & 0\end{smallmatrix}\right)$ *(*$\in \mathrm{PSL}_2(\mathbb{C})$*);*

(3) let $\theta := \mathrm{im}(\tilde\theta \in U(\tilde{\mathfrak{b}}_3) \to \mathrm{End}(U\mathfrak{t}_{1,2}^{\mathbb{C}}))$, *then*

$$e^{\frac{1}{4}(2\pi\,\mathrm{i})^2\delta_0}\left(\begin{smallmatrix}0 & -1\\ 1 & 0\end{smallmatrix}\right)\theta : (A_{KZ}, B_{KZ}) \mapsto (A_{KZ}B_{KZ}A_{KZ}^{-1}, A_{KZ}^{-1}),$$

where $((x,y) = (x_2^+, x_2^-))$

$$A_{KZ} = \Phi_{KZ}\Big(\frac{\mathrm{ad}\,2\pi\,\mathrm{i}\,x}{e^{\mathrm{ad}\,2\pi\,\mathrm{i}\,x}-1}(y), -2\pi\,\mathrm{i}\,t\Big)e^{\frac{\mathrm{ad}\,2\pi\,\mathrm{i}\,x}{e^{\mathrm{ad}\,2\pi\,\mathrm{i}\,x}-1}(2\pi\,\mathrm{i}\,y)}\Phi_{KZ}\Big(\frac{\mathrm{ad}\,2\pi\,\mathrm{i}\,x}{e^{\mathrm{ad}\,2\pi\,\mathrm{i}\,x}-1}(y), -2\pi\,\mathrm{i}\,t\Big)^{-1},$$

$$B_{KZ} = e^{-\frac{1}{2}(2\pi\,\mathrm{i})^2 t}\Phi_{KZ}\Big(\frac{\mathrm{ad}\,2\pi\,\mathrm{i}\,x}{e^{-\mathrm{ad}\,2\pi\,\mathrm{i}\,x}-1}(y), -2\pi\,\mathrm{i}\,t\Big)e^{2\pi\,\mathrm{i}\,x}\Phi_{KZ}\Big(\frac{\mathrm{ad}\,2\pi\,\mathrm{i}\,x}{e^{\mathrm{ad}\,2\pi\,\mathrm{i}\,x}-1}(y), -2\pi\,\mathrm{i}\,t\Big)^{-1}.$$

Remark 10. 1) According to [Za], $L_l^\sharp(1) = -L_l^\sharp(2l+1) = (\text{nonzero rational})\times\zeta(2l+1)$ and for $1 < k < 2l+1$, $L_l^\sharp(k) \in (2\pi\,\mathrm{i})^{2l+1}\mathbb{Q}$, where the rational numbers are related to Bernoulli numbers.

2) It can be shown that $\tilde{\mathfrak{b}}_3^+$ is freely generated by the $(\mathrm{ad}\,e_-)^k(\tilde\delta_{2l})$, $l \geq 0$, $k = 0,\dots,2l$. It follows that (1) can be translated as a shuffle relation

$$L^\sharp_{l_1,\dots,l_p}(k_1{+}1,\dots,k_p{+}1)L^\sharp_{l_{p+1},\dots,l_n}(k_{p+1}{+}1,\dots,k_n{+}1) = \sum_{\sigma\in S_{p,n-p}} L^\sharp_{l_{\sigma(1)},\dots,l_{\sigma(n)}}(k_{\sigma(1)}{+}1,\dots,k_{\sigma(n)}{+}1). \tag{4}$$

Similarly, (2) and (3) can be expressed as a family of polynomial relations on $2\pi\,\mathrm{i}$ and the $L^\sharp_{l_1,\dots,l_n}(k_1,\dots,k_n)$ of degree 2 (resp., 3) with rational coefficients, which may be viewed as homogeneous if $2\pi\,\mathrm{i}$ has weight 1 and $L^\sharp_{l_1,\dots,l_n}(k_1,\dots,k_n)$ has weight $2(l_1+\cdots+l_n)+n$.

3) It can be shown that $\psi := \mathrm{im}(\tilde\psi \in U(\tilde{\mathfrak{b}}_3) \to \mathrm{End}(U\mathfrak{t}^{\mathbb{C}}_{1,2}))$ is such that $e^{-\frac{1}{12}(2\pi\,\mathrm{i})^2\delta_0}\tilde\psi\left(\begin{smallmatrix}1 & 1\\ 0 & 1\end{smallmatrix}\right)$: $(A_{KZ}, B_{KZ}) \mapsto (A_{KZ}, B_{KZ}A_{KZ})$.

Define $\mathrm{Z}_n := \mathrm{Span}_{\mathbb{Q}}\{(2\pi\,\mathrm{i})^a\zeta(k_1,\dots,k_p) | a + k_1 + \cdots + k_p = n\} \subset \mathbb{C}$; then $\mathbf{k}_{MZV} := \sum_{n\geq 0}\mathrm{Z}_n$ is a subring of $\mathbb{C}$, and $\mathrm{Z}_n\,\mathrm{Z}_m \subset \mathrm{Z}_{n+m}$. Recall that $\mathfrak{t}_{1,2}$ is defined over $\mathbb{Q}$; we equip it with the degree $\deg x = (1,0)$, $\deg y = (0,1)$.

Relation (4) then gives:

Proposition 11. *Any* $\ell \in (\mathrm{Der}\,\mathfrak{t}_{1,2})^*_{l,k}$ *yields a relation*

$$\sum_{\substack{(n,\underline{k},\underline{l})|\\ k_1+\cdots+k_n+n=k,\\ l_1+\cdots+l_n+n=(k+l)/2}} r^\ell_{\underline{l},\underline{k}}(2\pi\,\mathrm{i})^n L^\sharp_{l_1,\dots,l_n}(k_1+1,\cdots,k_n+1) \in \mathrm{Z}_{k+l},$$

where the $r^\ell_{\underline{l},\underline{k}}$ *are in* $\mathbb{Q}$*.*

Relation (4) does not allow to compute the $L^\sharp_{l_1,\dots,l_n}(k_1,\cdots,k_n)$ in terms of MZVs, as there are nontrivial relations in $\mathfrak{b}_3$ between the $(\mathrm{ad}\,e_-)^k(\delta_{2l})$. For example, the space of quadratic Lie relations between the δ_{2l} of a given weight (where $\mathrm{wt}(\delta_{2l}) = 2l+1$) is isomorphic to the space of cusp forms for $\mathrm{SL}_2(\mathbb{Z})$ of the same weight, and the quadratic relation of lowest weight says that $[\delta_2, \delta_8]$ and $[\delta_4, \delta_6]$ are proportional.

In particular, we do not know whether $L^\sharp_{l_1,\ldots,l_n}(k_1, \cdots, k_n) \in \mathrm{Z}_{2(l_1+\cdots+l_n)+n}$ in general.

3. ELLIPTIC STRUCTURES OVER BRAIDED MONOIDAL CATEGORIES

We now develop the theory of elliptic associators, starting with elliptic structures over BMCs.

3.1. **Definition.** Let $(\mathcal{C}_0, \otimes, \beta_{-,-}, a_{-,-,-}, \mathbf{1})$ be a braided monoidal category (see e.g. [Ka]). Here $\otimes : \mathcal{C}_0 \times \mathcal{C}_0$ is the tensor product, $\beta_{X,Y} : X \otimes Y \to Y \otimes X$ and $a_{X,Y,Z} : (X \otimes Y) \otimes Z \to X \otimes (Y \otimes Z)$ are the braiding and associativity isomorphisms and $\mathbf{1}$ is the unit object. They satisfy in particular the pentagon and hexagon identities

$$a_{X,Y,Z\otimes T} a_{X\otimes Y,Z,T} = (\mathrm{id}_X \otimes a_{Y,Z,T}) a_{X,Y\otimes Z,T} (a_{X,Y,Z} \otimes \mathrm{id}_T),$$

$$(\mathrm{id}_Y \otimes \beta^\pm_{X,Z}) a_{Y,X,Z} (\beta^\pm_{X,Y} \otimes \mathrm{id}_Z) = a_{Y,Z,X} \beta^\pm_{X,Y\otimes Z} a_{X,Y,Z},$$

where $\beta^+_{X,Y} = \beta_{X,Y}$, $\beta^-_{X,Y} = \beta^{-1}_{Y,X}$.

Definition 12. *An elliptic structure over $\mathcal{C}_0$ is a set $(\mathcal{C}, F, A^+_{-,-}, A^-_{-,-})$, where $\mathcal{C}$ is a category, $F : \mathcal{C}_0 \to \mathcal{C}$ is a functor inducing a bijection*[1] $\mathrm{Ob}\,\mathcal{C}_0 \simeq \mathrm{Ob}\,\mathcal{C}$*, and $A^\pm_{-,-}$ are natural*[2] *assignments $(\mathrm{Ob}\,\mathcal{C}_0)^2 \ni (X, Y) \mapsto A^\pm_{X,Y} \in \mathrm{Aut}_\mathcal{C}(X \otimes Y)$, such that:*

$$\alpha^\pm_{Z,X,Y} \alpha^\pm_{Y,Z,X} \alpha^\pm_{X,Y,Z} = \mathrm{id}_{(X\otimes Y)\otimes Z}, \tag{5}$$

where $\alpha^\pm_{X,Y,Z} = F(\beta^\pm_{X,Y\otimes Z}) A^\pm_{X,Y\otimes Z} F(a_{X,Y,Z})$,

$$\begin{aligned} F(\beta_{Y,X}\beta_{X,Y} \otimes \mathrm{id}_Z) = \quad & \big(F(a^{-1}_{X,Y,Z}) A^-_{X,Y\otimes Z} F(a_{X,Y,Z}), \\ & F((\beta^{-1}_{X,Y} \otimes \mathrm{id}_Z) a^{-1}_{Y,X,Z}) (A^+_{Y,X\otimes Z})^{-1} F(a_{Y,X,Z}(\beta^{-1}_{Y,X} \otimes \mathrm{id}_Z))\big), \end{aligned} \tag{6}$$

(identities[3] *in $\mathrm{Aut}_\mathcal{C}((X \otimes Y) \otimes Z)$), for any $X, Y, Z \in \mathrm{Ob}\,\mathcal{C}_0$, and*

$$A^\pm_{X,\mathbf{1}} = \mathrm{id}_{X\otimes\mathbf{1}} \quad \textit{for any } X \in \mathrm{Ob}\,\mathcal{C}_0. \tag{7}$$

A morphism $(\mathcal{C}_0, \mathcal{C}, F, A^\pm_{-,-}) \to (\mathcal{C}'_0, \mathcal{C}', F', A'^\pm_{-,-})$ is then the data of a tensor functor $\mathcal{C}_0 \overset{\varphi_0}{\to} \mathcal{C}'_0$ and a functor $\mathcal{C} \overset{\varphi}{\to} \mathcal{C}'$, such that

$$\begin{tikzcd} \mathcal{C}_0 \arrow[r] \arrow[d] & \mathcal{C}'_0 \arrow[d] \\ \mathcal{C} \arrow[r] & \mathcal{C}' \end{tikzcd}$$

commutes, and $\varphi(A^\pm_{X,Y}) = A'^\pm_{\varphi_0(X),\varphi_0(Y)}$.

Remark 13. Taking (7) into account, axiom (5) can be replaced by

$$A^\pm_{X\otimes Y,Z} = F((\beta^\pm_{Y,X} \otimes \mathrm{id}_Z) a^{-1}_{Y,X,Z}) A^\pm_{Y,X\otimes Z} F(a_{Y,X,Z}(\beta^\pm_{X,Y} \otimes \mathrm{id}_Z) a^{-1}_{X,Y,Z}) A^\pm_{X,Y\otimes Z} F(a_{X,Y,Z}).$$

The axioms (5)-(7) also imply $F(\beta^\pm_{Y,X}) A^\pm_{Y,X} F(\beta^\pm_{X,Y}) A^\pm_{X,Y} = \mathrm{id}_{X\otimes Y}$ and $F(\beta_{Y,X}\beta_{X,Y}) = (A^-_{X,Y}, A^+_{X,Y})$.

---

[1] For $\mathcal{C}$ a category, $\mathrm{Ob}\,\mathcal{C}$ is its class of objects; for $X, Y \in \mathrm{Ob}\,\mathcal{C}$, $\mathrm{Iso}_\mathcal{C}(X, Y) \subset \mathcal{C}(X, Y)$ is the set of (iso)morphisms $X \to Y$, $\mathrm{Aut}_\mathcal{C}(X) = \mathrm{Iso}_\mathcal{C}(X, X)$.

[2] Natural means that if $\varphi \in \mathcal{C}_0(X, X')$, $\psi \in \mathcal{C}_0(Y, Y')$, then $A^\pm_{X',Y'} F(\varphi \otimes \psi) = F(\varphi \otimes \psi) A^\pm_{X,Y}$

[3] In (6) and later, we set $(g, h) := ghg^{-1}h^{-1}$.

3.2. **Relation with elliptic braid groups.** The reduced elliptic braid group on n strands $B_{1,n}$ can be presented by generators σ_i $(i = 1, ..., n-1)$, $X_1^{\pm}$, and relations

$$(\sigma_1^{\pm 1} X_1^{\pm})^2 = (X_1^{\pm}\sigma_1^{\pm 1})^2, \quad (X_1^{\pm}, \sigma_i) = 1 \text{ for } i = 2, ..., n-1, \quad (X_1^{-}, (X_2^{+})^{-1}) = \sigma_1^2, \quad X_1^{\pm}...X_n^{\pm} = 1,$$

$$(\sigma_i, \sigma_j) = 1 \text{ for } |i-j| > 1, \quad \sigma_i\sigma_{i+1}\sigma_i = \sigma_{i+1}\sigma_i\sigma_{i+1} \text{ for } i = 1, ..., n-2, \tag{8}$$

where $X_{i+1}^{\pm} = \sigma_i^{\pm 1} X_i^{\pm} \sigma_i^{\pm 1}$ for $i = 1, ..., n-1$ ([Bi]).

The braid group B_n on n strands is presented by generators σ_i, $i = 1, ..., n-1$ and the Artin relations (8). We have a morphism $B_n \to B_{1,n}$, $\sigma_i \mapsto \sigma_i$. If $\mathcal{C}_0$ is a braided monoidal category and $X \in \mathrm{Ob}\,\mathcal{C}_0$, then there is a unique group morphism $\varphi_0 : B_n \to \mathrm{Aut}_{\mathcal{C}}(X^{\otimes n})$ ($X^{\otimes n}$ is defined by right parenthesization, so $X^{\otimes n} = X \otimes X^{\otimes n-1}$), such that

$$\sigma_i \mapsto a_i((\mathrm{id}_{X^{\otimes i-1}} \otimes \beta_{X,X}) \otimes \mathrm{id}_{X^{\otimes n-i-1}})a_i^{-1},$$

where $a_i : (X^{\otimes i-1} \otimes X^{\otimes 2}) \otimes X^{\otimes n-i-1} \to X^{\otimes n}$ is the associativity constraint.

Proposition 14. *If $(\mathcal{C}, F, A^{\pm}_{-,-})$ is an elliptic structure over $\mathcal{C}_0$ and $X \in \mathrm{Ob}\,\mathcal{C}_0$, then there is a unique group morphism $B_{1,n} \to \mathrm{Aut}_{\mathcal{C}}(X^{\otimes n})$, such that*

$$X_1^{\pm} \mapsto A^{\pm}_{X,X^{\otimes n-1}}, \quad \sigma_i \mapsto F(\varphi_0(\sigma_i)).$$

Proof. Let us check that $(\sigma_1 X_1^+)^2 = (X_1^+\sigma_1)^2$, i.e., $(X_1^+, X_2^+) = 1$ is preserved (for simplicity, we omit the associativity constraints). By naturality, $(\beta_{X,Y} \otimes \mathrm{id}_Z)A^+_{X\otimes Y,Z} = A^+_{Y\otimes X,Z}(\beta_{X,Y} \otimes \mathrm{id}_Z)$. Plugging in this equality the relations $A^+_{X\otimes Y,Z} = (\beta_{Y,X} \otimes \mathrm{id}_Z)A^+_{Y,X\otimes Z}(\beta_{X,Y} \otimes \mathrm{id}_Z)A^+_{X,Y\otimes Z}$ and $A^+_{Y\otimes X,Z} = (\beta_{X,Y} \otimes \mathrm{id}_Z)A^+_{X,Y\otimes Z}(\beta_{Y,X} \otimes \mathrm{id}_Z)A^+_{Y,X\otimes Z}$, we obtain

$$(A^+_{X,Y\otimes Z}, (\beta_{Y,X} \otimes \mathrm{id}_Z)A^+_{Y,X\otimes Z}(\beta_{X,Y} \otimes \mathrm{id}_Z)) = 1;$$

if we set $Y := X$, $Z := X^{\otimes n-2}$, this says that $(X_1^+, X_2^+) = 1$ is preserved. Similarly, one proves that (5) with sign $-$ implies that $(\sigma_1^{-1}X_1^-)^2 = (X_1^-\sigma_1^{-1})^2 = 1$ is preserved. (6) immediately implies that $(X_1^-, (X_1^+)^{-1}) = \sigma_1^2$ is preserved. The naturality assumption implies that $(X_1^{\pm}, \sigma_i) = 1$ $(i > 1)$ is preserved. The image of $X_1^{\pm}...X_n^{\pm}$ is $A^{\pm}_{X^{\otimes n},\mathbf{1}}$, which is $\mathrm{id}_{X^{\otimes n}}$ by (7). Finally, as $B_n \to \mathrm{Aut}_{\mathcal{C}_0}(X^{\otimes n})$ is a group morphism, the Artin relations are preserved. ☐

3.3. **Universal elliptic structures.** Let $\mathcal{C}_0 := \mathbf{PaB}$ be the braided monoidal category of parenthesized braids (see [JS, Ba]). Recall that $\mathrm{Ob}\,\mathcal{C}_0 = \sqcup_{n\geq 0}\mathrm{Par}_n$, where $\mathrm{Par}_n = \{$parenthesizations of the word $\bullet\ldots\bullet$ of length $n\}$, so $\mathrm{Par}_0 = \{\mathbf{1}\}$, $\mathrm{Par}_1 = \{\bullet\}$, $\mathrm{Par}_2 = \{\bullet\bullet\}$, $\mathrm{Par}_3 = \{(\bullet\bullet)\bullet, \bullet(\bullet\bullet)\}$, etc. For $O, O' \in \mathrm{Ob}\,\mathcal{C}_0$, we set $|O| :=$ the integer such that $O \in \mathrm{Par}_{|O|}$, and $\mathcal{C}_0(O,O') := \begin{cases} B_{|O|} & \text{if } |O| = |O'| \\ \emptyset & \text{otherwise} \end{cases}$. The composition is the product in $B_{|O|}$. The tensor product is defined at the level of objects, as the juxtaposition, and at the level of morphisms, by the group morphism $B_n \times B_{n'} \to B_{n+n'}$, $(\sigma_i, 1) \mapsto \sigma_i$, $(1, \sigma_j) \mapsto \sigma_{n+j}$. We set $a_{O,O',O''} := 1 \in B_{|O|+|O'|+|O''|} = \mathcal{C}_0((O \otimes O') \otimes O'', O \otimes (O' \otimes O''))$ and $\beta_{O,O'} := \sigma_{|O|,|O'|} \in B_{|O|+|O'|} = \mathcal{C}_0(O \otimes O', O' \otimes O)$, where $\sigma_{n,n'} := (\sigma_n...\sigma_1)(\sigma_{n+1}...\sigma_2)...(\sigma_{n+n'-1}...\sigma_{n'}) \in B_{n+n'}$.

Let now $\mathcal{C} := \mathbf{PaB}_{ell}$ be the category with $\mathrm{Ob}\,\mathcal{C} := \mathrm{Ob}\,\mathcal{C}_0$, $\mathcal{C}(O,O') := \begin{cases} B_{1,|O|} & \text{if } |O| = |O'| \\ \emptyset & \text{otherwise} \end{cases}$ and whose product is the composition in $B_{1,|O|}$. Let $F : \mathcal{C}_0 \to \mathcal{C}$ be the functor induced by the identity at the level of objects, and by $B_n \to B_{1,n}$, $\sigma_i \mapsto \sigma_i$ at the level of morphisms. For $O, O' \in \mathrm{Ob}\,\mathcal{C}_0$, set $A^{\pm}_{O,O'} := X_1^{\pm}...X_{|O|}^{\pm} \in B_{1,|O|+|O'|}$. Then $(\mathbf{PaB}_{ell}, F, A^{\pm}_{-,-})$ is an elliptic structure over $\mathbf{PaB}$.

The pair $(\mathbf{PaB}, \bullet)$ has the following universal property: for any pair $(\tilde{\mathcal{C}}_0, M)$, where $\tilde{\mathcal{C}}_0$ is a braided monoidal category and $M \in \mathrm{Ob}\,\tilde{\mathcal{C}}_0$, there exists a unique tensor functor $\varphi_0 : \mathbf{PaB} \to \tilde{\mathcal{C}}_0$, such that $F(\bullet) = M$. Proposition 14 immediately implies that this property extends as follows.

Proposition 15. *If now $\tilde{\mathcal{C}}$ is an elliptic structure over $\tilde{\mathcal{C}}_0$, then there exists a unique morphism $(\mathbf{PaB}, \mathbf{PaB}_{ell}) \to (\tilde{\mathcal{C}}_0, \tilde{\mathcal{C}})$, extending φ_0.*

3.4. **Generalization: elliptic structures over** $(\mathcal{C}_0, \mathcal{D})$**.** Let $\mathcal{D}$ be a category and let $\mathcal{C}_0$ be a braided monoidal category.

Definition 16. *A $(\mathcal{C}_0, \mathcal{D})$-elliptic structure is the data of:*
(a) a category $\mathcal{C}$ and a functor $F : \mathcal{C}_0 \times \mathcal{D} \to \mathcal{C}$,
(b) natural[4] *assignments $(\mathrm{Ob}\,\mathcal{C}_0)^2 \times \mathrm{Ob}\,\mathcal{D} \ni (X, Y, M) \mapsto A^{\pm}_{X,Y,M} \in \mathrm{Aut}_{\mathcal{C}}(F(X \otimes Y, M))$, satisfying the analogues of (5)-(6) with $A^{\pm}_{X\otimes Y,Z}, \ldots$ replaced by $A^{\pm}_{X\otimes Y,Z,M}, \ldots$ and the various $F(\varphi)$ by $F(\varphi, \mathrm{id}_M)$, as well as $A^{\pm}_{X,\mathbf{1},M} = \mathrm{id}_{F(X,M)}$.*

If $M \in \mathrm{Ob}\,\mathcal{D}$, then we have a functor $F(M, -) : \mathcal{C}_0 \to \mathcal{C}$, and[5] $(F(M, -)^*\mathcal{C}, \mathcal{C}_0 \to F(M, -)^*\mathcal{C}, A^{\pm}_{-,-,M})$ is an elliptic structure over $\mathcal{C}_0$. In particular, we obtain morphisms $B_{1,n} \to \mathrm{Aut}_{\mathcal{C}}(F(X^{\otimes n}, M))$ for any $X \in \mathrm{Ob}\,\mathcal{C}_0$.

Example 17. If $\mathcal{D}$ is trivial and F induced a bijection between sets of objects, we recover the notion of an elliptic structure over $\mathcal{C}_0$.

Example 18. Let H be a QTQBA (quasi-triangular quasi-bialgebra) over a field $\mathbf{k}$ and let (D, A, B) be an elliptic structure over H (see [CEE], Def. 5.6). Let $\mathcal{D} := \{D\text{-modules}\}$, $\mathcal{C}_0 := \{H\text{-modules}\}$, $\mathcal{C} := \mathrm{Vect}$ (the category of vector spaces), $F : \mathcal{C}_0 \times \mathcal{D} \to \mathcal{C}$ be given by $F(X, M) := (M \otimes X)^H$, $A^{\pm}_{X,Y,M} := (\pi_M \otimes \pi_X \otimes \pi_Y)(A^{\pm})$ (we set $A^+ = A$, $A^- = B$, then $A^{\pm} \in D \otimes A^{\otimes 2}$).

4. The elliptic Grothendieck-Teichmüller group

4.1. **Reminders about $\underline{\mathrm{GT}}$ and its variants.** According to [Dr], $\underline{\mathrm{GT}}$ is the set of pairs $(\lambda, f) \in (1 + 2\mathbb{Z}) \times F_2$, such that

$$f(X_3, X_1)X_3^m f(X_2, X_3)X_2^m f(X_1, X_2)X_1^m = 1, \quad m = \frac{\lambda - 1}{2}, \quad X_1X_2X_3 = 1,$$

$$f(Y, X) = f(X, Y)^{-1}, \quad \partial_3(f)\partial_1(f) = \partial_0(f)\partial_2(f)\partial_4(f),$$

where[6] $\partial_i : F_2 \subset P_3 \to P_4$ are simplicial morphisms. It is equipped with a semigroup structure with $(\lambda, f)(\lambda', f') = (\lambda'', f'')$, with $\lambda'' := \lambda\lambda'$, $f''(X, Y) := f(f'(X, Y)X^{\lambda'} f'(X, Y)^{-1}, Y^{\lambda'})f'(X, Y)$

One defines similarly semigroups $\widehat{\underline{\mathrm{GT}}}$, $\underline{\mathrm{GT}}_l$, $\underline{\mathrm{GT}}(\mathbf{k})$ by replacing in the above definition $(\mathbb{Z}, F_2)$ by their profinite, pro-l, $\mathbf{k}$-prounipotent versions (where $\mathbf{k}$ is a $\mathbb{Q}$-ring). We then have morphisms $\underline{\mathrm{GT}} \hookrightarrow \widehat{\underline{\mathrm{GT}}} \to \underline{\mathrm{GT}}_l \hookrightarrow \underline{\mathrm{GT}}(\mathbb{Q}_l)$ and $\underline{\mathrm{GT}} \to \underline{\mathrm{GT}}(\mathbf{k})$ for any $\mathbf{k}$.

$\underline{\mathrm{GT}}$ acts on {braided monoidal categories (BMCs)} by $(\lambda, f)(\mathcal{C}_0, \beta, a) := (\mathcal{C}_0, \beta', a')$, where $\beta'_{X,Y} := \beta_{X,Y}(\beta_{Y,X}\beta_{X,Y})^m$, $a'_{X,Y,Z} := a_{X,Y,Z}f(\beta_{YX}\beta_{XY} \otimes \mathrm{id}_Z, a^{-1}_{X,Y,Z}(\mathrm{id}_X \otimes \beta_{ZY}\beta_{YZ})a_{X,Y,Z})$. Similarly, $\widehat{\underline{\mathrm{GT}}}$ (resp., $\underline{\mathrm{GT}}_l$, $\underline{\mathrm{GT}}(\mathbf{k})$) act on {BMCs $\mathcal{C}_0$ such that $\mathrm{Aut}_{\mathcal{C}_0}(X)$ is finite for any $X \in \mathrm{Ob}\,\mathcal{C}_0$} (resp., such that the image of $P_n \to \mathrm{Aut}_{\mathcal{C}_0}(X_1 \otimes \cdots \otimes X_n)$ is an l-group, is contained in a unipotent group).

---

[4] Natural means that if $f \in \mathcal{C}_0(X, X')$, $g \in \mathcal{C}_0(Y, Y')$, $\varphi \in \mathcal{D}(M, M')$, then $F(f \otimes g, \varphi)A^{\pm}_{X,Y,M} = A^{\pm}_{X',Y',M'}F(f \otimes g, \varphi)$.

[5] If $F : \mathcal{A} \to \mathcal{B}$ is a functor, then we define $F^*\mathcal{B}$ as the category with $\mathrm{Ob}\,F^*\mathcal{B} := \mathrm{Ob}\,\mathcal{A}$ and $(F^*B)(X, Y) := \mathcal{B}(F(X), F(Y))$; then F induces a functor $\mathcal{A} \to F^*\mathcal{B}$, which is the identity on objects.

[6] $P_n = \mathrm{Ker}(B_n \to S_n, \sigma_i \mapsto (i, i + 1))$ is the pure braid group on n strands.

4.2. **The semigroup $\underline{\mathrm{GT}}_{ell}$ and its variants.** Let us define $\underline{\mathrm{GT}}_{ell}$ as the set of all $(\lambda, f, g_\pm)$, where $(\lambda, f) \in \underline{\mathrm{GT}}$, $g_\pm \in F_2$ are such that

$$(\sigma_2^{\pm 1}\sigma_1^{\pm 1}(\sigma_1\sigma_2^2\sigma_1)^{\pm m} g_\pm(X_1^+, X_1^-) f(\sigma_1^2, \sigma_2^2))^3 = 1, \tag{9}$$

$$u^2 = (g_-, u^{-1} g_+^{-1} u^{-1}) \tag{10}$$

(identities in $B_{1,3}$) where $u = f(\sigma_1^2, \sigma_2^2)\sigma_1^\lambda f(\sigma_1^2, \sigma_2^2)^{-1}$, $g_\pm = g_\pm(X_1^+, X_1^-)$. Note that these conditions imply the identities

$$(\sigma_1^{\pm\lambda} g_\pm(X_1^+, X_1^-))^2 = 1, \quad \sigma_1^{2\lambda} = (g_-(X_1^+, X_1^-), g_+(X_1^+, X_1^-)) \tag{11}$$

in $B_{1,2}$.

We set $(\lambda, f, g_\pm)(\lambda', f', g'_\pm) := (\lambda'', f'', g''_\pm)$, where

$$g''_\pm(X, Y) = g_\pm(g'_+(X, Y), g'_-(X, Y)).$$

Proposition 19. *This defines a semigroup structure on $\underline{\mathrm{GT}}_{ell}$. We have a semigroup inclusion $\underline{\mathrm{GT}}_{ell} \subset \underline{\mathrm{GT}} \times \mathrm{End}(F_2)^{op}$, $(\lambda, f, g_\pm) \mapsto ((\lambda, f), \theta_{g_\pm})$, where $\theta_{g_\pm} = (X \mapsto g_+(X, Y), Y \mapsto g_-(X, Y))$.*

Proof. We first prove:

Lemma 20. *If $(\lambda, f, g_\pm) \in \underline{\mathrm{GT}}_{ell}$, then there is a unique endomorphism of $B_{1,3}$, such that*

$$\sigma_1 \mapsto \tilde\sigma_1 := f(\sigma_1^2, \sigma_2^2)\sigma_1^\lambda f(\sigma_1^2, \sigma_2^2)^{-1}, \quad \sigma_2 \mapsto \tilde\sigma_2 := \sigma_2^\lambda, \quad X_1^\pm \mapsto g_\pm(X_1^+, X_1^-).$$

For any $\lambda' \in 2\mathbb{Z} + 1$, we then have

$$f(\sigma_1^2, \sigma_2^2)\sigma_2^{\pm 1}\sigma_1^{\pm 1}(\sigma_1\sigma_2^2\sigma_1)^{\pm\frac{\lambda\lambda'-1}{2}} = \tilde\sigma_2^{\pm 1}\tilde\sigma_1^{\pm 1}(\tilde\sigma_1\tilde\sigma_2^2\tilde\sigma_1)^{\pm\frac{\lambda'-1}{2}}. \tag{12}$$

Proof. Recall that we have an elliptic structure $(\mathbf{PaB}, \mathbf{PaB}_{ell}, F, A^\pm_{-,-})$. Applying $(\lambda, f, g_\pm)$ to it, we get an elliptic structure $(\widetilde{\mathbf{PaB}}, \widetilde{\mathbf{PaB}}^{ell}, F, \tilde A^\pm_{-,-})$. The morphism $B_{1,3} \to \mathrm{Aut}_{\widetilde{\mathbf{PaB}}^{ell}}(\bullet(\bullet\bullet)) \simeq B_{1,3}$ is then the above endomorphism of $B_{1,3}$.

We now prove (12). The hexagon identity implies

$$(\sigma_2^2)^m f(\sigma_1^2, \sigma_2^2)(\sigma_1^2)^m f((\sigma_1^2\sigma_2^2)^{-1}, \sigma_1^2)(\sigma_1^2\sigma_2^2)^{-m} f(\sigma_2^2, (\sigma_1^2\sigma_2^2)^{-1}) = 1.$$

Now since $(\sigma_1^2\sigma_2^2)^{-1} \equiv \sigma_1\sigma_2^2\sigma_1^{-1} \equiv \sigma_2^{-1}\sigma_1^2\sigma_2$ mod $Z(B_3)$, since $f(a, b) = f(a', b')$ for any group G and any $a, a', b, b' \in G$ with $a \equiv a'$, $b \equiv b'$ mod $Z(G)$, and by the duality identity, this is rewritten $(\sigma_2^2)^m f(\sigma_1^2, \sigma_2^2)\sigma_1^{2m+1} f(\sigma_1^2, \sigma_2^2)^{-1}\sigma_1^{-1}(\sigma_1^2\sigma_2^2)^{-m}\sigma_2^{-1} f(\sigma_1^2, \sigma_2^2)^{-1}\sigma_2 = 1$, which yields (12) with $(\pm, \lambda') = (+, 1)$. (12) with $\pm = +$ then follows from $\tilde\sigma_1\tilde\sigma_2^2\tilde\sigma_1 = (\sigma_1\sigma_2^2\sigma_1)^\lambda$, which is proved as follows: $\tilde\sigma_1\tilde\sigma_2^2\tilde\sigma_1 = \tilde Z\tilde\sigma_2^{-2} = Z^\lambda\sigma_2^{-2\lambda} = (Z\sigma_2^{-2})^\lambda = (\sigma_1\sigma_2^2\sigma_1)^\lambda$, where $Z = (\sigma_1\sigma_2^2)^2$ is a generator of $Z(PB_3) \simeq \mathbb{Z}$, while $\tilde Z$ is its image by the endomorphism $\sigma_i \mapsto \tilde\sigma_i$ of B_3; $\tilde Z = Z^\lambda$ follows from the hexagon identity. Now each side of (12) with $\pm = -$ identifies with the same side of (12) with $\pm = +$ and λ' replaced by $-\lambda'$. This implies (12) with $\pm = -$. □

End of proof of Proposition 19. It suffices to prove that $(\lambda'', f'', g''_\pm) \in \underline{\mathrm{GT}}_{ell}$. It satisfies (10) because of the first part of Lemma 20 (existence of an endomorphism of $B_{1,3}$), and (9) because of (12) and of the existence of the same endomorphism. □

Then $\underline{\mathrm{GT}}_{ell}$ acts on $\{(\mathcal{C}_0, \mathcal{C}, F, A^\pm_{-,-}) | \mathcal{C}_0$ is a BMC, $(\mathcal{C}, F, A^\pm_{-,-})$ is an elliptic structure over it$\}$ as $(\lambda, f, g_\pm)(\mathcal{C}_0, \mathcal{C}, F, A^\pm_{-,-}) := (\mathcal{C}'_0, \mathcal{C}, F, A'^\pm_{-,-})$, where $\mathcal{C}'_0 = (\lambda, f)\mathcal{C}_0$, $A'^\pm_{X,Y} = g_\pm(A^+_{X,Y}, A^-_{X,Y})$.

As before, we define semigroups $\widehat{\underline{\mathrm{GT}}}_{ell}$, $\underline{\mathrm{GT}}_l^{ell}$, $\underline{\mathrm{GT}}(\mathbf{k})$ by replacing in the definition of $\underline{\mathrm{GT}}_{ell}$, $(\underline{\mathrm{GT}}, F_2)$ by[7] $(\widehat{\underline{\mathrm{GT}}}, \widehat{F}_2)$, $(\underline{\mathrm{GT}}_l, (F_2)_l)$, $(\underline{\mathrm{GT}}(\mathbf{k}), F_2(\mathbf{k}))$. They act on the sets of pairs $(\mathcal{C}_0, \mathcal{C})$, such that $\mathcal{C}_0$ satisfies the same conditions as above, together with: $\mathrm{Aut}_{\mathcal{C}}(X)$ is finite for any $X \in \mathrm{Ob}\,\mathcal{C}_0$ (resp., the image of $P_{1,n} \to \mathrm{Aut}_{\mathcal{C}}(X_1 \otimes \cdots \otimes X_n)$ is an l-group, is contained in a unipotent group). We have morphisms $\underline{\mathrm{GT}}_{ell} \hookrightarrow \widehat{\underline{\mathrm{GT}}}_{ell} \to \underline{\mathrm{GT}}_l^{ell} \hookrightarrow \underline{\mathrm{GT}}_{ell}(\mathbb{Q}_l)$ and $\underline{\mathrm{GT}}_{ell} \to \underline{\mathrm{GT}}_{ell}(\mathbf{k})$ compatible with the similar 'non-elliptic' morphisms.

4.3. **Computation of $\underline{\mathrm{GT}}_{ell}$.** We first introduce a group $\widetilde{\mathrm{GL}_2\mathbb{Z}}$.

Proposition 21. *1) Let* $\widetilde{\mathrm{SL}_2\mathbb{Z}} := \langle \Theta, \Psi \rangle / ((\Theta^2, \Psi) = 1, (\Theta\Psi)^3 = \Theta^4)$*, then we have a central extension* $1 \to \mathbb{Z} \to \widetilde{\mathrm{SL}_2\mathbb{Z}} \to \mathrm{SL}_2(\mathbb{Z}) \to 1$*, where* $\widetilde{\mathrm{SL}_2\mathbb{Z}} \to \mathrm{SL}_2(\mathbb{Z})$ *is given by* $\Theta \mapsto \left(\begin{smallmatrix} 0 & 1 \\ -1 & 0 \end{smallmatrix}\right)$, $\Psi \mapsto \left(\begin{smallmatrix} 1 & 1 \\ 0 & 1 \end{smallmatrix}\right)$*. The center of* $\widetilde{\mathrm{SL}_2\mathbb{Z}}$ *is* $\langle \Theta^2 \rangle \simeq \mathbb{Z}$*, and* $\mathrm{Ker}(\widetilde{\mathrm{SL}_2\mathbb{Z}} \to \mathrm{SL}_2(\mathbb{Z})) = \langle \Theta^4 \rangle \simeq \mathbb{Z}$.

2) Let $\widetilde{\mathrm{GL}_2\mathbb{Z}} := \langle \Theta, \Psi, \varepsilon \rangle / ((\Psi, \Theta^2) = (\varepsilon\Psi\Theta)^2 = (\varepsilon\Theta)^2 = \varepsilon^2 = 1, (\Theta\Psi)^3 = \Theta^4)$*, then we have a (non-central) extension* $1 \to \mathbb{Z} \to \widetilde{\mathrm{GL}_2\mathbb{Z}} \to \mathrm{GL}_2(\mathbb{Z}) \to 1$*, where* $\widetilde{\mathrm{GL}_2\mathbb{Z}} \to \mathrm{GL}_2(\mathbb{Z})$ *extends* $\widetilde{\mathrm{SL}_2\mathbb{Z}} \to \mathrm{SL}_2(\mathbb{Z})$ *by* $\varepsilon \mapsto \left(\begin{smallmatrix} 0 & 1 \\ 1 & 0 \end{smallmatrix}\right)$*, and a morphism* $\widetilde{\mathrm{GL}_2\mathbb{Z}} \overset{\mathrm{det}}{\to} \mathbb{Z}/2$*, given by* $\Psi, \Theta, \varepsilon \mapsto 1, 1, -1$*, fitting in*

$$\begin{tikzcd}
1 \arrow[r] & \mathbb{Z} \arrow[r] \arrow[d] & \widetilde{\mathrm{SL}_2\mathbb{Z}} \arrow[r] \arrow[d] & \mathrm{SL}_2(\mathbb{Z}) \arrow[r] \arrow[d] & 1 \\
1 \arrow[r] & \mathbb{Z} \arrow[r] \arrow[d] & \widetilde{\mathrm{GL}_2\mathbb{Z}} \arrow[r] \arrow[d] & \mathrm{GL}_2(\mathbb{Z}) \arrow[r] \arrow[d] & 1 \\
0 \arrow[r] & 0 \arrow[r] & \mathbb{Z}/2 \arrow[r] & \mathbb{Z}/2 \arrow[r] & 0
\end{tikzcd}$$

The proof is straightforward. Note that we have an isomorphism $B_3 \simeq \widetilde{\mathrm{SL}_2\mathbb{Z}}$, given by $\sigma_1 \mapsto \Psi$, $\sigma_2 \mapsto \Theta\Psi\Theta^{-1}$.

Proposition 22. *1) There is a unique semigroup morphism* $\widetilde{\mathrm{GL}_2\mathbb{Z}} \to \underline{\mathrm{GT}}_{ell}$*, such that:*

$$\Psi \mapsto (\lambda, f, g_+, g_-) = (1, 1, g_+(X,Y) = X, g_-(X,Y) = YX),$$

$$\Theta \mapsto (\lambda, f, g_+, g_-) = (1, 1, g_+(X,Y) = Y^{-1}, g_-(X,Y) = YXY^{-1}),$$

$$\varepsilon \mapsto (\lambda, f, g_+, g_-) = (-1, 1, g_+(X,Y) = Y, g_-(X,Y) = X),$$

It fits in a commutative diagram

$$\begin{tikzcd}
\widetilde{\mathrm{GL}_2\mathbb{Z}} \arrow[r] \arrow[d] & \underline{\mathrm{GT}}_{ell} \arrow[d] \\
\mathbb{Z}/2 \arrow[r] & \underline{\mathrm{GT}}
\end{tikzcd}$$

2) The horizontal maps in this diagram are isomorphisms.

Proof. 1) is straightforward, using the above presentation of $\widetilde{\mathrm{GL}_2\mathbb{Z}}$. Let us prove 2). The bijectivity of $\mathbb{Z}/2 \to \underline{\mathrm{GT}}$ is proved in [Dr], Proposition 4.1. Set $\underline{R}_{ell} := \mathrm{Ker}(\underline{\mathrm{GT}}_{ell} \to \underline{\mathrm{GT}})$, then the commutativity of the above diagram implies that its upper map restricts to a morphism $\widetilde{\mathrm{SL}_2\mathbb{Z}} \to \underline{R}_{ell}$, and we need to prove that it is bijective. According to the second identity in (11), $\underline{R}_{ell} \subset \{(g_+, g_-) \in (F_2)^2 | (g_-(X,Y), g_+(X,Y)) = (Y, X)\}$. We now recall some results due to Nielsen.

---

[7] For G a group (other than GT, GT_{ell} or R_{ell}), $\widehat{G}$ is its profinite completion. If G is a free or pure (elliptic) braid group, G_l, $G(\mathbf{k})$ are its pro-l, $\mathbf{k}$-prounipotent completions. Here $G(-)$ is the prounipotent $\mathbb{Q}$-group scheme associated to G; it is characterized by $\mathrm{Hom}_{groups}(G, U(\mathbb{Q})) \simeq \mathrm{Hom}_{gp\ schemes}(G(-), U)$ for any unipotent group scheme U. If $G = B_n$ or $B_{1,n}$, then $G_l := P_l *_P G$, $G(\mathbf{k}) := P(\mathbf{k}) *_P G$, where $P = \mathrm{Ker}(G \to S_n)$.

Theorem 23. *([N]) 1) The morphism* $\mathrm{Out}(F_2) \to \mathrm{GL}_2(\mathbb{Z})$ *induced by abelianization is an isomorphism.*

2) $\mathrm{Im}(\mathrm{Aut}(F_2) \to (F_2)^2) = \{(g_+, g_-) \in (F_2)^2 | \exists k \in F_2, \exists \epsilon \in \{\pm 1\}, (g_-(X,Y), g_+(X,Y)) = k(Y,X)^\epsilon k^{-1}\}$*, where the map* $\mathrm{Aut}(F_2) \to (F_2)^2$ *is* $\theta \mapsto (\theta(X), \theta(Y))$*.*

The bijectivity of $\widetilde{\mathrm{SL}_2 \mathbb{Z}} \to \underline{R}_{ell}$, together with the equality $\underline{R}_{ell} = \{(g_+, g_-) \in (F_2)^2 | (g_-(X,Y), g_+(X,Y)) = (Y,X)\}$ are then proven in the following corollary to Theorem 23:

Corollary 24. *We have bijections*

$$\widetilde{\mathrm{SL}_2 \mathbb{Z}} \to \mathrm{Aut}_{(X,Y)}(F_2) \to \{(g_+, g_-) \in (F_2)^2 | (g_-(X,Y), g_+(X,Y)) = (Y,X)\},$$

where $\mathrm{Aut}_{(X,Y)}(F_2) = \{\theta \in \mathrm{Aut}(F_2) | \theta((X,Y)) = (X,Y)\}$*, the first map is as in Proposition 22 and the second map is* $\theta \mapsto (\theta(X), \theta(Y))$*.*

Proof of Corollary 24. The bijectivity of the second map follows from the injectivity of $\mathrm{Aut}(F_2) \to (F_2)^2$, $\theta \mapsto (\theta(X), \theta(Y))$ and from Theorem 23, 2. Let us now prove the bijectivity of the map $\widetilde{\mathrm{SL}_2 \mathbb{Z}} \to \mathrm{Aut}_{(X,Y)}(F_2)$. The kernel of $\widetilde{\mathrm{SL}_2 \mathbb{Z}} \to \mathrm{Aut}_{(X,Y)}(F_2)$ is contained in $\mathrm{Ker}(\widetilde{\mathrm{SL}_2 \mathbb{Z}} \to \mathrm{Aut}_{(X,Y)}(F_2) \to \mathrm{Out}(F_2) \to \mathrm{GL}_2(\mathbb{Z})) = \langle \Theta^4 \rangle$. On the other hand, $\widetilde{\mathrm{SL}_2 \mathbb{Z}} \to \mathrm{Aut}_{(X,Y)}(F_2)$ takes Θ^4 to $x \mapsto (X,Y)^{-1} x (X,Y)$, so the restriction of $\widetilde{\mathrm{SL}_2 \mathbb{Z}} \to \mathrm{Aut}_{(X,Y)}(F_2)$ to $\langle \Theta^4 \rangle$ is injective. It follows that $\widetilde{\mathrm{SL}_2 \mathbb{Z}} \to \mathrm{Aut}_{(X,Y)}(F_2)$ is injective.

Let us now show that $\widetilde{\mathrm{SL}_2 \mathbb{Z}} \to \mathrm{Aut}_{(X,Y)}(F_2)$ is surjective. We have a commutative diagram

$$\begin{tikzcd}
\mathrm{Aut}(F_2) \arrow[r] & \mathrm{Out}(F_2) \arrow[r, "\simeq"] & \mathrm{GL}_2(\mathbb{Z}) \\
\mathrm{Aut}_{(X,Y)}(F_2) \arrow[u, hook] \arrow[rr] & & \mathrm{SL}_2(\mathbb{Z}) \arrow[u, hook]
\end{tikzcd}$$

where the isomorphism follows from Theorem 23, 1. It follows that $\mathrm{Ker}(\mathrm{Aut}_{(X,Y)}(F_2) \to \mathrm{SL}_2(\mathbb{Z})) = \mathrm{Ker}(\mathrm{Aut}_{(X,Y)}(F_2) \to \mathrm{Out}(F_2)) = \mathrm{Aut}_{(X,Y)}(F_2) \cap \mathrm{Inn}(F_2) = \{\theta \in \mathrm{Aut}(F_2) | \exists k \in F_2, \theta(x) = kxk^{-1}$ and $\theta((X,Y)) = (X,Y)\} = \{\theta \in \mathrm{Aut}(F_2) | \exists n \in \mathbb{Z}, \theta(x) = (X,Y)^n x (X,Y)^{-n}\} = \langle x \mapsto (X,Y) x (X,Y)^{-1} \rangle$. As $x \mapsto (X,Y)x(X,Y)^{-1}$ is the image of $\Theta^4 \in \widetilde{\mathrm{SL}_2 \mathbb{Z}} \to \mathrm{Aut}_{(X,Y)}(F_2)$, $\mathrm{Ker}(\mathrm{Aut}_{(X,Y)}(F_2) \to \mathrm{SL}_2(\mathbb{Z}))$ contains $\mathrm{Im}(\widetilde{\mathrm{SL}_2 \mathbb{Z}} \to \mathrm{Aut}_{(X,Y)}(F_2))$. Since the composed map $\widetilde{\mathrm{SL}_2 \mathbb{Z}} \to \mathrm{Aut}_{(X,Y)}(F_2) \to \mathrm{SL}_2(\mathbb{Z})$ is surjective (the second map being given by abelianization), this implies that $\widetilde{\mathrm{SL}_2 \mathbb{Z}} \to \mathrm{Aut}_{(X,Y)}(F_2)$ is surjective. □ □

Remark 25. As $\mathrm{GL}_2(\mathbb{Z})$ is the nonoriented mapping class group of the topological torus, we have a morphism $\mathrm{GL}_2(\mathbb{Z}) \to \mathrm{Out}(B_{1,n})$. It lifts to $\widetilde{\mathrm{GL}_2 \mathbb{Z}} \to \mathrm{Aut}(B_{1,n})$, where $\Theta^4 \mapsto$ (conjugation by the image of $z \in P_n \to B_{1,n}$), and z is a generator of $Z(P_n) \simeq \mathbb{Z}$. The assignment {elliptic categories} → {representations of $B_{1,n}$} is then $\widetilde{\mathrm{GL}_2 \mathbb{Z}}$-equivariant.

4.4. **The semigroup scheme** $\underline{\mathrm{GT}}_{ell}(-)$**.** For $\mathbf{k}$ a $\mathbb{Q}$-ring, we set $R_{ell}(\mathbf{k}) := \mathrm{Ker}(\underline{\mathrm{GT}}_{ell}(\mathbf{k}) \to \underline{\mathrm{GT}}(\mathbf{k}))$. The assignments $\mathbf{k} \mapsto \underline{\mathrm{GT}}_{(ell)}(\mathbf{k}), R_{ell}(\mathbf{k})$ are functors {$\mathbb{Q}$-rings} → {semigroups}, i.e., semigroup schemes over $\mathbb{Q}$.

Proposition 26. *We have a commutative diagram of morphisms of semigroup schemes*

$$\begin{tikzcd}
R_{ell}(-) \arrow[r] \arrow[d] & \underline{\mathrm{GT}}_{ell}(-) \arrow[r] \arrow[d] & \underline{\mathrm{GT}}(-) \arrow[d] \\
\mathrm{SL}_2 \arrow[r] & \mathrm{M}_2 \arrow[r, "\det"] & \mathbb{A}
\end{tikzcd}$$

where $\underline{\mathrm{GT}}(\mathbf{k}) \to \mathbf{k}$ *is* $(\lambda, f) \mapsto \lambda$ *and* $\underline{\mathrm{GT}}_{ell}(\mathbf{k}) \to \mathrm{M}_2(\mathbf{k})$ *is* $(\lambda, f, g_\pm) \mapsto \left(\begin{smallmatrix}\alpha_+ & \beta_+ \\ \alpha_- & \beta_-\end{smallmatrix}\right)$*, where* $\log g_\pm(X,Y) = \alpha_\pm \log X + \beta_\pm \log Y$ *mod*[8] $[\hat{\mathfrak{f}}_2^{\mathbf{k}}, \hat{\mathfrak{f}}_2^{\mathbf{k}}]$.

Proof. It suffices to show that the right square is commutative, which follows by abelianization from the second part of (11). ☐

Recall that[9] $\mathrm{GT}(\mathbf{k}) = \underline{\mathrm{GT}}(\mathbf{k})^\times$. We set

Definition 27. $\mathrm{GT}_{ell}(\mathbf{k}) := \underline{\mathrm{GT}}_{ell}(\mathbf{k})^\times$.

Proposition 28. *1)* $\mathrm{GT}_{ell}(\mathbf{k}) = \underline{\mathrm{GT}}_{ell}(\mathbf{k}) \times_{\mathrm{M}_2(\mathbf{k})} \mathrm{GL}_2(\mathbf{k})$.

2) $R_{ell}(\mathbf{k})$ *is a group.*

Proof. Let us prove 1). If $(\lambda, f, g_\pm) \in \underline{\mathrm{GT}}_{ell}(\mathbf{k})$ is invertible as an element of $\underline{\mathrm{GT}}(\mathbf{k}) \times \mathrm{End}(F_2(\mathbf{k}))^{op}$, then one proves as in Proposition 19 that its inverse belongs to $\underline{\mathrm{GT}}_{ell}(\mathbf{k})$. The invertibility condition is then equivalent to the image of $(\lambda, f, g_\pm) \in \underline{\mathrm{GT}}_{ell}(\mathbf{k}) \to \mathrm{M}_2(\mathbf{k})$ lying in $\mathrm{GL}_2(\mathbf{k})$. 2) is then immediate. ☐

Recall that for any $\mathbb{Q}$-ring $\mathbf{k}$, $\mathrm{GT}_1(\mathbf{k}) = \mathrm{Ker}(\underline{\mathrm{GT}}(\mathbf{k}) \to \mathbf{k})$. We also set

$$\mathrm{GT}_{I_2}^{ell}(\mathbf{k}) := \mathrm{Ker}(\underline{\mathrm{GT}}_{ell}(\mathbf{k}) \to \mathrm{M}_2(\mathbf{k})), \quad R_{I_2}^{ell}(\mathbf{k}) := \mathrm{Ker}(R_{ell}(\mathbf{k}) \to \mathrm{SL}_2(\mathbf{k})).$$

Then $\mathbf{k} \mapsto \mathrm{GT}_{(I_2)}^{(ell)}(\mathbf{k})$, $R_{(I_2)}^{ell}(\mathbf{k})$ are $\mathbb{Q}$-group schemes. It is known that $\mathrm{GT}_1(-)$ is unipotent.

Proposition 29. *The group schemes* $\mathrm{GT}_{I_2}^{ell}(-)$ *and* $R_{I_2}^{ell}(-)$ *are prounipotent.*

Proof. $\mathrm{GT}_{I_2}^{ell}(\mathbf{k}) \subset \mathrm{GT}_1(\mathbf{k}) \times \mathrm{Aut}_{I_2}(F_2(\mathbf{k}))^{op}$, where $\mathrm{Aut}_{I_2}(F_2(\mathbf{k})) = \mathrm{Ker}(\mathrm{Aut}(F_2(\mathbf{k})) \to \mathrm{GL}_2(\mathbf{k}))$; $\mathbf{k} \mapsto \mathrm{Aut}_{I_2}(F_2(\mathbf{k}))$ is prounipotent, so $\mathbf{k} \mapsto \mathrm{GT}_1(\mathbf{k})$ is prounipotent as the subgroup of a prounipotent group scheme. The same argument implies that $R_{I_2}^{ell}(-)$ is prounipotent. ☐

Proposition 30. *We have exact sequences* $1 \to R_{I_2}^{ell}(\mathbf{k}) \to R_{ell}(\mathbf{k}) \to \mathrm{SL}_2(\mathbf{k}) \to 1$ *and* $1 \to \mathrm{GT}_{I_2}^{ell}(\mathbf{k}) \to \mathrm{GT}_{ell}(\mathbf{k}) \to \mathrm{GL}_2(\mathbf{k}) \to 1$.

Proof. We need to prove that $R_{ell}(\mathbf{k}) \to \mathrm{SL}_2(\mathbf{k})$ is surjective. Set $G(\mathbf{k}) := \mathrm{Im}(R_{ell}(\mathbf{k}) \to \mathrm{SL}_2(\mathbf{k}))$, then $\mathbf{k} \mapsto G(\mathbf{k})$ is a group subscheme of SL_2. We have morphisms $\mathbb{G}_a \to R_{ell}(-)$, extending $\mathbb{Z} \to \widetilde{\mathrm{SL}_2 \mathbb{Z}}$, $1 \mapsto \Psi_\pm$ (where $\Psi_+ = \Psi$, $\Psi_- = \Theta\Psi\Theta^{-1}$) in the sense that

$$\begin{tikzcd}
\widetilde{\mathrm{SL}_2 \mathbb{Z}} \arrow[r] & R_{ell}(\mathbf{k}) \\
\mathbb{Z} \arrow[u] \arrow[r] & (\mathbf{k}, +) \arrow[u]
\end{tikzcd}$$

commutes; then $\mathbb{G}_a \to R_{ell}(-) \to \mathrm{SL}_2$ are the morphisms $t \mapsto \left(\begin{smallmatrix}1 & t \\ 0 & 1\end{smallmatrix}\right), \left(\begin{smallmatrix}1 & 0 \\ -t & 1\end{smallmatrix}\right)$. So the Lie algebra of $G(-)$ contains both $\left(\begin{smallmatrix}0 & 1 \\ 0 & 0\end{smallmatrix}\right)$ and $\left(\begin{smallmatrix}0 & 0 \\ 1 & 0\end{smallmatrix}\right)$, hence equals $\mathfrak{sl}_2$, so $G = \mathrm{SL}_2$.

Let us now prove that $\mathrm{GT}_{ell}(\mathbf{k}) \to \mathrm{GL}_2(\mathbf{k})$ is surjective. Set $\tilde{G}(\mathbf{k}) := \mathrm{Im}(\mathrm{GT}_{ell}(\mathbf{k}) \to \mathrm{GL}_2(\mathbf{k}))$, then $\mathrm{SL}_2 \subset \tilde{G}(-) \subset \mathrm{GL}_2$. We will construct in Section 4.6 a semigroup scheme morphism $\underline{\mathrm{GT}}(-) \xrightarrow{\sigma} \underline{\mathrm{GT}}_{ell}(-)$, such that

$$\begin{tikzcd}
\underline{\mathrm{GT}}(-) \arrow[r] \arrow[d, "\sigma"'] & \mathbb{A} \arrow[d] \\
\underline{\mathrm{GT}}_{ell}(-) \arrow[r] & \mathrm{M}_2
\end{tikzcd}$$

commutes, where $\mathbb{A} \to \mathrm{M}_2$ is $t \mapsto \left(\begin{smallmatrix}t & 0 \\ 0 & 1\end{smallmatrix}\right)$. Then $\tilde{G}(-)$ contains the image of $\mathbb{G}_m \to \mathrm{GT}(-) \xrightarrow{\sigma} \mathrm{GT}_{ell}(-) \to \mathrm{GL}_2$, where $\mathbb{G}_m \to \mathrm{GT}(-)$ is a section of $\mathrm{GT}(-) \to \mathbb{G}_m$ (see [Dr]), which is the image of $\mathbb{G}_m \to \mathrm{GL}_2$, $t \mapsto \left(\begin{smallmatrix}t & 0 \\ 0 & 1\end{smallmatrix}\right)$. Then $\mathrm{Lie}(\tilde{G}) = \mathfrak{gl}_2$, so $\tilde{G} = \mathrm{GL}_2$, as wanted. ☐

---

[8] Recall that $F_2(\mathbf{k}) = \exp(\hat{\mathfrak{f}}_2^{\mathbf{k}})$, where $\hat{\mathfrak{f}}_2^{\mathbf{k}}$ is the topologically free $\mathbf{k}$-Lie algebra in two generators $\log X$ and $\log Y$.

[9] If S is a semigroup with unit, $S^\times$ is the group of its invertible elements.

4.5. **The Zariski closure $\langle B_3\rangle \subset R_{ell}(-)$.** Recall that we have a group morphism $B_3 = R_{ell} \to R_{ell}(\mathbb{Q})$. The Zariski closure $\langle B_3\rangle \subset R_{ell}(-)$ is then the subgroup scheme defined as

$$\langle B_3\rangle := \bigcap_{\substack{G\subset R_{ell}(-) \text{ subgroup scheme}\,| \\ G(\mathbb{Q})\supset \mathrm{Im}(B_3\to R_{ell}(\mathbb{Q}))}} G$$

Let us compute the Lie algebra[10] inclusion $\mathrm{Lie}\langle B_3\rangle \subset \mathrm{Lie}\, R_{ell}(-)$. First, $\mathrm{Lie}\, R_{ell}(-)$ is a Lie subalgebra of[11] $\mathrm{Lie}\,\mathrm{Aut}(F_2(-))^{op} = \{(\alpha_+,\alpha_-) \in \hat{\mathfrak{f}}_2 \times \hat{\mathfrak{f}}_2\}$, with Lie bracket $[(\alpha_+,\alpha_-),(\alpha'_+,\alpha'_-)] := (D_{\alpha'_+,\alpha'_-}(\alpha_+), D_{\alpha'_+,\alpha'_-}(\alpha_-)) - (\alpha_\pm \leftrightarrow \alpha'_\pm)$, where $D_{\alpha_+,\alpha_-} \in \mathrm{Der}(\hat{\mathfrak{f}}_2)$ is given by $\xi_\pm \mapsto \alpha_\pm$. We find that $\mathrm{Lie}\, R_{ell}(-)$ is the set of (α_+,α_-) such that

$$\begin{aligned}&\mathrm{Ad}(X_1^+)^{-1}(\tilde\alpha_+(\log X_1^+, \log X_1^-)) + \mathrm{Ad}(X_3^+)(\tilde\alpha_+(\log X_2^+\sigma_1^{-2}, \log \sigma_1^2 X_2^-)) \\ &+\tilde\alpha_+(\log X_3^+\sigma_2^{-1}\sigma_1^{-2}\sigma_2^{-1}, \log \sigma_2\sigma_1^2\sigma_2 X_3^-) = 0,\end{aligned}$$

$$\begin{aligned}&\mathrm{Ad}(X_1^-)^{-1}(\tilde\alpha_-(\log X_1^+, \log X_1^-)) + \mathrm{Ad}(X_3^-)(\tilde\alpha_-(\log \sigma_1^{-2}X_2^+, \log X_2^-\sigma_1^2)) \\ &+\tilde\alpha_-(\log \sigma_2^{-1}\sigma_1^{-2}\sigma_2^{-1}X_3^+, X_3^- \log \sigma_2\sigma_1^2\sigma_2) = 0,\end{aligned}$$

$$(\mathrm{Ad}((X_2^-)^{-1}\sigma_1^2){-}1)(\tilde\alpha_+(\log X_1^+, \log X_1^-)){+}\mathrm{Ad}(\sigma_1^{-1}(X_1^-)^{-1})(\mathrm{Ad}(X_2^+\sigma_1^{-2}){-}1)(\tilde\alpha_-(\log X_1^+, \log X_1^-)) = 0$$

where $\tilde\alpha_\pm(\xi_+,\xi_-) = \frac{e^{\mathrm{ad}\,\xi_\pm}-1}{\mathrm{ad}\,\xi_\pm}(\alpha_\pm(\xi_+,\xi_-))$ (relations in $\mathrm{Lie}\, P_{1,3}(-)$). These relations imply

$$\tilde\alpha_+(\log X_1^+, \log X_1^-) + \mathrm{Ad}(X_1^+)(\tilde\alpha_+(\log X_2^+\sigma_1^{-2}, \log \sigma_1^2 X_2^-)) = 0,$$

$$\tilde\alpha_-(\log X_1^+, \log X_1^-) + \mathrm{Ad}(X_1^-)(\tilde\alpha_-(\log \sigma_1^{-2}X_2^+, \log X_2^-\sigma_1^2)) = 0,$$

$$\big(\mathrm{Ad}(X_1^-, X_1^+){-}\mathrm{Ad}\, X_1^-\big)(\tilde\alpha_+(\log X_1^+, \log X_1^-)){+}\big(\mathrm{Ad}(X_1^- X_1^+ (X_1^-)^{-1}{-}1)\big)(\tilde\alpha_-(\log X_1^+, \log X_1^-)) = 0$$

(relations in $\mathrm{Lie}\, P_{1,2}(-) \simeq \hat{\mathfrak{f}}_2$).

We now compute $\mathrm{Lie}\langle B_3\rangle \subset \mathrm{Lie}\, R_{ell}(-)$.

Lemma 31. *Let* $u_+ := (0, \frac{\mathrm{ad}\,\xi_-}{1-e^{-\mathrm{ad}\,\xi_-}}(\xi_+))$, $u_- := (\frac{\mathrm{ad}\,\xi_+}{1-e^{-\mathrm{ad}\,\xi_+}}(\xi_-), 0)$, *then* $u_\pm \in \mathrm{Lie}\langle B_3\rangle$.

Proof. We have morphisms $\mathbb{G}_a \to \langle B_3\rangle \subset \mathrm{Aut}(F_2(-))^{op}$, extending $\mathbb{Z} \to B_3$, $1 \mapsto \Psi_\pm^{\pm 1}$. The corresponding morphisms $(\mathbf{k},+) \to \mathrm{Aut}(F_2(\mathbf{k}))^{op}$ are $t \mapsto (X \mapsto X, Y \mapsto YX^t)$ and $t \mapsto (X \mapsto XY^t, Y \mapsto Y)$. The associated Lie algebra morphisms are $\mathbb{Q} \to \mathrm{Lie}\,\mathrm{Aut}(F_2(-))^{op}$, $1 \mapsto u_\pm$, which proves that $u_\pm \in \mathrm{Lie}\langle B_3\rangle$. $\square$

Proposition 32. $\mathrm{Lie}\langle B_3\rangle \subset \mathrm{Lie}\,\mathrm{Aut}(F_2(-))^{op} \simeq \hat{\mathfrak{f}}_2^2$ *is the smallest closed Lie subalgebra containing* u_+ *and* u_-.

We first prove:

Lemma 33. *Let G be a proalgebraic group scheme over $\mathbb{Q}$ fitting in $1 \to U \to G \to G_0 \to 1$, where G_0 is semisimple and U is prounipotent. Let $0 \to \mathfrak{u} \to \mathfrak{g} \to \mathfrak{g}_0 \to 0$ be the corresponding exact sequence of Lie algebras. Then $H \mapsto \mathrm{Lie}\, H$ sets up a bijection* {*proalgebraic subgroups* $H \subset G$, *such that* $\mathrm{Im}(H \subset G \to G_0) = G_0$} $\overset{\sim}{\to}$ {*closed Lie subalgebras* $\mathfrak{h} \subset \mathfrak{g}$, *such that* $\mathrm{Im}(\mathfrak{h} \subset \mathfrak{g} \to \mathfrak{g}_0) = \mathfrak{g}_0$}.

---

[10] Recall that the Lie algebra of a $\mathbb{Q}$-group scheme G is $\mathrm{Ker}(G(\mathbb{Q}[\varepsilon]/(\varepsilon^2)) \to G(\mathbb{Q}))$.

[11] We set $\hat{\mathfrak{f}}_2 := \hat{\mathfrak{f}}_2^{\mathbb{Q}}$, and $\xi_+ := \log X$, $\xi_- := \log Y$.

Proof. If H is in the first set, then we have an exact sequence $1 \to H \cap U \to H \to G_0 \to 1$, where $H \cap U$ is necessarily prounipotent, hence connected, which implies that H is connected. So the map $H \mapsto \mathrm{Lie}\, H$ is injective.

Let us prove its surjectivity. Let $\mathfrak{h}$ belong to the second set.

First note that according to the Levi-Mostow decomposition ([Mo], [BS] prop. 5.1), there exists a section $\tilde\sigma : G_0 \to G$ of $G \to G_0$. We denote by $\sigma : \mathfrak{g}_0 \to \mathfrak{g}$ its infinitesimal. Any section of $\mathfrak{g} \to \mathfrak{g}_0$ is then conjugate to σ by an element of $U(\mathbb{Q})$.

Then we have an exact sequence $0 \to \mathfrak{h} \cap \mathfrak{u} \to \mathfrak{h} \to \mathfrak{g}_0 \to 0$; applying the Levi decomposition theorem for Lie algebras, we obtain a section $\tau : \mathfrak{g}_0 \to \mathfrak{h}$ of $\mathfrak{h} \to \mathfrak{g}_0$. Now the composite map $\mathfrak{g}_0 \xrightarrow{\tau} \mathfrak{h} \hookrightarrow \mathfrak{g}$ is a section of $\mathfrak{g} \to \mathfrak{g}_0$, hence of the form $\mathrm{Ad}(x) \circ \sigma$, where $x \in U(\mathbb{Q})$. If $\mathfrak{v} := \mathfrak{h} \cap \mathfrak{u}$, we then have $[\mathrm{Ad}(x)(\sigma(\mathfrak{g}_0)), \mathfrak{v}] \subset \mathfrak{v}$.

Let then $V \subset U$ be the subgroup with Lie algebra $\mathfrak{v}$; if we set $H := V \cdot \mathrm{Ad}(x)(\tilde\sigma(G_0)) = \mathrm{Ad}(x)(\tilde\sigma(G_0)) \cdot V$, then H is in the first set, and has Lie algebra $\mathfrak{h}$. □

Proof of Proposition 32. Let $\mathrm{Lie}(u_+, u_-)$ be the smallest closed subalgebra of $\mathrm{Lie}\,\mathrm{Aut}(F_2(-))^{op}$ containing u_+ and u_-. Then $\mathrm{Lie}\langle B_3 \rangle \supset \mathrm{Lie}(u_+, u_-)$. Apply now Lemma 33 with $G = R_{ell}(-)$, $G_0 = \mathrm{SL}_2$. The map $\mathfrak{g} \to \mathfrak{g}_0$ is such that $u_+ \mapsto \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$ and $u_- \to \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$, so if $\mathfrak{h} := \mathrm{Lie}(u_+, u_-)$, then $\mathrm{Im}(\mathfrak{h} \subset \mathfrak{g} \to \mathfrak{g}_0) = \mathfrak{g}_0$. Let then $H \subset R_{ell}(-)$ be the proalgebraic subgroup corresponding to $\mathfrak{h}$ by Lemma 33; then $\langle B_3 \rangle \supset H$. On the other hand, we have group morphisms $\mathbb{G}_a \to H$ corresponding to $\mathbb{Q} \to \mathfrak{h}$, $1 \mapsto u_\pm$, whose versions over $\mathbb{Q}$ are $(\mathbb{Q}, +) \to H(\mathbb{Q}) \subset \mathrm{Aut}(\widetilde{F_2(\mathbb{Q})})^{op}$, $t \mapsto \Psi_\pm^t$. Setting $t = 1$, we obtain $H(\mathbb{Q}) \ni \Psi_\pm$, and as Ψ_+, Ψ_- generate B_3, $H(\mathbb{Q}) \supset \widetilde{\mathrm{SL}_2 \mathbb{Z}}$. So $\langle B_3 \rangle = H$. Taking Lie algebras, we obtain Proposition 32. □

Remark 34. Let $d_\pm := [[u_+, u_-], u_\pm] \pm 2u_\pm$. Then for any $(\alpha_\pm, \beta) \in \mathbb{N}^3$,

$$x^\pm_{\alpha_\pm, \beta} := \mathrm{ad}(u_+^{\alpha_+} u_-^{\alpha_-} [u_+, u_-]^\beta)(d_\pm) \in \mathrm{Ker}(\mathrm{Lie}\langle \widetilde{\mathrm{SL}_2 \mathbb{Z}} \rangle \to \mathfrak{sl}_2).$$

Then $\mathrm{Ker}(\mathrm{Lie}\langle B_3 \rangle \to \mathfrak{sl}_2)$ is topologically generated by these elements, more precisely, it is equal to $\{\sum_{n \geq 1} P_n((x^\pm_{\alpha_\pm, \beta})) | (P_n)_n \in \prod_{n \geq 1} \mathfrak{f}_n\}$, where $\mathfrak{f}$ is the free Lie algebra with generators indexed by $\mathbb{N}^3 \times \{\pm\}$, and $\mathfrak{f}_n$ is its part of degree n (each generator having degree 1). Then $\mathrm{Lie}\langle B_3 \rangle = \mathrm{Ker}(\mathrm{Lie}\langle B_3 \rangle \to \mathfrak{sl}_2) \oplus \mathrm{Span}_{\mathbb{Q}}(u_+, u_-, [u_+, u_-])$.

4.6. **A morphism $\underline{\mathrm{GT}} \to \underline{\mathrm{GT}}_{ell}$ and its variants.** We now construct a section of the semigroup morphism $\underline{\mathrm{GT}}_{ell} \to \underline{\mathrm{GT}}$ and of its variants.

Proposition 35. *There exists a unique semigroup morphism $\underline{\mathrm{GT}} \to \underline{\mathrm{GT}}_{ell}$, defined by $(\lambda, f) \mapsto (\lambda, f, g_\pm)$, where*

$$g_+(X, Y) = f(X, (Y, X)) X^\lambda f(X, (Y, X))^{-1},$$
$$g_-(X, Y) = (Y, X)^{\frac{\lambda - 1}{2}} f(Y X^{-1} Y^{-1}, (Y, X)) Y f(X, (Y, X))^{-1}.$$

The same formulas define semigroup morphisms $\widehat{\underline{\mathrm{GT}}} \to \widehat{\underline{\mathrm{GT}}}_{ell}$, $\underline{\mathrm{GT}}_l \to \underline{\mathrm{GT}}_l^{ell}$, and a semigroup scheme morphism $\underline{\mathrm{GT}}(-) \to \underline{\mathrm{GT}}_{ell}(-)$, compatible with the natural maps between the various versions of $\underline{\mathrm{GT}}_{(ell)}$.

We have commutative diagrams

$$\begin{tikzcd}
\underline{\mathrm{GT}} \arrow[r] \arrow[d, "\simeq"'] & \underline{\mathrm{GT}}_{ell} \arrow[d, "\simeq"] & \text{and} & \underline{\mathrm{GT}}(-) \arrow[r] \arrow[d] & \underline{\mathrm{GT}}_{ell}(-) \arrow[d] \\
\mathbb{Z}/2 \arrow[r] & \widetilde{\mathrm{GL}_2 \mathbb{Z}} & & \mathbb{A} \arrow[r] & \mathrm{M}_2
\end{tikzcd}$$

where the bottom morphisms are $\bar{1} \mapsto \varepsilon\Theta$ and $\lambda \mapsto \begin{pmatrix} \lambda & 0 \\ 0 & 1 \end{pmatrix}$.

Proof. 1) We show that g_+ satisfies (9) with $\pm = +$, i.e.,

$$\big(\sigma_2\sigma_1(\sigma_1\sigma_2^2\sigma_1)^m f(X_1^+, (X_1^-, X_1^+))(X_1^+)^\lambda f^{-1}(X_1^+, (X_1^-, X_1^+)) f(\sigma_1^2, \sigma_2^2)\big)^3 = 1 \tag{13}$$

in $B_{1,3}$.

Denote by $\sigma_0,\dots,\sigma_n$ the generators of B_{n+1}, identify $S_{n+1}\simeq \mathrm{Perm}(\{0,\dots,n\})$ and let $S_n\subset S_{n+1}$ be $\{\sigma|\sigma(0)=0\}$. We have a unique morphism

$$(B_{n+1}/Z(B_{n+1}))\times_{S_{n+1}} S_n\to B_{1,n},$$

defined by $\sigma_0^2\to X_1^+$, $\sigma_i\mapsto\sigma_i$ $(i>0)$. Indeed, using a presentation of $B_{n+1}\times_{S_{n+1}}S_n$ (a braid group of type B), one proves that the same formulas define a morphism $B_{n+1}\times_{S_{n+1}}S_n\to B_{1,n}$; a generator of $Z(B_{n+1})\simeq\mathbb{Z}$ then maps to $X_1^+\cdots X_n^+=1$. This morphism may be viewed as induced by the map $Cf_{[n]}(\mathbb{R}\times S^1)\to Cf_{[n]}(S^1\times S^1)$ by taking fundamental groups, where $Cf_{[n]}(X)=(X^n-\{\text{diagonals}\})/(S_n\times X)$ (where X is assumed to be a group, acting diagonally).

The l.h.s. of (13) is then the image by $B_{n+1}\times_{S_{n+1}}S_n\to B_{1,n}$ of

$$\big(\sigma_1\sigma_2(\sigma_1\sigma_2^2\sigma_1)^m f(\sigma_0^2,\sigma_1\sigma_2^2\sigma_1)\sigma_0^{2\lambda}f^{-1}(\sigma_0^2,\sigma_1\sigma_2^2\sigma_1)f(\sigma_1^2,\sigma_2^2)\big)^3. \tag{14}$$

Now $(\lambda,f)\in\underline{\mathrm{GT}}$ induces an endomorphism of $\mathbf{PaB}$, and therefore an endomorphism $F^O_{\lambda,f}$ of $B_n=\mathbf{PaB}(O,O)$ for any $O\in\mathbf{PaB}$, $|O|=n$. It is such that $F^O_{\lambda,f}(Z)=Z^\lambda$ for any $Z\in Z(B_n)$. Then one proves that $(14)=(F_{\lambda,f}^{\bullet((\bullet\bullet)\bullet)}(\sigma_1\sigma_2\sigma_0^2))^3=F_{\lambda,f}^{\bullet((\bullet\bullet)\bullet)}((\sigma_1\sigma_2\sigma_0^2)^3)$. Now $(\sigma_1\sigma_2\sigma_0^2)^3$ is a generator of $Z(B_4)$ so $(14)=((\sigma_1\sigma_2\sigma_0^2)^3)^\lambda\in Z(B_4)$. As $Z(B_4)$ is in the kernel of $B_4\times_{S_4}S_3\to B_{1,3}$, the l.h.s. of (13) equals 1.

2) We show that g_- satisfies (9) with $\pm=-$, i.e.,

$$(\sigma_2^{-1}\sigma_1^{-1}(\sigma_1\sigma_2^2\sigma_1)^{-m}g_-(X_1^+,X_1^-)f(\sigma_1^2,\sigma_2^2))^3=1$$

in $B_{1,3}$. As $(X_1^-,X_1^+)=\sigma_1\sigma_2^2\sigma_1$, this is rewritten

$$(Y_1A\sigma_2^{-1}\sigma_1^{-1})^3=1, \tag{15}$$

where $A=f^{-1}(X_1^+,\sigma_1\sigma_2^2\sigma_1)f(\sigma_1^2,\sigma_2^2)f((X_3^+)^{-1},\sigma_2\sigma_1^2\sigma_2)$. The pentagon identity implies that

$$A=f(X_2^+,\sigma_2^2)^{-1}g(X_1^+,\sigma_1^2). \tag{16}$$

On the other hand, (15) is rewritten

$$X_1^-AX_3^-(\sigma_2\sigma_1A\sigma_1^{-1}\sigma_2^{-1})X_2^-(\sigma_1\sigma_2A\sigma_2^{-1}\sigma_1^{-1})=1. \tag{17}$$

Using (16) and $X_1^-X_2^+(X_1^-)^{-1}=X_2^+\sigma_1^{-2}$, $(X_1^-,\sigma_2)=1$, $(X_3^-)^{-1}X_1^+X_3^-=\sigma_2\sigma_1^2\sigma_2X_1^+$, $((X_3^-)^{-1},\sigma_1)=1$, we get

$$X_1^-AX_3^-=f(X_2^+\sigma_1^{-2},\sigma_2^2)(X_2^-)^{-1}f(\sigma_2\sigma_1^2\sigma_2^{-1}X_1^+,\sigma_1^2),$$

so (17) is rewritten

$$(X_2^-)^{-1}f(\sigma_2\sigma_1^2\sigma_2^{-1}X_1^+,\sigma_1^2)(\sigma_2\sigma_1A\sigma_1^{-1}\sigma_2^{-1})X_2^-(\sigma_1\sigma_2A\sigma_2^{-1}\sigma_1^{-1})f(X_2^+\sigma_1^{-2},\sigma_2^2)^{-1}=1$$

or using $\sigma_1\sigma_2f(X_1^+,\sigma_1^2)\sigma_2^{-1}\sigma_1^{-1}=f(X_2^+\sigma_1^{-2},\sigma_2^2)$ and $X_2^-=\sigma_1^{-1}X_1^-\sigma_1^{-1}$, as follows

$$f(\sigma_2^2X_2^+\sigma_1^{-2},\sigma_1^2)(\sigma_1\sigma_2\sigma_1A\sigma_1^{-1}\sigma_2^{-1}\sigma_1^{-1})X_1^-f^{-1}(\sigma_2X_2^+\sigma_2^{-1},\sigma_2^2)(X_1^-)^{-1}=1,$$

which using $X_1^-(\sigma_2X_2^+\sigma_2^{-1})(X_1^-)^{-1}=X_1^-(X_3^+\sigma_2^{-2})(X_1^-)^{-1}=X_3^+\sigma_2^{-1}\sigma_1^{-2}\sigma_2^{-1}$ and $(X_1^-,\sigma_2)=1$ is rewritten

$$f(\sigma_2^2X_2^+\sigma_1^{-2},\sigma_1^2)(\sigma_1\sigma_2\sigma_1A\sigma_1^{-1}\sigma_2^{-1}\sigma_1^{-1})f^{-1}(X_3^+\sigma_2^{-1}\sigma_1^{-2}\sigma_2^{-1},\sigma_2^2)=1.$$

The latter identity holds as $\sigma_1\sigma_2\sigma_1X_1^+(\sigma_1\sigma_2\sigma_1)^{-1}=X_3^+(\sigma_2\sigma_1^2\sigma_2)^{-1}$, $(\sigma_1\sigma_2\sigma_1)\sigma_1^2(\sigma_1\sigma_2\sigma_1)^{-1}=\sigma_2^2$, $\sigma_1\sigma_2\sigma_1X_2^+(\sigma_1\sigma_2\sigma_1)^{-1}=\sigma_2^2X_2^+\sigma_1^{-2}$, $(\sigma_1\sigma_2\sigma_1)\sigma_2^2(\sigma_1\sigma_2\sigma_1)^{-1}=\sigma_1^2$.

3) We show that (g_+,g_-) satisfies (10), i.e., $u^2=(ug_+^{-1}u^{-1},g_-^{-1})$, where $u=f(\sigma_1^2,\sigma_2^2)\sigma_1^\lambda f(\sigma_1^2,\sigma_2^2)^{-1}$, $g_\pm=g_\pm(X_1^+,X_1^-)$. This is rewritten as $g_-ug_+ug_-^{-1}=ug_+u^{-1}$, or

$$\begin{aligned}&X_1^-\cdot f(X_1^+,(X_1^-,X_1^+))^{-1}ug_+uf(X_1^+,(X_1^-,X_1^+))\cdot(X_1^-)^{-1}\\&=\mathrm{Ad}_{f((X_1^-,X_1^+),(X_1^+)^{-1}(X_1^-,X_1^+)^{-1})(X_1^-,X_1^+)^{\frac{1-\lambda}{2}}f(\sigma_1^2,\sigma_2^2)\sigma_1^\lambda f(\sigma_1^2,\sigma_2^2)^{-1}f(X_1^+,(X_1^-,X_1^+))}(X_1^+)^\lambda.\end{aligned} \tag{18}$$

Now $f(X_1^+,(X_1^-,X_1^+))^{-1}ug_+uf(X_1^+,(X_1^-,X_1^+)) = F_{\lambda,f}^{(\bullet\bullet)(\bullet\bullet)}(\sigma_1\sigma_0^2\sigma_1) = f(X_2^+,\sigma_2^2)(X_2^+)^\lambda f(X_2^+,\sigma_2^2)^{-1}$ and as $(X_1^-,\sigma_2)=1$, $(X_1^-,(X_2^+)^{-1})=\sigma_1^2$, the l.h.s. of (18) is

$$\mathrm{Ad}_{f(X_2^+\sigma_1^{-2},\sigma_2^2)}(X_2^+\sigma_1^{-2})^\lambda = \mathrm{Ad}_{f(X_2^+\sigma_1^{-2},\sigma_2^2)\sigma_1}(X_1^+)^\lambda.$$

To show (18), it is then enough to prove that

$$\begin{aligned} A \quad := \quad & \sigma_1^{-1}f^{-1}(X_2^+\sigma_1^{-2},\sigma_2^2)f((X_1^-,X_1^+),(X_1^+)^{-1}(X_1^-,X_1^+)^{-1})(X_1^-,X_1^+)^{\frac{1-\lambda}{2}} \\ & f(\sigma_1^2,\sigma_2^2)\sigma_1^\lambda f^{-1}(\sigma_1^2,\sigma_2^2)f(X_1^+,(X_1^-,X_1^+)) \end{aligned}$$

commutes with X_1^+.

We will prove that

$$A = \sigma_2 f^{-1}(X_2^+,\sigma_2^2)\sigma_2^{-\lambda}, \tag{19}$$

which by $(X_1^+,\sigma_2)=(X_1^+,X_2^+)=1$ implies our claim. We have

$$\begin{aligned} & \sigma_2^\lambda f(X_2^+,\sigma_2^2)\sigma_2^{-1}A \\ & = \sigma_2^\lambda f(X_2^+,\sigma_2^2)f^{-1}(X_1^+,\sigma_1^2)\sigma_2^{-1}\sigma_1^{-1}f((X_1^-,X_1^+),(X_1^+)^{-1}(X_1^-,X_1^+)^{-1}) \\ & (X_1^-,X_1^+)^{\frac{1-\lambda}{2}}f(\sigma_1^2,\sigma_2^2)\sigma_1^\lambda f^{-1}(\sigma_1^2,\sigma_2^2)f(X_1^+,(X_1^-,X_1^+)), \end{aligned}$$

which is the image under $B_4\times_{S_4}S_3\to B_{1,3}$ of

$$\begin{aligned} X := \qquad & \sigma_2^\lambda f(\sigma_1\sigma_0^2\sigma_1,\sigma_2^2)f^{-1}(\sigma_0^2,\sigma_1^2)\sigma_2^{-1}\sigma_1^{-1}f(\sigma_1\sigma_2^2\sigma_1,\sigma_0^{-2}\sigma_1^{-1}\sigma_2^{-2}\sigma_1^{-1}) \\ & (\sigma_1\sigma_2^2\sigma_1)^{\frac{1-\lambda}{2}}f(\sigma_1^2,\sigma_2^2)\sigma_1^\lambda f^{-1}(\sigma_1^2,\sigma_2^2)f(\sigma_0^2,\sigma_1\sigma_2^2\sigma_1). \end{aligned}$$

We have

$$(\sigma_1\sigma_2^2\sigma_1)^{\frac{1-\lambda}{2}}f(\sigma_1^2,\sigma_2^2)\sigma_1^\lambda f^{-1}(\sigma_1^2,\sigma_2^2) = \sigma_1\sigma_2 f^{-1}(\sigma_1^2,\sigma_2^2)\sigma_2^{-\lambda}$$

because the hexagon identity, together with $\sigma_1^{-1}\sigma_2^2\sigma_1 = \sigma_2\sigma_1^2\sigma_2^{-1} \equiv (\sigma_2^2\sigma_1^2)^{-1} \mod Z(P_3)$, implies $(\sigma_2^2)^m f(\sigma_2\sigma_1^2\sigma_2^{-1},\sigma_2^2)(\sigma_2^2\sigma_1^2)^{\frac{1-\lambda}{2}}f(\sigma_1^2,\sigma_1^{-1}\sigma_2^2\sigma_1)(\sigma_1^2)^m f(\sigma_2^2,\sigma_1^2)=1$.

Then

$$X = \sigma_2^\lambda f(\sigma_1\sigma_0^2\sigma_1,\sigma_2^2)f^{-1}(\sigma_0^2,\sigma_1^2)\sigma_2^{-1}\sigma_1^{-1}f(\sigma_1\sigma_2^2\sigma_1,\sigma_0^{-2}\sigma_1^{-1}\sigma_2^{-2}\sigma_1^{-1})\sigma_1\sigma_2 f^{-1}(\sigma_1^2,\sigma_2^2)\sigma_2^{-\lambda}f(\sigma_0^2,\sigma_1\sigma_2^2\sigma_1).$$

Define Y as the r.h.s. of this equality with the factors $\sigma_2^{\pm\lambda}$ removed. As σ_2 commutes zith σ_0^2 and $\sigma_1\sigma_2^2\sigma_1$, the equality $Y=1$ implies $X=1$. Y is rewritten as

$$Y = f(\sigma_1\sigma_0^2\sigma_1,\sigma_2^2)f^{-1}(\sigma_0^2,\sigma_1^2)f(\sigma_2\sigma_1^2\sigma_2,\sigma_2^{-1}\sigma_1^{-1}\sigma_0^{-2}\sigma_1^{-1}\sigma_2^{-1})f^{-1}(\sigma_1^2,\sigma_2^2)f(\sigma_0^2,\sigma_1\sigma_2^2\sigma_1).$$

Now

$$\begin{aligned} f(\sigma_2\sigma_1^2\sigma_2,\sigma_2^{-1}\sigma_1^{-1}\sigma_0^{-2}\sigma_1^{-1}\sigma_2^{-1}) \qquad & = f(\sigma_2\sigma_1^2\sigma_2,(\sigma_0\sigma_1^2\sigma_0)\sigma_1^2) \\ & \text{as} \quad \sigma_2^{-1}\sigma_1^{-1}\sigma_0^{-2}\sigma_1^{-1}\sigma_2^{-1} \equiv (\sigma_0\sigma_1^2\sigma_0)\sigma_1^2 \quad \mathrm{mod} \quad Z(P_4) \\ & = f(\sigma_2\sigma_1^2\sigma_2,\sigma_0\sigma_1^2\sigma_0) \\ & \text{as} \quad \sigma_1^2 \text{ commutes with } \sigma_2\sigma_1^2\sigma_2 \text{ and } \sigma_0\sigma_1^2\sigma_0. \end{aligned}$$

So

$$\begin{aligned} Y \quad & = \quad f(\sigma_1\sigma_0^2\sigma_1,\sigma_2^2)f^{-1}(\sigma_0^2,\sigma_1^2)f(\sigma_2\sigma_1^2\sigma_2,\sigma_0\sigma_1^2\sigma_1)f^{-1}(\sigma_1^2,\sigma_2^2)f(\sigma_0^2,\sigma_1\sigma_2^2\sigma_1) \\ & = \quad 1 \text{ by the pentagon identity.} \end{aligned}$$

It follows that $Y=1$, so (19) holds, which implies (18).

4) The fact that $\underline{\mathrm{GT}}\to\underline{\mathrm{GT}}_{ell}$, $(\lambda,f)\mapsto(\lambda,f,g_\pm)$ is a morphism of semigroups follows from the identity $(g_-(X,Y),g_+(X,Y))=(Y,X)^\lambda$. It is straightforward to check the commutativity of the first diagram; in the second diagram follows from $((\lambda,f)\in\underline{\mathrm{GT}}(\mathbf{k}))\Rightarrow(\log f\in[\hat{\mathfrak{f}}_2^{\mathbf{k}},\hat{\mathfrak{f}}_2^{\mathbf{k}}])$.

5) The arguments used in the case of $\underline{\mathrm{GT}}_{(ell)}$ extend *mutatis* to their profinite, pro-l and prounipotent versions. ☐

Remark 36. We have compatible group morphisms $\underline{\mathrm{GT}} \to \mathrm{Aut}(R_{ell})$, $\underline{\mathrm{GT}}_l \to \mathrm{Aut}(R_l^{ell})$, $\underline{\mathrm{GT}}(\mathbf{k}) \to \mathrm{Aut}(R_{ell}(\mathbf{k}))$ (where $R_l^{ell} = \mathrm{Ker}(\underline{\mathrm{GT}}_l^{ell} \to \underline{\mathrm{GT}}_l)$), defined by $(\lambda, f) \mapsto \theta_{\lambda,f} :=$ conjugation by the image of $(\lambda, f) \mapsto (\lambda, f, g_\pm)$ from Proposition 35. They are such that $\theta_{\lambda,f}(\Psi_+) = \Psi_+^\lambda$ and $\theta_{\lambda,f}((\Psi_+\Psi_-)^3) = ((\Psi_+\Psi_-)^3)^\lambda$, where $(\Psi_+\Psi_-)^3$ is a generator of $Z(\widetilde{\mathrm{SL}_2\,\mathbb{Z}}) = Z(B_3) = \mathbb{Z}$. □

The semigroup scheme morphism from Proposition 35 restricts to a group scheme morphis, which yields an action of $\mathrm{GT}(-)$ on $R_{ell}(-)$. The group scheme $\mathrm{GT}_{ell}(-)$ has then a semidirect product structure, fitting in the diagram

$$\begin{tikzcd}
\mathrm{GRT}_{ell}(-) \arrow[r, phantom, "\simeq"] \arrow[d] & R_{ell}(-) \rtimes \mathrm{GRT}(-) \arrow[d] \\
\mathrm{GL}_2 \arrow[r, phantom, "\simeq"] & \mathrm{SL}_2(-) \rtimes \mathbb{G}_m
\end{tikzcd}$$

where the bottom map is induced by $\mathbb{G}_m \to \mathrm{GL}_2$, $\lambda \mapsto \left(\begin{smallmatrix}\lambda & 0\\ 0 & 1\end{smallmatrix}\right)$.

5. Elliptic associators

5.1. **Lie algebras $\mathfrak{t}_n$ and $\mathfrak{t}_{1,n}$.** Let $\mathbf{k}$ be a $\mathbb{Q}$-ring. If S is a finite set, we define $\mathfrak{t}_S^{\mathbf{k}}$ as the $\mathbf{k}$-Lie algebra with generators t_{ij}, $i \neq j \in S$ and relations $t_{ji} = t_{ij}$, $[t_{ij}, t_{ik} + t_{jk}] = 0$ for i, j, k distinct, $[t_{ij}, t_{kl}] = 0$ for i, j, k, l distinct. We define $\hat{\mathfrak{t}}_S^{\mathbf{k}}$ as its degree completion, where $\deg(t_{ij}) = 1$.

For $S' \supset D_\phi \xrightarrow{\phi} S$ a partially defined map, there is a unique Lie algebra morphism $\mathfrak{t}_S^{\mathbf{k}} \to \mathfrak{t}_{S'}^{\mathbf{k}}$, $x \mapsto x^\phi$, defined by $(t_{ij})^\phi := \sum_{i' \in \phi^{-1}(i), j' \in \phi^{-1}(j)} t_{i'j'}$. Then $S \mapsto \mathfrak{t}_S^{\mathbf{k}}$ is a contravariant functor (finite sets, partially defined maps) $\to$ {Lie algebras}.

We also define $\mathfrak{t}_{1,S}^{\mathbf{k}}$ as the $\mathbf{k}$-Lie algebra with generators $x_i^\pm$, $i \in S$ and relations $\sum_{i \in S} x_i^\pm = 0$, $[x_i^\pm, x_j^\pm] = 0$ for $i \neq j$, $[x_i^+, x_j^-] = [x_j^+, x_i^-]$ for $i \neq j$, $[x_k^\pm, [x_i^+, x_j^-]] = 0$ for i, j, k distinct. We then have a Lie algebra morphism $\mathfrak{t}_S^{\mathbf{k}} \to \mathfrak{t}_{1,S}^{\mathbf{k}}$, $t_{ij} \mapsto [x_i^+, x_j^-]$, which we denote by $x \mapsto \{x\}$. We will also write $t_{ij} = [x_i^+, x_j^-]$. We define $\hat{\mathfrak{t}}_{1,S}^{\mathbf{k}}$ as the degree completion of $\mathfrak{t}_{1,S}^{\mathbf{k}}$, where $\deg(x_i^\pm) = 1$.

For $S' \xrightarrow{\phi} S$ a map, there is a unique Lie algebra morphism $\mathfrak{t}_{1,S}^{\mathbf{k}} \to \mathfrak{t}_{1,S'}^{\mathbf{k}}$, $x \mapsto x^\phi$, such that $(x_i^\pm)^\phi := \sum_{i' \in \phi^{-1}(i)} x_{i'}^\pm$. Then $S \mapsto \mathfrak{t}_{1,S}$ is a contravariant functor (finite sets, maps) $\to$ {Lie algebras}. By restriction, $S \mapsto \mathfrak{t}_S$ may be viewed as a contravariant functor of the same type, and the morphism $\mathfrak{t}_S \to \mathfrak{t}_{1,S}$ is then functorial; i.e., we have $\{x\}^\phi = \{x^\phi\}$ for $x \in \mathfrak{t}_S$ and any map $S' \xrightarrow{\phi} S$.

We set $\mathfrak{t}_n^{\mathbf{k}} := \mathfrak{t}_{\{1,\ldots,n\}}^{\mathbf{k}}$, $\mathfrak{t}_{1,n} := \mathfrak{t}_{1,\{1,\ldots,n\}}^{\mathbf{k}}$ and we write x^ϕ as $x^{I_1,\ldots,I_n}$, where $I_i = \phi^{-1}(i)$ for $x \in \mathfrak{t}_n^{\mathbf{k}}$ or $x \in \mathfrak{t}_{1,n}^{\mathbf{k}}$.

5.2. **Elliptic associators.** Recall that the set $\underline{M}(\mathbf{k})$ of associators defined over $\mathbf{k}$ is the set of $(\mu, \Phi) \in \mathbf{k} \times \exp(\hat{\mathfrak{f}}_2^{\mathbf{k}})$, such that $\Phi^{3,2,1} = \Phi^{-1}$,

$$e^{\mu t_{23}/2}\Phi^{1,2,3}e^{\mu t_{12}/2}\Phi^{3,1,2}e^{\mu t_{31}/2}\Phi^{2,3,1} = e^{\mu(t_{12}+t_{13}+t_{23})/2}, \tag{20}$$

$$\Phi^{2,3,4}\Phi^{1,23,4}\Phi^{1,2,3} = \Phi^{1,2,34}\Phi^{12,3,4}, \tag{21}$$

where we use the inclusion $\hat{\mathfrak{f}}_2^{\mathbf{k}} \subset \hat{\mathfrak{t}}_3^{\mathbf{k}}$, $A, B \mapsto t_{12}, t_{23}$.

Definition 37. *The set $\underline{Ell}(\mathbf{k})$ of elliptic associators defined over $\mathbf{k}$ is the set of quadruples $(\mu, \Phi, A_\pm)$, where $(\mu, \Phi) \in \underline{M}(\mathbf{k})$ and $A_\pm \in \exp(\hat{\mathfrak{t}}_{1,2}^{\mathbf{k}})$, such that:*

$$\alpha_\pm^{3,1,2}\alpha_\pm^{2,3,1}\alpha_\pm^{1,2,3} = 1, \text{ where } \alpha_\pm = \{e^{\pm\mu(t_{12}+t_{13})/2}\}A_\pm^{1,23}\{\Phi^{1,2,3}\}, \tag{22}$$

$$\{e^{\mu t_{12}}\} = \big(\{\Phi\}^{-1}A_-^{1,23}\{\Phi\}, \{e^{-\mu t_{12}/2}(\Phi^{2,1,3})^{-1}\}(A_+^{2,13})^{-1}\{\Phi^{2,1,3}e^{-\mu t_{12}/2}\}\big). \tag{23}$$

Remark 38. We then have $\{e^{\pm\mu t_{12}/2}\}A_\pm^{2,1}\{e^{\pm\mu t_{12}/2}\}A_\pm^{1,2}=1$ and $\{e^{\mu t_{12}}\}=(A_-,A_+)$.

Then $\mathbf{k}\mapsto \underline{M}(\mathbf{k}),\underline{Ell}(\mathbf{k})$ are functors $\{\mathbb{Q}\text{-rings}\}\to\{\text{sets}\}$, i.e., $\mathbb{Q}$-schemes. We have an obvious scheme morphism $\underline{Ell}\to\underline{M}$, $(\mu,\Phi,A_\pm)\mapsto(\mu,\Phi)$.

Define also a scheme morphism $\underline{Ell}\to \mathrm{M}_2$ by $(\mu,\Phi,A_\pm)\mapsto \left(\begin{smallmatrix} u_+ & v_+ \\ u_- & v_-\end{smallmatrix}\right)$, where $\log A_\pm\equiv u_\pm x_1^+ + v_\pm x_1^-$ mod $[\hat{\mathfrak{t}}_{1,2},\hat{\mathfrak{t}}_{1,2}]$. Then the diagram

$$\begin{tikzcd}
\underline{Ell} \arrow[r] \arrow[d] & \underline{M} \arrow[d] \\
\mathrm{M}_2 \arrow[r, "\det"] & \mathbb{A}
\end{tikzcd}$$

commutes in view of (23).

5.3. **Categorical interpretations.**

Definition 39. *(see* [Dr]*) A infinitesimally braided monoidal category (IBMC) over* $\mathbf{k}$ *is a set* $(\mathcal{C}_0,\otimes,c_{-,-},a_{-,-,-},U_-,t_{-,-})$*, such that:*

1) $(\mathcal{C}_0,\otimes,c_{-,-},a_{-,-,-})$ *is a symmetric monoidal category (i.e.,* $c_{Y,X}c_{X,Y}=\mathrm{id}_{X\otimes Y}$*);*

2) $\mathrm{Ob}\,\mathcal{C}_0\ni X\mapsto U_X\lhd\mathrm{Aut}_{\mathcal{C}_0}(X)$ *is such that* U_X *is a* $\mathbf{k}$*-prounipotent group, and* $iU_Xi^{-1}=U_Y$ *for any* $i\in\mathrm{Iso}_{\mathcal{C}_0}(X,Y)$*;*

3) $(\mathrm{Ob}\,\mathcal{C}_0)^2\ni(X,Y)\mapsto t_{X,Y}\in\mathrm{Lie}\,U_{X\otimes Y}$ *is a natural assignment;*

4) $t_{Y,X}=c_{X,Y}t_{X,Y}c_{X,Y}^{-1}$ *and*

$$t_{X\otimes Y,Z}=a_{X,Y,Z}(\mathrm{id}_X\otimes t_{Y,Z})a_{X,Y,Z}^{-1}+(c_{Y,X}\otimes\mathrm{id}_Z)a_{Y,X,Z}(\mathrm{id}_Y\otimes t_{X,Z})((c_{Y,X}\otimes\mathrm{id}_Z)a_{Y,X,Z})^{-1}.$$

A functor $f:\mathcal{C}_0\to\mathcal{C}_0'$ between IBMCs is then a tensor functor, such that $f(U_X)\subset U'_{f(X)}$ and $f(t_{X,Y})=t'_{f(X),f(Y)}$. An example of IMBC is constructed as follows: $\mathcal{C}_0=\mathbf{PaCD}$ is the category with the same objects as **PaB**, $\mathbf{PaCD}(O,O'):=\begin{cases}\exp(\hat{\mathfrak{t}}_{|O|})\rtimes S_{|O|} & \text{if } |O|=|O'| \\ \emptyset & \text{otherwise}\end{cases}$, $c_{O,O'}=s_{n,n'}\in S_{n+n'}\subset\mathrm{Aut}_{\mathbf{PaCD}}(O\otimes O')$ is the permutation $i\mapsto i+n'$ for $i\in[1,n]$, $i\mapsto i-n$ for $i\in[n+1,n+n']$, $a_{O,O',O''}:=1$, $U_O=\exp(\hat{\mathfrak{t}}_n^{\mathbf{k}})\lhd\mathrm{Aut}_{\mathcal{C}_0}(O)$, $t_{O,O'}:=\sum_{i=1}^n\sum_{i'=n+1}^{n+n'}t_{ii'}$. The pair $(\mathbf{PaCD},\bullet)$ is initial among pairs (an IBMC, a distinguished object).

We then set:

Definition 40. *An elliptic structure over the IBMC* $\mathcal{C}_0$ *is a set* $(\mathcal{C},F,\tilde{U}_-,x^\pm_{-,-})$*, where* $\mathcal{C}$ *is a category,* $F:\mathcal{C}_0\to\mathcal{C}$ *is a functor inducing the identity on objects,* $\mathrm{Ob}\,\mathcal{C}_0\ni X\mapsto\tilde{U}_X\lhd\mathrm{Aut}_{\mathcal{C}}(X)$ *is the assignment of a* $\mathbf{k}$*-prounipotent group, where* $\tilde{i}\tilde{U}_X\tilde{i}^{-1}=\tilde{U}_Y$ *for* $\tilde{i}\in\mathrm{Iso}_{\mathcal{C}}(X,Y)$ *and* $F(U_X)\subset\tilde{U}_X$*, and* $(\mathrm{Ob}\,\mathcal{C}_0)^2\ni(X,Y)\mapsto x^\pm_{X,Y}\in\mathrm{Lie}\,\tilde{U}_{X\otimes Y}$ *is a natural assignment, such that*

$$x^\pm_{Y,X}=c_{X,Y}x^\pm_{X,Y}c_{X,Y}^{-1},\quad x^\pm_{X,\mathbf{1}}=0,$$

$$x^\pm_{X\otimes Y,Z}+(c_{X,Y\otimes Z}a_{X,Y,Z})^{-1}x^\pm_{Y\otimes Z,X}c_{X,Y\otimes Z}a_{X,Y,Z}+(a_{Z,X,Y}^{-1}c_{X\otimes Y,Z})^{-1}x^\pm_{Z\otimes X,Y}a_{Z,X,Y}^{-1}c_{X\otimes Y,Z}=0,$$

$$F(t_{X,Y}\otimes\mathrm{id}_Z)=[a_{X,Y,Z}^{-1}x^+_{X,Y\otimes Z}a_{X,Y,Z},(c_{X,Y}\otimes\mathrm{id}_Z)^{-1}a_{Y,X,Z}x^-_{Y,X\otimes Z}a_{Y,X,Z}^{-1}(c_{X,Y}\otimes\mathrm{id}_Z)].$$

Functors between pairs (an IBMC, an elliptic structure over it) are defined in the obvious way. An elliptic structure over **PaCD** is defined as follows: $\mathcal{C}:=\mathbf{PaCD}_{ell}$ is the category with the same objects as **PaB**, $\mathbf{PaCD}_{ell}(O,O'):=\begin{cases}\exp(\hat{\mathfrak{t}}^{\mathbf{k}}_{1,|O|})\rtimes S_{|O|} & \text{if } |O|=|O'| \\ \emptyset & \text{otherwise,}\end{cases}$ $\tilde{U}_O=\exp(\hat{\mathfrak{t}}_{1,|O|})\lhd\mathrm{Aut}_{\mathbf{PaCD}_{ell}}(O)$, F is induced by $\mathfrak{t}_n\to\mathfrak{t}_{1,n}$, $x\mapsto\{x\}$ and the identity of symmetric groups, $x^\pm_{O,O'}=\sum_{i=1}^{|O|}x_i^\pm\in\mathrm{Lie}\,\tilde{U}_{O\otimes O'}$. The triple $(\mathbf{PaCD},\mathbf{PaCD}_{ell},\bullet)$ is universal for triples (an IBMC, an elliptic structure over it, a distinguished object).

Let us say that a $\mathbf{k}$-BMC is a braided monoidal category (BMC) $\mathcal{C}_0$, such that the image of each morphism $P_n\to\mathrm{Aut}_{\mathcal{C}_0}(X_1\otimes\cdots\otimes X_n)$ is contained in a $\mathbf{k}$-prounipotent group. Then each

$(\mu, \Phi) \in \underline{M}(\mathbf{k})$ gives rise to a map $\{\text{IBMCs}\} \to \{\mathbf{k}\text{-BMCs}\}$, $\mathcal{C}_0 \mapsto (\mu, \Phi) * \mathcal{C}_0$, where $(\mu, \Phi) * \mathcal{C}_0 = (\mathcal{C}_0, \otimes, \beta_{X,Y} := c_{X,Y} e^{\mu t_{X,Y}/2}, \tilde{a}_{X,Y,Z} := \Phi(a_{X,Y,Z}(t_{X,Y} \otimes \mathrm{id}_Z) a^{-1}_{X,Y,Z}, \mathrm{id}_X \otimes t_{Y,Z}) a_{X,Y,Z})$.

In the same say, a $\mathbf{k}$-elliptic structure over a $\mathbf{k}$-BMC is an elliptic structure, such that the image of each morphism $P_{1,n} \to \mathrm{Aut}_{\mathcal{C}}(X_1 \otimes \cdots \otimes X_n)$ is contained in a $\mathbf{k}$-prounipotent group. Then each $(\mu, \Phi, A_\pm) \in \underline{Ell}(\mathbf{k})$ gives rise to a map $\{$(an IBMC, an elliptic structure over it)$\} \to \{$(a $\mathbf{k}$-BMC, an elliptic structure over it)$\}$, $(\mathcal{C}_0, \mathcal{C}) \mapsto (\mu, \Phi, A_\pm) * (\mathcal{C}_0, \mathcal{C}) = (\tilde{\mathcal{C}_0}, \tilde{\mathcal{C}})$, where $\tilde{\mathcal{C}_0} = (\mu, \Phi) * \mathcal{C}_0$ and $\tilde{\mathcal{C}} = (\mathcal{C}, F, \tilde{A}^\pm_{X,Y} := A_\pm(x^+_{X,Y}, x^-_{X,Y}))$.

5.4. **Action of $\underline{\mathrm{GT}}_{ell}(-)$ on $\underline{Ell}$.** Recall first that we have an action of $\underline{\mathrm{GT}}(\mathbf{k})$ on $\underline{M}(\mathbf{k})$, defined by

$$(\lambda, f) * (\mu, \Phi) := (\lambda\mu, \Phi(A, B) f(e^{\mu A}, \Phi(A, B)^{-1} e^{\mu B} \Phi(A, B))) = (\mu', \Phi').$$

For $(\lambda, f, g_\pm) \in \underline{\mathrm{GT}}_{ell}(\mathbf{k})$ and $(\mu, \Phi, A_\pm) \in \underline{Ell}(\mathbf{k})$, we set

$$(\lambda, f, g_\pm) * (\mu, \Phi, A_\pm) := (\mu', \Phi', A'_\pm)$$

where $A'_\pm := g_\pm(A_+, A_-)$.

Proposition 41. *This defines an action of* $\underline{\mathrm{GT}}_{ell}(\mathbf{k})$ *on* $\underline{Ell}(\mathbf{k})$.

Proof. For $g_{ell} \in \underline{\mathrm{GT}}_{ell}(\mathbf{k})$, and $(\mathcal{C}_0, \mathcal{C}) \in \{$(a $\mathbf{k}$-BMC, an elliptic structure over it)$\}$, we have $g_{ell} * ((\mu, \Phi, A_\pm) * (\mathcal{C}_0, \mathcal{C})) = (g_{ell} * (\mu, \Phi, A_\pm)) * (\mathcal{C}_0, \mathcal{C})$. When $(\mathcal{C}_0, \mathcal{C}) = (\mathbf{PaCD}, \mathbf{PaCD}_{ell})$, $(\mu, \Phi, A_\pm)$ can be recovered uniquely from $(\mu, \Phi, A_\pm) * (\mathcal{C}_0, \mathcal{C})$, as $e^{\mu t_{12}} = \beta^2_{\bullet,\bullet}$, $\Phi = \tilde{a}_{\bullet,\bullet,\bullet}$, and $A_\pm = A^\pm_{\bullet,\bullet}$, which implies that the above formula defines an action. □

Remark 42. The actions of $\underline{\mathrm{GT}}(\mathbf{k})$ on $\{\mathbf{k}\text{-BMCs}\}$ and on $\underline{M}(\mathbf{k})$ are compatible, in the sense that for $g \in \underline{\mathrm{GT}}(\mathbf{k})$, $g * ((\mu, \Phi) * \mathcal{C}_0) = (g * (\mu, \Phi)) * \mathcal{C}_0$. In the same way, the action of $\underline{\mathrm{GT}}_{ell}(\mathbf{k})$ on $\{$(a $\mathbf{k}$-BMC, an elliptic structure over it)$\}$ and on $\underline{Ell}(\mathbf{k})$ are compatible.

Remark 43. The morphism $\underline{Ell} \to \mathrm{M}_2$ from Section 5.2 is compatible with the semigroup scheme morphism $\underline{\mathrm{GT}}_{ell}(-) \to \mathrm{M}_2$ from Proposition 26, with the action of $\underline{\mathrm{GT}}_{ell}$ on $\underline{Ell}$, and with the left multiplication action of M_2 on itself.

5.5. **A morphism $\underline{M} \to \underline{Ell}$.** The scheme morphism $\underline{Ell} \to \underline{M}$, $(\mu, \Phi, A_\pm) \to (\mu, \Phi)$ is clearly compatible with the semigroup scheme morphism $\underline{\mathrm{GT}}_{ell}(-) \to \underline{\mathrm{GT}}(-)$. We now construct a section of this morphism.

Proposition 44. *There is a unique scheme morphism $\sigma : \underline{M} \to \underline{Ell}$, $(\mu, \Phi) \to (\mu, \Phi, A_\pm)$, where*

$$A_+ = \Phi(\frac{\mathrm{ad}\, x_1}{e^{\mathrm{ad}\, x_1} - 1}(y_2), t_{12}) \cdot e^{\mu \frac{\mathrm{ad}\, x_1}{e^{\mathrm{ad}\, x_1} - 1}(y_2)} \cdot \Phi(\frac{\mathrm{ad}\, x_1}{e^{\mathrm{ad}\, x_1} - 1}(y_2), t_{12})^{-1},$$

$$A_- = e^{\mu t_{12}/2} \Phi(\frac{\mathrm{ad}\, x_2}{e^{\mathrm{ad}\, x_2} - 1}(y_1), t_{21}) e^{x_1} \Phi(\frac{\mathrm{ad}\, x_1}{e^{\mathrm{ad}\, x_1} - 1}(y_2), t_{12})^{-1}$$

(we set $x_i := x_i^+, y_i := x_i^-$). It is compatible with the semigroup scheme morphism $\underline{\mathrm{GT}}(-) \to \underline{\mathrm{GT}}_{ell}(-)$ from Proposition 35.

One checks that σ fits in a diagram

$$\begin{tikzcd}
\underline{M} \arrow[r, "\sigma"] \arrow[d] & \underline{Ell} \arrow[d] \\
\mathbb{A} \arrow[r] & \mathrm{M}_2
\end{tikzcd}$$

where the bottom map is $\mu \mapsto \begin{pmatrix} 0 & -\mu \\ 1 & 0 \end{pmatrix}$, compatible with the last diagram of Proposition 35.

Proof. By [CEE], Prop. 5.3, $(\mu, \Phi, A_\pm)$ satisfies

$$A_\pm^{12,3} = \{e^{\pm\mu t_{12}/2} (\Phi^{-1})^{2,1,3}\} A_\pm^{2,13} \{\Phi^{2,1,3} e^{\pm\mu t_{12}/2} \Phi^{-1}\} A_\pm^{1,23} \{\Phi\},$$

and therefore (22).

The last identity of *loc. cit.* can be rewritten as follows (using the commutation of $\{t_{12}\}$ with $A_+^{12,3}$)

$$A_-^{2,13}\{\Phi^{2,1,3}\}A_+^{12,3}\{(\Phi^{2,1,3})^{-1}\}(A_-^{2,13})^{-1} = \{(\Phi^{3,1,2})^{-1}e^{\mu t_{12}/2}\Phi^{3,2,1}e^{\mu t_{23}}\Phi^{1,2,3}e^{-\mu t_{12}/2}\}A_+^{12,3}\{\Phi^{3,1,2}\}.$$

Now the hexagon and duality identities imply

$$(\Phi^{3,2,1})^{-1}e^{\mu t_{12}/2}\Phi^{3,2,1}e^{\mu t_{23}}\Phi^{1,2,3}e^{-\mu t_{12}/2} = e^{-\mu t_{13}/2}\Phi^{2,3,1}e^{\mu t_{23}/2}(\Phi^{3,2,1})^{-1}e^{\mu t_{3,12}/2}, \tag{24}$$

$$\Phi^{3,1,2} = e^{\mu t_{3,21}/2}\Phi^{3,2,1}e^{-\mu t_{23}/2}(\Phi^{2,3,1})^{-1}e^{-\mu t_{13}/2},$$

and

$$\Phi^{2,1,3} = e^{\mp\mu t_{13}/2}\Phi^{2,3,1}e^{\mp\mu t_{23}/2}(\Phi^{3,2,1})^{-1}e^{\pm\mu t_{3,12}/2},$$

so (24) is rewritten (using the commutation of $\{t_{13}\}$ with $A_-^{2,13}$)

$$\begin{aligned}&\{e^{-\mu t_{13}/2}\}A_-^{2,13}\{\Phi^{2,3,1}e^{-\mu t_{23}/2}(\Phi^{3,2,1})^{-1}e^{\mu t_{3,12}/2}\}A_+^{12,3}\{e^{\mu t_{3,12}/2}\Phi^{3,2,1}e^{-\mu t_{23}/2}(\Phi^{2,3,1})^{-1}\}\\ &\qquad (A_-^{2,13})^{-1}\{e^{-\mu t_{13}/2}\}\\ &= \{e^{-\mu t_{13}/2}\Phi^{2,3,1}e^{\mu t_{23}/2}(\Phi^{3,2,1})^{-1}e^{\mu t_{3,21}/2}\}A_+^{12,3}\{e^{\mu t_{3,21}/2}\Phi^{3,2,1}e^{-\mu t_{23}/2}(\Phi^{2,3,1})^{-1}e^{-\mu t_{13}/2}\}.\end{aligned} \tag{25}$$

As $A_+^{2,1}e^{\mu t_{12}/2}A_+e^{\mu t_{12}/2} = 1$, we have $e^{\mu t_{3,12}/2}A_+^{12,3}e^{\mu t_{3,12}/2} = (A_+^{3,12})^{-1}$; using this identity and performing the transformation of indices $(1,2,3) \to (3,1,2)$, (25) yields (23). So $(\mu, \Phi, A_\pm) \in \underline{Ell}(\mathbf{k})$. The compatibility of $\sigma : \underline{M}(\mathbf{k}) \to \underline{Ell}(\mathbf{k})$ with the semigroup morphism $\underline{\mathrm{GT}}(\mathbf{k}) \to \underline{\mathrm{GT}}_{ell}(\mathbf{k})$ follows from $(A_-, A_+) = e^{\mu t_{12}}$. ☐

5.6. **A subscheme** $Ell \subset \underline{Ell}$ **and its torsor structure under** $\mathrm{GT}_{ell}(-)$**.** Set $M(\mathbf{k}) := \{(\mu, \Phi) | \mu \in \mathbf{k}^\times\} \subset \underline{M}(\mathbf{k})$ and $Ell(\mathbf{k}) := \{(\mu, \Phi, A_\pm) | \mu \in \mathbf{k}^\times\} \subset \underline{Ell}(\mathbf{k})$. The actions of $\underline{\mathrm{GT}}_{(ell)}$ restrict to actions of $\mathrm{GT}(\mathbf{k})$ on $M(\mathbf{k})$ and $\mathrm{GT}_{ell}(\mathbf{k})$ on $Ell(\mathbf{k})$. Recall that $M(\mathbf{k})$ is a principal homogeneous space under the action of $\mathrm{GT}(\mathbf{k})$ ([Dr]). Similarly:

Proposition 45. *$Ell(\mathbf{k})$ is empty or a principal homogeneous space under the action of* $\mathrm{GT}_{ell}(\mathbf{k})$.

Proof. Assume $Ell(\mathbf{k}) \neq \emptyset$ and let us show that the action is free. If $(\lambda, f, g_\pm) \in \mathrm{Stab}(\mu, \Phi, A_\pm)$, then by the freeness of the action of $\mathrm{GT}(\mathbf{k})$ on $M(\mathbf{k})$, $(\lambda, f) = 1$. Then $A_\pm = g_\pm(A_+, A_-)$. Relation (23) implies that if $a_\pm, b_\pm \in \mathbf{k}$ are such that $\log A_\pm \equiv a_\pm x_1^+ + b_\pm x_1^-$ mod degree ≥ 2 (where $x_1^\pm$ have degree 1), then $a_+b_- - a_-b_+ = \mu$, which implies that $(\log A_+, \log A_-)$ generate $\hat{\mathfrak{t}}_{1,2}^{\mathbf{k}}$, and therefore that $g_\pm = 1$.

We now prove that the action is transitive. As the action of $\mathrm{GT}(\mathbf{k})$ on $M(\mathbf{k})$ is transitive, and as $\mathrm{GT}_{ell}(\mathbf{k}) \to \mathrm{GT}(\mathbf{k})$ is surjective (as the morphism defined in Proposition 35 restricts to a section of it), it suffices to prove that for any $(\mu, \Phi) \in Ell(\mathbf{k})$, the action of $R_{ell}(\mathbf{k})$ on $\{(A_+, A_-) | (\mu, \Phi, A_\pm) \in Ell(\mathbf{k})\}$ is transitive. If (A_+, A_-) and (A'_+, A'_-) belong to this set, we have a unique $(g_+, g_-) \in F_2(\mathbf{k})^2 \simeq P_{1,2}(\mathbf{k})^2$ such that $A'_\pm = g_\pm(A_+, A_-)$. Then

$$\alpha_\pm^{1,2,3}\alpha_\pm^{3,1,2}\alpha_\pm^{2,3,1} = 1, \text{ where } \alpha_\pm = g_\pm(A_+^{1,23}, A_-^{1,23})\{\Phi^{1,2,3}e^{\pm\mu t_{12,3}/2}\}.$$

The canonical morphism $B_{1,3} \to \mathrm{Aut}_{(\mu,\Phi,A_\pm)*\mathbf{PaCD}}(\bullet(\bullet\bullet)) = \exp(\hat{\mathfrak{t}}_{1,3}^{\mathbf{k}}) \rtimes S_3$ extends to an isomorphism $B_{1,3}(\mathbf{k}) \simeq \exp(\hat{\mathfrak{t}}_{1,3}^{\mathbf{k}}) \rtimes S_3$, given by $X_1^\pm \mapsto A_\pm^{1,23}$, $\sigma_1 \mapsto \{\Phi e^{\mu t_{12}/2}\}(12)\{\Phi\}^{-1}$, $\sigma_2 \mapsto \{e^{\mu t_{23}/2}\}(23)$. It is such that $\sigma_2^{\pm 1}\sigma_1^{\pm 1} \mapsto \{\Phi e^{\pm(\mu/2)t_{3,12}}\}(23)(12)$. The preimage of the above identity by this isomorphism then yields $(g_\pm(X_1^+, X_1^-)\sigma_2^{\pm 1}\sigma_1^{\pm 1})^3 = 1$. Similarly, the preimage of the identity

$$\{e^{\mu t_{12}}\} = \big(\{\Phi^{-1}\}g_-(A_+^{1,23}, A_-^{1,23})\{\Phi\}, \{e^{-(\mu/2)t_{12}(\Phi^{2,1,3})^{-1}}\}g_+^{-1}(A_+^{2,13}, A_-^{2,13})\{\Phi^{2,1,3}e^{-(\mu/2)t_{12}}\}\big)$$

yields $\sigma_1^2 = (\sigma_1 g_+^{-1}(X_1^+, X_1^-)\sigma_1, g_-(X_1^+, X_1^-))$. ☐

Recall the following definition:

Definition 46. *A $\mathbb{Q}$-torsor is the data of: $\mathbb{Q}$-group schemes G, H, a $\mathbb{Q}$-scheme X, commuting left and right actions of G, H on X, such that: for any $\mathbf{k}$ with $X(\mathbf{k}) \neq \emptyset$, the action of $G(\mathbf{k})$ and $H(\mathbf{k})$ on $X(\mathbf{k})$ is free and transitive.*

Morphisms of torsors are then defined in the obvious way.

The above $\mathbb{Q}$-scheme morphisms between $\underline{Ell}$ and $\underline{M}$ restrict to a torsor morphism $Ell \to M$ and a section of it $M \overset{\sigma}{\to} Ell$, fitting in commutative diagrams

$$\begin{tikzcd} Ell \arrow[r] \arrow[d] & M \arrow[d] \\ \mathrm{GL}_2 \arrow[r, "\det"] & \mathbb{G}_m \end{tikzcd} \text{ and } \begin{tikzcd} M \arrow[r] \arrow[d] & Ell \arrow[d] \\ \mathbb{G}_m \arrow[r, "\mu \mapsto \begin{pmatrix} 0 & -1 \\ \mu & 0 \end{pmatrix}"] & \mathrm{GL}_2 \end{tikzcd}$$

6. The group $\mathrm{GRT}_{ell}(\mathbf{k})$

6.1. **Reminders about** $\mathrm{GRT}(\mathbf{k})$**.** Let again $\mathbf{k}$ be a $\mathbb{Q}$-ring. Recall ([Dr]) that $\mathrm{GRT}_1(\mathbf{k})$ is defined as the set of all $g \in \exp(\hat{\mathfrak{f}}_2^{\mathbf{k}}) \subset \exp(\hat{\mathfrak{t}}_3^{\mathbf{k}})$, such that:

$$g^{3,2,1} = g^{-1}, \quad g^{3,1,2} g^{2,3,1} g^{1,2,3} = 1, \quad t_{12} + \mathrm{Ad}(g^{1,2,3})^{-1}(t_{23}) + \mathrm{Ad}(g^{2,1,3})^{-1}(t_{13}) = t_{12} + t_{13} + t_{23},$$
$$g^{2,3,4} g^{1,23,4} g^{1,2,3} = g^{1,2,34} g^{12,3,4}.$$

This is a group with law $(g_1 * g_2)(A,B) := g_1(\mathrm{Ad}(g_2(A,B))(A), B) g_2(A,B)$. Note that $g \in \mathrm{GRT}_1(\mathbf{k})$ gives rise to $\theta_g \in \mathrm{Aut}(\hat{\mathfrak{t}}_3^{\mathbf{k}})$, defined by

$$\theta_g : t_{12} \mapsto t_{12}, \quad t_{23} \mapsto \mathrm{Ad}(g^{1,2,3})^{-1}(t_{23}), \quad t_{13} \mapsto \mathrm{Ad}(g^{2,1,3})^{-1}(t_{13}).$$

Then $g_1 * g_2 = g_1 \theta_{g_2}(g_1)$, and $\theta_{g_1 * g_2} = \theta_{g_2} \theta_{g_1}$, so $g \mapsto \theta_g$ is a group antimorphism.

The group $\mathbf{k}^\times$ acts on $\mathrm{GRT}_1(\mathbf{k})$ by $(c \cdot g)(A,B) := g(c^{-1}A, c^{-1}B)$, and one sets $\mathrm{GRT}(\mathbf{k}) := \mathrm{GRT}_1(\mathbf{k}) \rtimes \mathbf{k}^\times$. $\mathrm{GRT}_1(-)$ is a prounipotent group scheme.

6.2. **The group** $\mathrm{GRT}_{ell}(\mathbf{k})$**.** Define $\mathrm{GRT}_1^{ell}(\mathbf{k})$ as the set of all $(g, u_\pm)$, such that $g \in \mathrm{GRT}_1(\mathbf{k})$, $u_\pm \in \hat{\mathfrak{t}}_{1,2}^{\mathbf{k}}$, and

$$\mathrm{Ad}(g^{1,2,3})^{-1}(u_\pm^{1,23}) + \mathrm{Ad}(g^{2,1,3})^{-1}(u_\pm^{2,13}) + u_\pm^{3,12} = 0, \tag{26}$$

$$[\mathrm{Ad}(g^{1,2,3})^{-1}(u_\pm^{1,23}), u_\pm^{3,12}] = 0, \tag{27}$$

$$[\mathrm{Ad}(g^{1,2,3})^{-1}(u_+^{1,23}), \mathrm{Ad}(g^{2,1,3})^{-1}(u_-^{2,13})] = t_{12}. \tag{28}$$

Set $(g_1, u_\pm^1) * (g_2, u_\pm^2) := (g, u_\pm)$, where

$$u_\pm(x_1, y_1) := u_\pm^1(u_+^2(x_1,y_1), u_-^2(x_1,y_1)) \tag{29}$$

(where $\mathfrak{t}_{1,2}^{\mathbf{k}}$ is viewed as the free Lie algebra generated by x_1, y_1).

We first prove:

Lemma 47. $(g, u_\pm) \in \mathrm{GRT}_1^{ell}(\mathbf{k})$ *iff there exists an automorphism of $\hat{\mathfrak{t}}_{1,3}^{\mathbf{k}}$ (henceforth denoted $\theta_{g,u_\pm}$), such that*

$$x_1^\pm \mapsto \mathrm{Ad}(g^{1,2,3})^{-1}(u_\pm^{1,23}), \quad x_2^\pm \mapsto \mathrm{Ad}(g^{2,1,3})^{-1}(u_\pm^{2,13}), \quad x_3^\pm \mapsto u_\pm^{3,12},$$
$$t_{12} \mapsto t_{12}, \quad t_{23} \mapsto \mathrm{Ad}(g^{1,2,3})^{-1}(t_{23}), \quad t_{13} \mapsto \mathrm{Ad}(g^{2,1,3})^{-1}(t_{13}).$$

Proof. The condition that the relations $x_1^\pm + x_2^\pm + x_3^\pm = 0$ (resp., $[x_1^\pm, x_3^\pm] = 0$, $[x_1^+, x_2^-] = t_{12}$) are preserved is equivalent to condition (26) (resp., (27), (28)), and the relation $[t_{12}, x_3^\pm] = 0$ is automatically preserved. Then the relation $g^{3,1,2} g^{2,3,1} g^{1,2,3} = 1$ implies that $\theta_{g,u_\pm}(x^{2,3,1}) = \mathrm{Ad}(g^{1,2,3})^{-1}(\theta_{g,u_\pm}(x)^{2,3,1})$ for $x \in \{x_i^\pm, t_{ij}\}$. So the other relations $[t_{ij}, x_k^\pm] = 0$ are also preserved. □

Proposition 48. $\mathrm{GRT}_1^{ell}(\mathbf{k})$, *equipped with the above product, is a group.*

Proof. The product is that of the group $\mathrm{GRT}_1(\mathbf{k}) \times \mathrm{Aut}(\hat{\mathfrak{t}}_{1,2}^{\mathbf{k}})^{op}$, so it remains to prove that $\mathrm{GRT}_1^{ell}(\mathbf{k})$ is stable under the operations of product and inverse. If now $(g_i, u_\pm^i) \in \mathrm{GRT}_1^{ell}(\mathbf{k})$ $(i = 1, 2)$, then the action of $\theta_{g_2,u_2^\pm}\theta_{g_1,u_1^\pm}$ on the generators of $\hat{\mathfrak{t}}_{1,3}^{\mathbf{k}}$ is given by the formulas of Lemma 47, with $g = g_1 * g_2$ and $u_\pm$ as in (29). So $(g, u_\pm) \in \mathrm{GRT}_1^{ell}(\mathbf{k})$, as claimed. Similarly, if $(g, u_\pm) \in \mathrm{GRT}_1^{ell}(\mathbf{k})$, then the action of $\theta_{g,u_\pm}^{-1}$ on the generators of $\hat{\mathfrak{t}}_{1,3}^{\mathbf{k}}$ is as in Lemma 47, with $(g, u_\pm)$ replaced by (inverse of g in $\mathrm{GRT}_1(\mathbf{k})$, inverse of (u_+, u_-) in $\mathrm{Aut}(\hat{\mathfrak{t}}_{1,2}^{\mathbf{k}})$), so $(g, u_\pm)$ is invertible. ☐

In particular, we have

$$\theta_{(g_2,u_2^\pm)}\theta_{(g_1,u_1^\pm)} = \theta_{(g_1,u_1^\pm)*(g_2,u_2^\pm)}. \tag{30}$$

The assignments $\mathbf{k} \mapsto \mathrm{GRT}_1^{(ell)}(\mathbf{k})$ are then $\mathbb{Q}$-group schemes.

For $(g, u_\pm) \in \mathrm{GRT}_1^{ell}(\mathbf{k})$, define $a_\pm, b_\pm$ by $u_\pm(x_1, y_1) = a_\pm x_1 + b_\pm y_1 \mod [\hat{\mathfrak{t}}_{1,2}^{\mathbf{k}}, \hat{\mathfrak{t}}_{1,2}^{\mathbf{k}}]$.

Lemma 49. *1) There is a unique group scheme morphism* $\mathrm{GRT}_1^{ell}(-) \to \mathrm{SL}_2$, $(g, u_\pm) \mapsto \left(\begin{smallmatrix} a_+ & b_+ \\ a_- & b_- \end{smallmatrix}\right)$.

2) This morphism has a section $\mathrm{SL}_2 \to \mathrm{GRT}_1^{ell}(-)$, *given by* $\left(\begin{smallmatrix} a_+ & b_+ \\ a_- & b_- \end{smallmatrix}\right) \mapsto (1, u_\pm(x_1, y_1) = a_\pm x_1 + b_\pm y_1)$.

Proof. 1) $a_+b_- - a_-b_+ = 1$ follows from (28); the morphism property is clear. 2) is straightforward. ☐

We now set $\mathrm{GRT}_{I_2}^{ell}(\mathbf{k}) := \mathrm{Ker}\left(\mathrm{GRT}_1^{ell}(\mathbf{k}) \to \mathrm{SL}_2(\mathbf{k})\right)$. This defines a group scheme $\mathrm{GRT}_{I_2}^{ell}(-)$.

Lemma 50. $\mathrm{GRT}_{I_2}^{ell}(-)$ *is a prounipotent group scheme; we have* $\mathrm{GRT}_1^{ell}(-) = \mathrm{GRT}_{I_2}^{ell}(-) \rtimes \mathrm{SL}_2$.

Proof. $\mathrm{GRT}_{I_2}^{ell}(\mathbf{k})$ is a subgroup of $\mathrm{GRT}_1(\mathbf{k}) \times \mathrm{Ker}\left(\mathrm{Aut}(\hat{\mathfrak{t}}_{1,2}^{\mathbf{k}}) \to \mathrm{GL}_2(\mathbf{k})\right)$; the assignment $\mathbf{k} \mapsto$ (the latter group) is a prounipotent group scheme, hence so is $\mathrm{GRT}_{I_2}^{ell}(-)$. The second statement follows from Lemma 49. ☐

The group $\mathbf{k}^\times$ acts on $\mathrm{GRT}_1^{ell}(\mathbf{k})$ by $c \cdot (g, u_\pm) := (c \cdot g, c \cdot u_\pm)$, where $c \cdot g$ is as above, $(c \cdot u_+)(x_1^+, x_1^-) := u_+(x_1^+, c^{-1}x_1^-)$, $(c \cdot u_-)(x_1^+, x_1^-) := cu_-(x_1^+, c^{-1}x_1^-)$. We then set $\mathrm{GRT}_{ell}(\mathbf{k}) := \mathrm{GRT}_1^{ell}(\mathbf{k}) \rtimes \mathbf{k}^\times$. Then $\mathbf{k} \mapsto \mathrm{GRT}_{ell}(\mathbf{k})$ is a $\mathbb{Q}$-group scheme, we have $\mathrm{GRT}_{ell}(-) = \mathrm{GRT}_1^{ell}(-) \rtimes \mathbb{G}_m$.

We then have a group scheme morphism

$$\mathrm{GRT}_{ell}(-) \to \mathrm{GRT}(-), \tag{31}$$

induced by $(g, u_\pm) \mapsto g$ and the identity on $\mathbb{G}_m$.

To elucidate the structure of $\mathrm{GRT}_{ell}(-)$, we use the following statement on iterated semidirect products:

Lemma 51. *Let G_i be groups $(i = 1, 2, 3)$. The following data are equivalent:*

(a) actions[12] $G_j \to \mathrm{Aut}(G_i)$ *for* $i < j$, *such that* $g_3 * (g_2 * g_1) = (g_3 * g_2) * (g_3 * g_1)$;

(b) actions $G_j \to \mathrm{Aut}(G_i)$ *for* $(i, j) = (1, 2)$ *and* $(2, 3)$, *and an action* $G_{23} \to \mathrm{Aut}(G_{12})$ *(where* $G_{ij} := G_i \rtimes G_j$*), compatible with the actions of G_j on G_i for* $(i, j) = (1, 2)$ *or* $(2, 3)$, *and with the adjoint action of G_2 on itself.*

These equivalent data yield actions $G_3 \to \mathrm{Aut}(G_{12})$ *and* $G_{23} \to \mathrm{Aut}(G_1)$, *and we then have a canonical isomorphism* $(G_1 \rtimes G_2) \rtimes G_3 \simeq G_1 \rtimes (G_2 \rtimes G_3)$.

---

[12] The action of $g_j \in G_j$ on $g_i \in G_i$ is denoted $g_j * g_i \in G_i$.

Proof. Straightforward. $\square$

We then have an action of GL_2 on $\mathrm{GRT}_1^{ell}(-)$, given by $\gamma \cdot (g, u_+, u_-) := (|\gamma| \cdot g, \tilde{u}_+, \tilde{u}_-)$, where $\left(\begin{smallmatrix} \tilde{u}_+(x_1,y_1) \\ \tilde{u}_-(x_1,y_1) \end{smallmatrix}\right) := \gamma \left(\begin{smallmatrix} u_+(\tilde{x}_1,\tilde{y}_1) \\ u_-(\tilde{x}_1,\tilde{y}_1) \end{smallmatrix}\right)$ and $\left(\begin{smallmatrix} \tilde{x}_1 \\ \tilde{y}_1 \end{smallmatrix}\right) := \gamma^{-1}\left(\begin{smallmatrix} x_1 \\ y_1 \end{smallmatrix}\right)$. It satisfies the conditions of Lemma 51, (b), where: $G_1 = \mathrm{GRT}_{I_2}^{ell}(-)$, $G_2 = \mathrm{SL}_2$, $G_3 = \mathbb{G}_m$, the isomorphism $G_2 \rtimes G_3 \simeq \mathrm{GL}_2$ being given by $\mathbb{G}_m \to \mathrm{GL}_2$, $c \mapsto \left(\begin{smallmatrix} 1 & 0 \\ 0 & c \end{smallmatrix}\right)$. We have therefore an isomorphism

$$\mathrm{GRT}_{ell}(-) \simeq \mathrm{GRT}_{I_2}^{ell}(-) \rtimes \mathrm{GL}_2,$$

where we recall that $\mathrm{GRT}_{I_2}^{ell}(-)$ is prounipotent.

The morphism (31) then fits in a commutative diagram

$$\begin{tikzcd}
\mathrm{GRT}_{ell}(-) \arrow[r] \arrow[d] & \mathrm{GRT}(-) \arrow[d] \\
\mathrm{GL}_2 \arrow[r, "\det"] & \mathbb{G}_m
\end{tikzcd}$$

as the morphism $G_2 \rtimes G_3 \to G_3$ coincides with det.

6.3. A morphism $\mathrm{GRT}(\mathbf{k}) \to \mathrm{GRT}_{ell}(\mathbf{k})$.

We now construct a section of (31). We first set

$$t_{0i} := -\frac{\mathrm{ad}\, x_i}{e^{\mathrm{ad}\, x_i} - 1}(y_i) \in \hat{\mathfrak{t}}_{1,n}^{\mathbf{k}} \text{ for } i \in \{1, ..., n\}. \tag{32}$$

For $g \in \exp(\hat{\mathfrak{f}}_2^{\mathbf{k}}) \subset \exp(\hat{\mathfrak{t}}_3^{\mathbf{k}})$ (where the inclusion is $A \mapsto t_{12}$, $B \mapsto t_{23}$), we set $g^{0,1,2} := g(t_{01}, t_{12}) \in \exp(\hat{\mathfrak{t}}_{1,2}^{\mathbf{k}})$ $g^{0,2,1} := g(t_{02}, t_{21}) \in \exp(\hat{\mathfrak{t}}_{1,2}^{\mathbf{k}})$.

Lemma-Definition 52. *For $g \in \exp(\hat{\mathfrak{f}}_2^{\mathbf{k}})$, there exists $\alpha_g \in \mathrm{Aut}(\hat{\mathfrak{t}}_{1,2}^{\mathbf{k}})$, uniquely defined by $\alpha_g(x_1) = \log(g^{0,2,1} e^{x_1} (g^{0,1,2})^{-1})$, $\alpha_g(t_{01}) = g^{0,1,2} t_{01} (g^{0,1,2})^{-1}$. We set $(u_+^g, u_-^g) := (\alpha_g(x_1), \alpha_g(y_1))$.*

Proof. This follows from the fact that $\mathfrak{t}_{1,2}^{\mathbf{k}}$ is freely generated by x_1 and t_{01}. $\square$

Proposition 53. *The map $\mathrm{GRT}_1(\mathbf{k}) \to \mathrm{GRT}_1^{ell}(\mathbf{k})$, $g \mapsto (g, u_+^g, u_-^g)$ is a group morphism. It is compatible with the action of $\mathbf{k}^\times$, hence extends to a group morphism $\mathrm{GRT}(\mathbf{k}) \to \mathrm{GRT}_{ell}(\mathbf{k})$, section of (31). The latter fits in a commutative diagram*

$$\begin{tikzcd}
\mathrm{GRT}(-) \arrow[r] \arrow[d] & \mathrm{GRT}_{ell}(-) \arrow[d] \\
\mathbb{G}_m \arrow[r] & \mathrm{GL}_2,
\end{tikzcd}$$

where the bottom morphism is $c \mapsto \left(\begin{smallmatrix} 1 & 0 \\ 0 & c \end{smallmatrix}\right)$.

Proof. We first prove:

Lemma 54. *$\hat{\mathfrak{t}}_{1,n}^{\mathbf{k}}$ is isomorphic to the topological Lie algebra with generators $x_i, t_{\alpha\beta}$ ($i \in \{1, ..., n\}$, $\alpha \neq \beta \in \{0, ..., n\}$) and relations (we set $X_i := e^{x_i}$; $i, j, ...$ run over $\{1, ..., n\}$ while $\alpha, \beta, ...$ run over $\{0, \dots, n\}$):*

$$t_{\beta\alpha} = t_{\alpha\beta} \text{ for } \alpha \neq \beta, \quad [t_{\alpha\beta}, t_{\gamma\delta}] = [t_{\alpha\beta}, t_{\alpha\gamma} + t_{\beta\gamma}] = 0 \text{ for } \alpha, ..., \delta \text{ all different,} \tag{33}$$

$$\log(X_i, X_j) = \log(\prod_i X_i) = 0, \tag{34}$$

$$X_i (t_{0j} + t_{ij}) X_i^{-1} = t_{0j} \text{ if } i \neq j, \quad X_i t_{0i} X_i^{-1} = \sum_{\alpha \neq i} t_{\alpha i}, \tag{35}$$

$$X_i t_{jk} X_i^{-1} = t_{jk} \text{ for } i, j, k \text{ distinct,} \quad (X_j X_k) t_{jk} (X_j X_k)^{-1} = t_{jk} \text{ for } i \neq j, \tag{36}$$

$$\sum_{0\leq\alpha<\beta\leq n} t_{\alpha\beta}=0. \tag{37}$$

Proof. One first checks that if one defines t_{0i} as in (32), then the above relations are satisfied; conversely, if one sets $y_i := -\frac{e^{\operatorname{ad} x_i}-1}{\operatorname{ad} x_i}(t_{0i})$, then the above relations lead to the defining relations of $\hat{\mathfrak{t}}^{\mathbf{k}}_{1,n}$. ☐

Lemma 55. *Let* $(g,u_\pm)\in \mathrm{GRT}_1^{ell}(\mathbf{k})$ *and* $\alpha\in\mathrm{Aut}(\hat{\mathfrak{t}}^{\mathbf{k}}_{1,2})$ *be defined by* $\alpha(x_1^\pm)=u_\pm$. *Then* $\theta_{g,u_\pm}\in\mathrm{Aut}(\hat{\mathfrak{t}}^{\mathbf{k}}_{1,3})$ *(see Lemma 47) may be defined by*

$$\theta_{g,u_\pm}: X_1\mapsto \mathrm{Ad}(g^{1,2,3})^{-1}(\alpha(X_1)^{1,23}),\quad X_2\mapsto \mathrm{Ad}(g^{2,1,3})^{-1}(\alpha(X_1)^{2,13}),\quad X_3\mapsto\alpha(X_1)^{3,12},$$
$$t_{01}\mapsto \mathrm{Ad}(g^{1,2,3})^{-1}(\alpha(t_{01})^{1,23}),\quad t_{02}\mapsto \mathrm{Ad}(g^{2,1,3})^{-1}(\alpha(t_{01})^{2,13}),\quad t_{03}\mapsto\alpha(t_{01})^{3,12},$$
$$t_{12}\mapsto t_{12},\quad t_{23}\mapsto \mathrm{Ad}(g^{1,2,3})^{-1}(t_{23}),\quad t_{13}\mapsto \mathrm{Ad}(g^{2,1,3})^{-1}(t_{13}).$$

Proof. Immediate. ☐

Lemma 56. *Let* $g\in\mathrm{GRT}_1(\mathbf{k})$. *There is a unique* $\tilde\theta_g\in\mathrm{Aut}(\hat{\mathfrak{t}}^{\mathbf{k}}_{1,3})$, *such that*

$$\tilde\theta_g: X_1\mapsto (g^{1,2,3})^{-1}g^{0,23,1}X_1(g^{0,1,23})^{-1}g^{1,2,3},\quad X_2\mapsto (g^{2,1,3})^{-1}g^{0,13,2}X_2(g^{0,2,13})^{-1}g^{2,1,3},$$
$$X_3\mapsto g^{0,12,3}X_3(g^{0,3,12})^{-1},$$

$$t_{01}\mapsto\mathrm{Ad}((g^{1,2,3})^{-1}g^{0,1,23})(t_{01}),\quad t_{02}\mapsto\mathrm{Ad}((g^{2,1,3})^{-1}g^{0,2,13})(t_{02}),\quad t_{03}\mapsto\mathrm{Ad}(g^{0,3,12})(t_{03}), \tag{38}$$

$$t_{12}\mapsto t_{12},\quad t_{23}\mapsto\mathrm{Ad}(g^{1,2,3})^{-1}(t_{23}),\quad t_{13}\mapsto\mathrm{Ad}(g^{2,1,3})^{-1}(t_{13}). \tag{39}$$

Proof. Let us first prove that relations (33) and (37) (for $n=3$) are preserved. In Subsection 6.4, we will construct a functor $\mathbf{PaCD}\to g*\mathbf{PaCD}$, which gives rise to an automorphism $\exp(\hat{\mathfrak{t}}^{\mathbf{k}}_n)\rtimes S_n=\mathbf{PaCD}(O)\to(\mathbf{PaCD}*g)(O)=\exp(\hat{\mathfrak{t}}^{\mathbf{k}}_n)\rtimes S_n$ for any $O\in\mathbf{PaCD}(O)$, $|O|=n$. When $O=\bullet((\bullet\bullet)\bullet)$, the resulting automorphism of $\hat{\mathfrak{t}}^{\mathbf{k}}_4$ is given by (38), (39). So relations (33) are preserved. The automorphism necessarily preserves $Z(\hat{\mathfrak{t}}^{\mathbf{k}}_4)=\mathbf{k}\cdot\sum_{\alpha<\beta}t_{\alpha\beta}$, so relation (37) is also preserved.

Note for later use that

$$\tilde\theta_g(x^{2,3,1})=\mathrm{Ad}(g^{1,2,3})^{-1}(\tilde\theta_g(x)^{2,3,1})\text{ for }x\in\{x_i,t_{\alpha\beta}\}. \tag{40}$$

We have

$$\begin{aligned}
\tilde\theta_g(X_2)\tilde\theta_g(X_3) &= (g^{2,1,3})^{-1}g^{0,13,2}X_2(g^{0,2,13})^{-1}g^{2,1,3}g^{0,21,3}X_3(g^{0,3,12})^{-1}\\
&= (g^{2,1,3})^{-1}g^{0,13,2}X_2g^{02,1,3}(g^{0,2,1})^{-1}X_3(g^{0,3,12})^{-1}\\
&= (g^{2,1,3})^{-1}g^{0,13,2}g^{0,1,3}X_2X_3(g^{03,2,1})^{-1}(g^{0,3,12})^{-1}\\
&= (g^{2,1,3})^{-1}(g^{1,3,2})^{-1}g^{0,1,32}g^{01,3,2}X_2X_3(g^{0,3,2})^{-1}(g^{0,32,1})^{-1}(g^{3,2,1})^{-1}\\
&= (g^{2,1,3})^{-1}(g^{1,3,2})^{-1}g^{0,1,32}X_2X_3(g^{0,32,1})^{-1}(g^{3,2,1})^{-1},
\end{aligned}$$

while

$$\begin{aligned}
\tilde\theta_g(X_3)\tilde\theta_g(X_2) &= g^{0,21,3}X_3(g^{0,3,12})^{-1}(g^{2,1,3})^{-1}g^{0,13,2}X_2(g^{0,2,13})^{-1}g^{2,1,3}\\
&= g^{0,21,3}X_3g^{03,1,2}(g^{0,3,1})^{-1}X_2(g^{0,2,13})^{-1}g^{2,1,3}\\
&= g^{0,12,3}g^{0,1,2}X_3X_2(g^{02,3,1})^{-1}(g^{0,2,13})^{-1}g^{2,1,3}\\
&= (g^{1,2,3})^{-1}g^{0,1,23}g^{01,2,3}X_3X_2(g^{0,2,3})^{-1}(g^{0,23,1})^{-1}(g^{2,3,1})^{-1}g^{2,1,3}\\
&= (g^{1,2,3})^{-1}g^{0,1,23}X_3X_2(g^{0,23,1})^{-1}(g^{2,3,1})^{-1}g^{2,1,3},
\end{aligned}$$

which implies $(\tilde\theta_g(X_2),\tilde\theta_g(X_3))=1$. Then (40) implies that $(\tilde\theta_g(X_i),\tilde\theta_g(X_j))=1$ for any i,j.

The above computation of $\tilde{\theta}_g(X_2)\tilde{\theta}_g(X_3)$ implies that

$$\begin{aligned}
&\tilde{\theta}_g(X_1)\tilde{\theta}_g(X_2)\tilde{\theta}_g(X_3)\\
&= (g^{1,2,3})^{-1}g^{0,23,1}X_1(g^{0,1,23})^{-1}g^{1,2,3}(g^{2,1,3})^{-1}(g^{1,3,2})^{-1}X_2X_3(g^{0,32,1})^{-1}(g^{3,2,1})^{-1} = 1
\end{aligned}$$

as $X_1X_2X_3 = 1$. So $X_1X_2X_3 = 1$ is preserved.

$\tilde{\theta}_g(X_3)$ clearly commutes with $\tilde{\theta}_g(t_{12})$, which implies that $X_jt_{jk}X_i^{-1} = t_{jk}$ is preserved in view of (40), as well as $X_jX_kt_{jk}(X_jX_k)^{-1} = t_{jk}$ (as the X_i commute and $X_1X_2X_3 = 1$).

Now

$$\begin{aligned}
&\tilde{\theta}_g(t_{02} + t_{12}) = \mathrm{Ad}((g^{2,1,3})^{-1}g^{0,2,31})(t_{02}) + t_{12} = \mathrm{Ad}(g^{0,21,3})(\mathrm{Ad}(g^{0,2,1})(t_{02}) + t_{12})\\
&= \mathrm{Ad}(g^{0,21,3})(t_{01} + t_{02} + t_{12} - \mathrm{Ad}(g^{0,1,2})(t_{01})) = t_{12} + \mathrm{Ad}(g^{0,21,3})(t_{01} + t_{02}) - \mathrm{Ad}(g^{0,12,3}g^{0,1,2})(t_{01}).
\end{aligned}$$

Then

$$\begin{aligned}
&\tilde{\theta}_g(X_1)\tilde{\theta}_g(t_{02} + t_{12}) = (g^{1,2,3})^{-1}g^{0,23,1}X_1\Big((g^{0,1,23})^{-1}g^{1,2,3}t_{12} + (g^{0,1,23})^{-1}g^{1,2,3}g^{0,12,3}(t_{01} + t_{02})(g^{0,12,3})^{-1}\\
&-(g^{0,1,23})^{-1}g^{1,2,3}g^{0,12,3}g^{0,1,2}t_{01}(g^{0,1,2})^{-1}(g^{0,12,3})^{-1}\Big)\\
&= (g^{1,2,3})^{-1}g^{0,23,1}X_1\Big(g^{01,2,3}(g^{0,1,2})^{-1}t_{12}(g^{0,12,3})^{-1} + g^{01,2,3}(g^{0,1,2})^{-1}(t_{01} + t_{02})(g^{0,12,3})^{-1}\\
&-g^{01,2,3}t_{01}(g^{0,1,2})^{-1}(g^{0,12,3})^{-1}\Big)\\
&= (g^{1,2,3})^{-1}g^{0,23,1}X_1g^{01,2,3}(t_{02} + t_{12})(g^{0,1,2})^{-1}(g^{0,12,3})^{-1}\\
&= (g^{1,2,3})^{-1}g^{0,23,1}g^{0,2,3}t_{02}X_1(g^{0,1,2})^{-1}(g^{0,12,3})^{-1},
\end{aligned}$$

while

$$\begin{aligned}
\tilde{\theta}_g(t_{02})\tilde{\theta}_g(X_1) &= (g^{2,1,3})^{-1}g^{0,2,31}t_{02}(g^{0,2,31})^{-1}g^{2,1,3}(g^{1,2,3})^{-1}g^{0,23,1}X_1(g^{0,1,23})^{-1}g^{1,2,3}\\
&= (g^{2,1,3})^{-1}g^{0,2,31}t_{02}g^{02,3,1}(g^{0,2,3})^{-1}X_1(g^{0,1,23})^{-1}g^{1,2,3}\\
&= (g^{2,1,3})^{-1}g^{0,2,31}g^{02,3,1}t_{02}X_1(g^{01,2,3})^{-1}(g^{0,1,23})^{-1}g^{1,2,3}\\
&= g^{3,1,2}g^{2,3,1}g^{0,23,1}g^{0,2,3}t_{02}X_1(g^{0,1,2})^{-1}(g^{0,12,3})^{-1},
\end{aligned}$$

so the relation $X_1(t_{02} + t_{12})X_1^{-1} = t_{02}$ is preserved. (40) then implies that the relations $X_i(t_{0j} + t_{ij})X_i^{-1} = t_{0j}$ are preserved. Together with the other relations, these relations imply $X_it_{0i}X_i^{-1} = \sum_{\alpha\neq i} t_{\alpha i}$, which are therefore also preserved. □

End of proof of Proposition 53. If $g \in \mathrm{GRT}_1(\mathbf{k})$, then one checks that the automorphisms $\tilde{\theta}_g$ from Lemma 56 and $\alpha_g \in \mathrm{Aut}(\hat{\mathfrak{t}}^{\mathbf{k}}_{1,2})$ from Lemma-Definition 52 are related in the same way as $\theta_{g,u_\pm}$ and α are in Lemma 55. It follows that if $u_\pm^g$ are as in Lemma-Definition 52, then $(g, u_\pm^g) \in \mathrm{GRT}_1^{ell}(\mathbf{k})$. This defines a map $\mathrm{GRT}_1(\mathbf{k}) \to \mathrm{GRT}_1^{ell}(\mathbf{k})$.

We now show that $\mathrm{GRT}_1(\mathbf{k}) \to \mathrm{GRT}_1^{ell}(\mathbf{k})$ is a group morphism. In view of (30), it suffices to prove that $\tilde{\theta}_{g_2}\tilde{\theta}_{g_1} = \tilde{\theta}_{g_1*g_2}$, which can be checked directly, e.g.,

$$\begin{aligned}
&\tilde{\theta}_{g_2}(\tilde{\theta}_{g_1}(X_1)) = \tilde{\theta}_{g_2}(g_1^{0,2,1}X_1(g^{0,1,2})^{-1}) = \tilde{\theta}_{g_2}(g_1(t_{02}, t_{21})X_1g_1^{-1}(t_{01}, t_{12}))\\
&= g_1(\mathrm{Ad}(g_2^{0,2,1})(t_{02}), t_{21})g_2^{0,2,1}X_1(g_2^{0,2,1})^{-1}g_1^{-1}(\mathrm{Ad}(g_2^{0,2,1})(t_{01}), t_{12})\\
&= (g_1 * g_2)^{0,2,1}X_1((g_1 * g_2)^{0,1,2})^{-1} = \tilde{\theta}_{g_1*g_2}(X_1),
\end{aligned}$$

etc.

We now prove that $\mathrm{GRT}_1(\mathbf{k}) \to \mathrm{GRT}_1^{ell}(\mathbf{k})$ is compatible with the actions of $\mathbf{k}^\times$. If $c \cdot (g, u_\pm) = (\tilde{g}, \tilde{u}_\pm)$, then $\theta_{\tilde{g},\tilde{u}_\pm}$ and $\theta_{g,u_\pm}$ are related by $\theta_{\tilde{g},\tilde{u}_\pm} = \gamma_c\theta_{g,u_\pm}\gamma_c^{-1}$, where $\gamma_c \in \mathrm{Aut}(\hat{\mathfrak{t}}^{\mathbf{k}}_{1,3})$ is $\gamma_c(x_i^+) = x_i^+$, $\gamma_c(x_i^-) = c^{-1}x_i^-$. It then suffices to prove that $\tilde{\theta}_{\tilde{g}} = \gamma_c\tilde{\theta}_g\gamma_c^{-1}$, where we recall that $\tilde{g}(A, B) = g(c^{-1}A, c^{-1}B)$, which follows from $\gamma_c(x_i) = x_i$, $\gamma_c(t_{\alpha\beta}) = c^{-1}t_{\alpha\beta}$ for $0 \leq \alpha \neq \beta \leq 3$.

The final commutative diagram follows from

$$\begin{tikzcd}
\mathrm{GRT}_1^{ell}(\mathbf{k}) \rtimes \mathbf{k}^\times \arrow[d] \arrow[dr] & \\
\mathrm{SL}_2(\mathbf{k}) \rtimes \mathbf{k}^\times \arrow[r, "\sim"] & \mathrm{GL}_2(\mathbf{k})
\end{tikzcd}$$

where $\mathrm{SL}_2(\mathbf{k}) \rtimes \mathbf{k}^\times \simeq \mathrm{GL}_2(\mathbf{k})$ is induced by $\mathbf{k}^\times \to \mathrm{GL}_2(\mathbf{k})$, $c \mapsto \left(\begin{smallmatrix} 1 & 0 \\ 0 & c \end{smallmatrix}\right)$. □

We set

$$R^{gr}_{ell}(\mathbf{k}) := \mathrm{Ker}\left(\mathrm{GRT}_{ell}(\mathbf{k}) \to \mathrm{GRT}(\mathbf{k})\right). \tag{41}$$

Explicitly,

$$\begin{aligned} R^{gr}_{ell}(\mathbf{k}) &= \{(u_+, u_-) \in (\hat{\mathfrak{t}}^{\mathbf{k}}_{1,2})^2 | u_\pm^{1,23} + u_\pm^{2,31} + u_\pm^{3,12} = 0, [u_\pm^{1,23}, u_\pm^{2,13}] = 0, [u_+^{1,23}, u_-^{2,13}] = t_{12}\} \\ &\subset \mathrm{Aut}(\hat{\mathfrak{t}}^{\mathbf{k}}_{1,2})^{op}. \end{aligned} \tag{42}$$

Then $\mathbf{k} \mapsto R^{gr}_{ell}(\mathbf{k})$ is $\mathbb{Q}$-group scheme, and we have a commutative diagram

$$\begin{tikzcd}
1 \arrow[r] & R^{gr}_{ell}(-) \arrow[r] \arrow[d] & \mathrm{GRT}_{ell}(-) \arrow[r] \arrow[d] & \mathrm{GRT}(-) \arrow[r] \arrow[d] & 1 \\
1 \arrow[r] & \mathrm{SL}_2 \arrow[r] & \mathrm{GL}_2 \arrow[r, "\det"] & \mathbb{G}_m \arrow[r] & 1
\end{tikzcd}$$

The lift of $\mathrm{GRT}_{ell}(-) \to \mathrm{GL}_2$ restricts to a morphism $\mathrm{SL}_2 \to R^{gr}_{ell}(-)$, and the structure of $R^{gr}_{ell}(-)$ is therefore

$$R^{gr}_{ell}(-) = \mathrm{Ker}\left(R^{gr}_{ell}(-) \to \mathrm{SL}_2\right) \rtimes \mathrm{SL}_2,$$

in which the kernel is prounipotent.

The morphism from Proposition 53 enables us to define an action of $\mathrm{GRT}(-)$ on $R^{gr}_{ell}(-)$. $\mathrm{GRT}_{ell}(-)$ has then the structure of a semidirect product, fitting in

$$\begin{tikzcd}
\mathrm{GRT}_{ell}(-) \arrow[r, phantom, "\simeq"] \arrow[d] & R^{gr}_{ell}(-) \rtimes \mathrm{GRT}(-) \arrow[d] \\
\mathrm{GL}_2 \arrow[r, phantom, "\simeq"] & \mathrm{SL}_2 \rtimes \mathbb{G}_m
\end{tikzcd}$$

where the bottom morphism is induced by $\mathbb{G}_m \to \mathrm{GL}_2$, $c \mapsto \left(\begin{smallmatrix} 1 & 0 \\ 0 & c \end{smallmatrix}\right)$.

6.4. **Categorical interpretations.** A left action of $\mathrm{GRT}(\mathbf{k})$ on {IMBCs} is defined as follows: $g \in \mathrm{GRT}_1(\mathbf{k})$ acts on $(\mathcal{C}_0, c_{-,-}, a_{-,-,-}, t_{-,-})$ by only modifying a_{XYZ} into $\tilde{a}_{X,Y,Z} := a_{XYZ} g(t_{XY} \otimes \mathrm{id}_Z, a^{-1}_{XYZ}(\mathrm{id}_X \otimes t_{YZ}) a_{XYZ})$ and $c \in \mathbf{k}^\times$ acts by only modifying t_{XY} into ct_{XY}.

Similarly, one can show that a left action of $\mathrm{GRT}_{ell}(\mathbf{k})$ on {(an IBMC, an elliptic structure over it)} is defined as follows: $(g, u_\pm) \in \mathrm{GRT}_1^{ell}(\mathbf{k})$ acts on $(\mathcal{C}_0, \mathcal{C})$ as $(g, u_\pm) * (\mathcal{C}_0, \mathcal{C}) := (g * \mathcal{C}_0, \tilde{\mathcal{C}})$, where for $\mathcal{C} = (\mathcal{C}, F, x^\pm_{-,-})$, we set $\tilde{\mathcal{C}} = (\mathcal{C}, F, \tilde{x}^\pm_{-,-})$, where $\tilde{x}^\pm_{X,Y} = u^\pm(x^+_{X,Y}, x^-_{X,Y})$, and $c \in \mathbf{k}^\times$ acts on $(\mathcal{C}_0, \mathcal{C})$ as $c * (\mathcal{C}_0, \mathcal{C}) := (c * \mathcal{C}_0, \tilde{\mathcal{C}})$, where $\tilde{\mathcal{C}} = (\mathcal{C}, F, x^+_{X,Y}, cx^-_{X,Y})$.

6.5. **Action of** $\mathrm{GRT}_{ell}(\mathbf{k})$ **on** $\underline{\mathit{Ell}}(\mathbf{k})$**.** Recall that $\mathrm{GRT}(\mathbf{k})$ acts on $\underline{M}(\mathbf{k})$ from the right as follows: for $g \in \mathrm{GRT}_1(\mathbf{k})$ and $(\mu, \Phi) \in \underline{M}(\mathbf{k})$, $(\mu, \Phi) * g := (\mu, \tilde{\Phi})$, where

$$\tilde{\Phi}(t_{12}, t_{23}) = \Phi(\mathrm{Ad}(g^{1,2,3})(t_{12}), t_{23}) g^{1,2,3},$$

and for $c \in \mathbf{k}^\times$,$(\mu, \Phi) * c := (c\mu, c * \Phi)$, where $(c * \Phi)(A, B) = \Phi(cA, cB)$. This action is compatible with the maps {IBMCs} → {BMCs} induced by elements of $\underline{M}(\mathbf{k})$ as follows: $\Phi * (g * \mathcal{C}_0) = (\Phi * g) * \mathcal{C}_0$ for any $\Phi \in \underline{M}(\mathbf{k})$, $g \in \mathrm{GRT}(\mathbf{k})$ and IBMC $\mathcal{C}_0$.

For $(g, u_\pm) \in \mathrm{GRT}_1^{ell}(\mathbf{k})$ and $(\mu, \Phi, A_\pm) \in \underline{\mathit{Ell}}(\mathbf{k})$, we set $(\mu, \Phi, A_\pm) * (g, u_\pm) := (\mu, \tilde{\Phi}, \tilde{A}_\pm)$, where

$$\tilde{A}_\pm(x_1, y) := A_\pm(u_+(x_1, y_1), u_-(x_1, y_1))$$

(in other terms, $\tilde{A}_\pm = \theta(A_\pm)$, where $\theta \in \mathrm{Aut}(\hat{\mathfrak{t}}^{\mathbf{k}}_{1,2})$ is $x_1^\pm \mapsto u_\pm(x_1^+, x_1^-)$) and for $c \in \mathbf{k}^\times$, we set $(\mu, \Phi, A_\pm) := (\mu, c * \Phi, c\sharp A_\pm)$, where $(c\sharp A_\pm)(x_1^+, x_1^-) := A_\pm(x_1^+, cx_1^-)$.

Proposition 57. *This defines a right action of* $\mathrm{GRT}_{ell}(\mathbf{k})$ *on* $\underline{Ell}(\mathbf{k})$*, commuting with the left action of* $\underline{\mathrm{GT}}_{ell}(\mathbf{k})$ *and compatible with the right action of* $\mathrm{GL}_2(\mathbf{k})$ *on* $\mathrm{M}_2(\mathbf{k})$*.*

Proof. Let us show that $(\mu, \tilde\Phi, \tilde A_\pm) \in \underline{Ell}(\mathbf{k})$. If $\theta \in \mathrm{Aut}(\hat{\mathfrak{t}}^{\mathbf{k}}_{1,2})$ is defined by $\theta(x_1^\pm) = u_\pm$, and $\tilde\theta := \theta_{g,u_\pm}$, then

$$\tilde\theta(x^{1,23}) = \mathrm{Ad}(g^{1,2,3})^{-1}(\theta(x)^{1,23}), \quad \tilde\theta(x^{2,31}) = \mathrm{Ad}(g^{2,1,3})^{-1}(\theta(x)^{2,31}), \quad \tilde\theta(x^{3,12}) = \theta(x)^{3,12}$$

for any $x \in \hat{\mathfrak{t}}^{\mathbf{k}}_{1,2}$. Applying $\tilde\theta$ to (22), we get

$$\theta(\{e^{\pm\mu t_{12}/2}\}A_\pm)^{3,12}\tilde\theta(\Phi^{3,1,2})(g^{2,1,3})^{-1}\theta(\{e^{\pm\mu t_{12}/2}\}A_\pm)^{2,31}g^{2,1,3}\tilde\theta(\Phi^{2,3,1})(g^{1,2,3})^{-1}\theta(\{e^{\pm\mu t_{12}/2}\}A_\pm)^{1,23} \\ g^{1,2,3}\tilde\theta(\Phi^{1,2,3}) = 1.$$

Now $\tilde\theta(\Phi^{3,1,2})(g^{2,1,3})^{-1} = \tilde\Phi^{3,1,2}$, $g^{2,1,3}\tilde\theta(\Phi^{2,3,1})(g^{1,2,3})^{-1} = \tilde\Phi^{2,3,1}$, $g^{1,2,3}\tilde\theta(\Phi^{1,2,3}) = \tilde\Phi^{1,2,3}$. Since $\theta(\{e^{\pm\mu t_{12}/2}\}A_\pm) = \{e^{\pm\mu t_{12}/2}\}\tilde A_\pm$, $(\mu, \tilde\Phi, \tilde A_\pm)$ satisfies (22).

Applying now $\tilde\theta$ to (23), one gets

$$e^{\mu t_{12}} = \big(\tilde\theta(\Phi)^{-1}g^{-1}\theta(A_-)^{1,23}g\tilde\theta(\Phi), e^{-\mu t_{12}/2}\tilde\theta(\Phi^{2,1,3})^{-1}(g^{2,1,3})^{-1}(\theta(A)^{2,13})^{-1}g^{2,1,3}\tilde\theta(\Phi^{2,1,3})e^{-\mu t_{12}/2}\big).$$

Using again $g\tilde\theta(\Phi) = \tilde\Phi$ and $g^{2,1,3}\tilde\theta(\Phi^{2,1,3}) = \tilde\Phi^{2,1,3}$, together with $\theta(A_\pm) = \tilde A_\pm$, we see that $(\mu, \tilde\Phi, \tilde A_\pm)$ satisfies (23).

Similarly, applying the automorphism $x_i^+ \mapsto x_i^+$, $x_i^- \mapsto cx_i^-$ to identities (22), (23), we see that $(\mu, \Phi, A_\pm) * c$ again satisfies these identities, hence belongs to $\underline{Ell}(\mathbf{k})$. It is then immediate to check that we have a right action of $\mathrm{GRT}_{ell}(\mathbf{k})$, commuting with the left action of $\underline{\mathrm{GT}}_{ell}(\mathbf{k})$. □

Proposition 58. *The action of* $\mathrm{GRT}_{ell}(\mathbf{k})$ *restricts to an action on* $Ell(\mathbf{k}) \subset \underline{Ell}(\mathbf{k})$*, which is free and transitive.*

Proof. Given that the action of $\mathrm{GRT}(\mathbf{k})$ on $M(\mathbf{k})$ is free and transitive, it suffices to prove that the action of $R^{gr}_{ell}(\mathbf{k})$ on $Ell_{(\mu,\Phi)}(\mathbf{k}) := Ell(\mathbf{k}) \times_{M(\mathbf{k})} \{(\mu, \Phi)\}$ is free and transitive for any $(\mu, \Phi) \in Ell(\mathbf{k})$.

Recall that $R^{gr}_{ell}(\mathbf{k})$ is explicitly described by (42); its inclusion into $\mathrm{Aut}(\hat{\mathfrak{t}}^{\mathbf{k}}_{1,2})$ is by $(u_+, u_-) \mapsto \theta_{u_+,u_-} = (x_1^\pm \mapsto u_\pm)$. On the other hand, $Ell_{(\mu,\Phi)}(\mathbf{k}) = \{(A_+, A_-)$ satisfying (22), (23)$\}$. Then

$$(A_+, A_-) * (u_+, u_-) = (\theta_{u_\pm}(A_+), \theta_{u_\pm}(A_-)). \tag{43}$$

Relation (23) implies that $(A_-, A_+) = e^{\mu t_{12}}$, which together with $\mu \in \mathbf{k}^\times$ implies that $\hat{\mathfrak{t}}^{\mathbf{k}}_{1,2}$ is generated by $\log A_\pm$. Together with (43), this implies that the action of $R^{gr}_{ell}(\mathbf{k})$ on $Ell_{(\mu,\Phi)}(\mathbf{k})$ is free.

Let us now show that this action is transitive. We first observe that $R^{gr}_{ell}(\mathbf{k})$ can be described as $\{\theta \in \mathrm{Aut}(\hat{\mathfrak{t}}^{\mathbf{k}}_{1,2}) | \exists\tilde\theta \in \mathrm{Aut}(\hat{\mathfrak{t}}^{\mathbf{k}}_{1,3})$ with $\tilde\theta(t_{ij}) = t_{ij}$ for $1 \le i \ne j \le 3$ and $\tilde\theta(x^{i,jk}) = \theta(x)^{i,jk}$ for $\{i, j, k\} = \{1, 2, 3\}$ and $x \in \hat{\mathfrak{t}}^{\mathbf{k}}_{1,2}\}$. Let (A_+, A_-) and $(\tilde A_+, \tilde A_-) \in Ell_{(\mu,\Phi)}(\mathbf{k})$ and let $\theta \in \mathrm{Aut}(\hat{\mathfrak{t}}^{\mathbf{k}}_{1,2})$ be the automorphism such that $\theta(A_\pm) = \tilde A_\pm$. Let us show that there exists $\tilde\theta \in \mathrm{Aut}(\hat{\mathfrak{t}}^{\mathbf{k}}_{1,3})$, such that

$$\tilde\theta(t_{ij}) = t_{ij} \text{ for } 1 \le i \ne j \le 3 \quad \text{and} \quad \tilde\theta(x^{i,jk}) = \theta(x)^{i,jk} \text{ for } \{i, j, k\} = \{1, 2, 3\} \text{ and } x \in \hat{\mathfrak{t}}^{\mathbf{k}}_{1,2}. \tag{44}$$

Let $i_{(\mu,\Phi)} : B_3(\mathbf{k}) \to \exp(\hat{\mathfrak{t}}_3) \rtimes S_3$, $i_{(\mu,\Phi,A_\pm)} : B_{1,3}(\mathbf{k}) \to \exp(\hat{\mathfrak{t}}_{1,3}) \rtimes S_3$ be the isomorphisms induced by (μ, Φ), $(\mu, \Phi, A_\pm)$ and the object $\bullet(\bullet\bullet)$. We have a commutative diagram

$$\begin{tikzcd}
P_3(\mathbf{k}) \arrow[r, "\tilde{i}_{(\mu,\Phi)}"'] \arrow[d] \arrow[rr, bend left] & \exp(\hat{\mathfrak{t}}_3^{\mathbf{k}}) \arrow[d] \arrow[rr, bend left] & B_3(\mathbf{k}) \arrow[r, "i_{(\mu,\Phi)}"'] \arrow[d] & \exp(\hat{\mathfrak{t}}_3^{\mathbf{k}}) \rtimes S_3 \arrow[d] \\
P_{1,3}(\mathbf{k}) \arrow[r, "\tilde{i}_{(\mu,\Phi,A_\pm)}"] \arrow[rr, bend right] & \exp(\hat{\mathfrak{t}}_{1,3}^{\mathbf{k}}) \arrow[rr, bend right] & B_{1,3}(\mathbf{k}) \arrow[r, "i_{(\mu,\Phi,A_\pm)}"] & \exp(\hat{\mathfrak{t}}_{1,3}^{\mathbf{k}}) \rtimes S_3
\end{tikzcd}$$

where the maps 'i' are isomorphisms. Note that for $\sigma \in B_3$, $i_{(\mu,\Phi)}(\sigma)[\sigma]^{-1} \in \exp(\hat{\mathfrak{t}}_3^{\mathbf{k}})$ (where $\sigma \mapsto [\sigma]$ is the canonical morphism $B_3 \to S_3$).

Then

$$\tilde{i}_{(\mu,\Phi,A_\pm)}(X_1^\pm) = A_\pm^{1,23}, \quad \tilde{i}_{(\mu,\Phi,A_\pm)}(X_2^\pm) = \{i_\Phi(\sigma_1^{\pm1})s_1\}A_\pm^{2,13}\{s_1 i_\Phi(\sigma_1^{\pm1})\},$$
$$\tilde{i}_{(\mu,\Phi,A_\pm)}(X_3^\pm) = \{i_\Phi(\sigma_2^{\pm1}\sigma_1^{\pm1})s_1 s_2\}A_\pm^{3,12}\{s_2 s_1 i_\Phi(\sigma_1^{\pm1}\sigma_2^{\pm1})\},$$

where we recall that $x \mapsto \{x\}$ is induced by the canonical morphism $\mathfrak{t}_3 \to \mathfrak{t}_{1,3}$. Also $\tilde{i}_{(\mu,\Phi,A_\pm)}(\sigma_i^2) = \{\tilde{i}_{(\mu,\Phi)}(\sigma_i^2)\}$, for $i = 1, 2$ and $\tilde{i}_{(\mu,\Phi,A_\pm)}(\sigma_1\sigma_2^2\sigma_1) = \{\tilde{i}_{(\mu,\Phi)}(\sigma_1\sigma_2^2\sigma_1)\}$.

Let $\tilde{\theta} := \tilde{i}_{(\mu,\Phi,\tilde{A}_\pm)} \circ \tilde{i}^{-1}_{(\mu,\Phi,A_\pm)}$. Then $\tilde{\theta} \in \mathrm{Aut}(\hat{\mathfrak{t}}_{1,3}^{\mathbf{k}})$, and:

(a) $\tilde{\theta}$ leaves $\{\tilde{i}_{(\mu,\Phi)}(\sigma_i^2)\}$ $(i = 1, 2)$ and $\{\tilde{i}_{(\mu,\Phi)}(\sigma_1\sigma_2^2\sigma_1)\}$ fixed, so it leaves the image of $\hat{\mathfrak{t}}_3 \to \hat{\mathfrak{t}}_{1,3}$ pointwise fixed;

(b) $\tilde{\theta}(A_\pm^{1,23}) = \tilde{A}_\pm^{1,23}$, $\tilde{\theta}(\{i_{(\mu,\Phi)}(\sigma_1^{\pm1})s_1\}A_\pm^{2,13}\{s_1 i_{(\mu,\Phi)}(\sigma_1^{\pm1})\}) = \{i_{(\mu,\Phi)}(\sigma_1^{\pm1})s_1\}\tilde{A}_\pm^{2,13}\{s_1 i_{(\mu,\Phi)}(\sigma_1^{\pm1})\}$, which implies, as $\{i_{(\mu,\Phi)}(\sigma_1^{\pm1})s_1\}$, $\{s_1 i_{(\mu,\Phi)}(\sigma_1^{\pm1})\} \in \mathrm{im}(\exp(\hat{\mathfrak{t}}_3^{\mathbf{k}}) \to \exp(\hat{\mathfrak{t}}_{1,3}^{\mathbf{k}}))$, that $\tilde{\theta}(A_\pm^{2,13}) = \tilde{A}_\pm^{2,13}$; one proves similarly that $\tilde{\theta}(A_\pm^{3,12}) = \tilde{A}_\pm^{3,12}$.

(b) implies that $\tilde{\theta}(x^{i,jk}) = \theta(x)^{i,jk}$ holds for $x = A_\pm$, therefore also for x in the topological group generated by $A_\pm$. As $\mu \in \mathbf{k}^\times$, this group equals $\exp(\hat{\mathfrak{t}}_{1,2}^{\mathbf{k}})$, so $\tilde{\theta}$ satisfies (44). So $\theta \in R_{ell}^{gr}(\mathbf{k})$. □

Proposition 59. *The scheme morphisms* $\underline{Ell} \to \underline{M}$ *and* $\underline{M} \overset{\sigma}{\to} \underline{Ell}$ *(see Proposition 44) are compatible with the morphisms* $\mathrm{GRT}_{ell}(-) \to \mathrm{GRT}(-)$ *and* $\mathrm{GRT}(-) \to \mathrm{GRT}_{ell}(-)$ *(see Proposition 53).*

Proof. We need to prove the second statement only. Let $\underline{M}(\mathbf{k}) \overset{\sigma}{\to} \underline{Ell}(\mathbf{k})$ be given by $(\mu, \Phi) \mapsto (\mu, \Phi, A_\pm(\mu, \Phi))$, then we must show that for $g \in \mathrm{GRT}_1(\mathbf{k})$ and $(\mu, \tilde{\Phi}) = (\mu, \Phi) * g$, we have $A_\pm(\mu, \tilde{\Phi}) = \alpha_g(A(\mu, \Phi))$, where α_g is as in Lemma-Definition 52. This follows from the fact that α_g satisfies $\alpha_g(t_{02}) = \mathrm{Ad}(g^{0,2,1})(t_{02})$, $\alpha_g(t_{12}) = t_{12}$. It is also clear that $\underline{M}(\mathbf{k}) \overset{\sigma}{\to} \underline{Ell}(\mathbf{k})$ is compatible with the action of $\mathbf{k}^\times$. □

Remark 60. In fact, the commutative diagrams $\begin{array}{ccc} \underline{Ell} & \to & \underline{M} \\ \downarrow & & \downarrow \\ \mathrm{M}_2 & \overset{\det}{\to} & \mathbb{A} \end{array}$ and $\begin{array}{ccc} \underline{M} & \overset{\sigma}{\to} & \underline{Ell} \\ \downarrow & & \downarrow \\ \mathbb{A} & \overset{c \mapsto \left(\begin{smallmatrix} 0 & -c \\ 1 & 0 \end{smallmatrix}\right)}{\to} & \mathrm{M}_2 \end{array}$ are compatible with the right actions of the diagrams

$$\begin{array}{ccc} \mathrm{GRT}_{ell}(-) & \to & \mathrm{GRT}(-) \\ \downarrow & & \downarrow \\ \mathrm{GL}_2 & \overset{\det}{\to} & \mathbb{G}_m \end{array} \quad \text{and} \quad \begin{array}{ccc} \mathrm{GRT}(-) & \to & \mathrm{GRT}_{ell}(-) \\ \downarrow & & \downarrow \\ \mathbb{G}_m & \overset{c \mapsto \left(\begin{smallmatrix} 1 & 0 \\ 0 & c \end{smallmatrix}\right)}{\to} & \mathrm{GL}_2 \end{array}$$

6.6. Lie algebras. The graded Grothendieck-Teichmüller Lie algebra is[13]

$$\begin{aligned}\mathfrak{grt}_1 = \{ \quad & \psi \in \mathfrak{f}_2 | \psi + \psi^{3,2,1} = 0, \psi + \psi^{2,3,1} + \psi^{3,1,2} = 0, [t_{23}, \psi^{1,2,3}] + [t_{13}, \psi^{2,1,3}] = 0, \\ & \psi^{2,3,4} - \psi^{12,3,4} + \psi^{1,23,4} - \psi^{1,2,34} + \psi^{1,2,3} = 0\},\end{aligned}$$

where we use the inclusion $\mathfrak{f}_2 \subset \mathfrak{t}_3$, $A \mapsto t_{12}$, $B \mapsto t_{23}$; it is equipped with the Lie bracket $\langle \psi_1, \psi_2 \rangle = [\psi_1, \psi_2] + D_{\psi_2}(\psi_1) - D_{\psi_1}(\psi_2)$, where $D_\psi : A \mapsto [\psi, A]$, $B \mapsto 0$.

The Lie algebra $\mathbb{Q}$ acts on $\mathfrak{grt}_1$ by $[1, \psi] = -(\deg \psi)\psi$ (where $\deg A = \deg B = 1$), and we set $\mathfrak{grt} := \mathfrak{grt}_1 \rtimes \mathbb{Q}$.

The Lie algebras $\mathfrak{grt}_{(1)}$ are $\mathbb{N}$-graded (where deg is extended to be 0 on $\mathbb{Q}$), we then have $\mathrm{Lie}\,\mathrm{GRT}_{(1)}(-) = \widehat{\mathfrak{grt}}_{(1)}$ (the degree completions).

Let

$$\begin{aligned}\mathfrak{grt}_1^{ell} := \{ \quad & (\psi, \alpha_\pm) \in \mathfrak{f}_2 \times (\mathfrak{t}_{1,2})^2 | \psi \in \mathfrak{grt}_1, \\ & \alpha_\pm^{1,23} + \alpha_\pm^{2,31} + \alpha_\pm^{3,12} + [x_\pm^1, \psi^{1,2,3}] + [x_\pm^2, \psi^{2,1,3}] = 0, \\ & [x_\pm^1, \alpha_\pm^{3,12}] + [\alpha_\pm^{1,23}, x_\pm^3] - [x_\pm^1, [x_\pm^3, \psi^{1,2,3}]] = 0, \\ & [x_+^1, \alpha_-^{2,13}] - [x_-^2, \alpha_+^{1,23}] - [x_-^2, [x_+^1, \psi^{1,2,3}]] = 0\}.\end{aligned}$$

It is a Lie algebra when equipped with the bracket

$$[(\psi_1, \alpha_1^\pm), (\psi_2, \alpha_2^\pm)] = (\langle \psi_1, \psi_2 \rangle, D_{\alpha_2^\pm}(\alpha_1^\pm) - D_{\alpha_1^\pm}(\alpha_2^\pm)),$$

where for $\alpha_\pm \in \mathfrak{t}_{1,2}$, $D_{\alpha_\pm} \in \mathrm{Der}(\mathfrak{t}_{1,2})$ is given by $x_1^\pm \mapsto \alpha_\pm$.

The Lie algebra $\mathbb{Q}e_{22}$ acts on $\mathfrak{grt}_1^{ell}$ by $[e_{22}, (\psi, \alpha_+, \alpha_-)] = (-(\deg \psi)\psi, -(\deg_- \alpha_+)\alpha_+, (1 - \deg_- \alpha_-)\alpha_-)$, where $\deg \psi$ is as above, and $\deg_- \alpha_\pm$ is defined by $\deg_- x_1^+ = 0$, $\deg_- x_1^- = 1$. We then set $\mathfrak{grt}_{ell} := \mathfrak{grt}_1^{ell} \rtimes \mathbb{Q}e_{22}$.

The Lie algebras $\mathfrak{grt}_{(1)}^{ell}$ are $\mathbb{N}$-graded, where $(\psi, \alpha_\pm)$ has degree n if $2 \deg \psi = \deg \alpha_\pm - 1 = n$ ($\deg \alpha_\pm$ being defined by $\deg x_1^\pm = 1$ and $\deg \psi$ by $\deg t_{12} = \deg t_{23} = 1$) and e_{22} has degree 0. Then $\mathrm{Lie}\,\mathrm{GRT}_{(1)}^{ell}(-) = \widehat{\mathfrak{grt}}_{(1)}^{ell}$.

We have a morphism $\mathfrak{grt}_1^{ell} \to \mathfrak{sl}_2$, $(\psi, \alpha_+, \alpha_-) \mapsto \left(\begin{smallmatrix} a_+ & b_+ \\ a_- & b_- \end{smallmatrix}\right)$, where $\alpha_+ \equiv a_\pm x_1 + b_\pm y_1$ modulo degree ≥ 2. It extends to a morphism $\mathfrak{grt}_{ell} \to \mathfrak{gl}_2$ via $e_{22} \mapsto \left(\begin{smallmatrix} 0 & 0 \\ 0 & 1 \end{smallmatrix}\right)$. We denote by $\mathfrak{grt}_{I_2}^{ell}$ the common kernel of these morphisms; it coincides with the part of $\mathfrak{grt}_{(1)}^{ell}$ of positive degree.

These morphisms admit sections $\mathfrak{sl}_2 \to \mathfrak{grt}_1^{ell}$ given by $\left(\begin{smallmatrix} a_+ & b_+ \\ a_- & b_- \end{smallmatrix}\right) \mapsto (0, a_\pm x_1 + b_\pm y_1)$ and $\mathfrak{gl}_2 \to \mathfrak{grt}_{ell}$ given by its extension by $\left(\begin{smallmatrix} 0 & 0 \\ 0 & 1 \end{smallmatrix}\right) \mapsto e_{22}$. We then have $\mathfrak{grt}_1^{ell} \simeq \mathfrak{grt}_{I_2}^{ell} \rtimes \mathfrak{sl}_2$, $\mathfrak{grt}_{ell} \simeq \mathfrak{grt}_{I_2}^{ell} \rtimes \mathfrak{gl}_2$.

$\mathbb{Z}^2$-gradings may be defined on $\mathfrak{grt}_{(1)}^{ell}$ as follows. We have a Lie algebra inclusion $\mathfrak{grt}_1^{ell} \subset \mathfrak{grt}_1 \oplus \mathrm{Der}(\mathfrak{t}_{1,2}) =: \mathfrak{G}$. Recall that $\mathfrak{grt}_1$ is $\mathbb{N}$-graded while $\mathrm{Der}(\mathfrak{t}_{1,2})$ is $\mathbb{Z}^2$-graded by the $\mathbb{Z}^2$-grading of $\mathfrak{t}_{1,2}$ given by $(\deg_+, \deg_-)(x_1^+) = (1, 0)$, $(\deg_+, \deg_-)(x_1^-) = (0, 1)$. We then define a $\mathbb{Z}^2$-grading on $\mathfrak{G}$ by $\mathfrak{G}[p, q] := \begin{cases} \mathrm{Der}(\mathfrak{t}_{1,2})[p, q] & \text{if } q \neq p \\ \mathfrak{grt}_1[p] \oplus \mathrm{Der}(\mathfrak{t}_{1,2})[p, p] & \text{if } q = p \end{cases}$. This restricts to a $\mathbb{Z}^2$-grading $(\deg_+, \deg_-)$ of $\mathfrak{grt}_1^{ell}$, which extends to $\mathfrak{grt}_{ell}$ by $(\deg_+, \deg_-)(e_{22}) = (0, 0)$.

The $\mathbb{Z}^2$-grading of $\mathfrak{grt}_{ell}$ is compatible with the action of the Cartan subalgebra of $\mathfrak{gl}_2$: we have $[e_{11}, x] = -(\deg_+ x)x$, $[e_{22}, x] = -(\deg_- x)x$ for $e_{11} = \left(\begin{smallmatrix} 1 & 0 \\ 0 & 0 \end{smallmatrix}\right) \in \mathfrak{gl}_2$ and $x \in \mathfrak{grt}_{ell}$ homogeneous.

We have a morphism $\mathfrak{grt}_1^{ell} \to \mathfrak{grt}_1$, $(\psi, u_\pm) \mapsto \psi$. It extends to a morphism $\mathfrak{grt}_{ell} \to \mathfrak{grt}$ by $e_{22} \mapsto 1$. Using Proposition 53, sections of these morphisms are constructed as follows:

Proposition 61. *There is a unique Lie algebra morphism* $\mathfrak{grt}_1 \to \widehat{\mathfrak{grt}}_1^{ell}$, $\psi \mapsto (\psi, u_+^\psi, u_-^\psi)$, *where* $u_\pm^\psi := D_\psi(x_1^\pm)$ *and* $D_\psi \in \mathrm{Der}(\hat{\mathfrak{t}}_{1,2})$ *is defined by* $D_\psi(e^{x_1}) = \psi^{0,2,1} e^{x_1} - e^{x_1} \psi^{0,1,2}$, $D_\psi(t_{01}) =$

---

[13] As before, $\mathfrak{f}_2 = \mathfrak{f}_2^{\mathbb{Q}}$, etc.

$[\psi^{0,1,2}, t_{01}]$. *It extends to a Lie algebra morphism* $\mathfrak{grt} \to \widehat{\mathfrak{grt}}_{ell}$ *by* $1 \mapsto e_{22}$. *It is homogeneous,* $\mathfrak{grt}$ *being equipped with its degree and* $\widehat{\mathfrak{grt}}_{ell}$ *with degree* $\deg_-$.

Set now $\mathfrak{r}_{ell}^{gr} := \mathrm{Ker}(\mathfrak{grt}_{ell} \to \mathfrak{grt})$. We have

$$\begin{aligned}\mathfrak{r}_{ell}^{gr} = \{(\alpha_+, \alpha_-) \in (\mathfrak{t}_{1,2})^2 | \qquad & \alpha_\pm^{1,23} + \alpha_\pm^{2,31} + \alpha_\pm^{3,12} = 0, \\ & [x_\pm^1, \alpha_\pm^{2,13}] + [\alpha_\pm^{1,23}, x_\pm^2] = 0, \\ & [x_+^1, \alpha_-^{2,13}] + [\alpha_+^{1,23}, x_-^2] = 0\} \subset \mathrm{Der}(\mathfrak{t}_{1,2})^{op}.\end{aligned}$$

This is a $\mathbb{Z}^2$-graded Lie subalgebra of $\mathfrak{grt}_{ell}$; it is also $\mathbb{N}$-graded by $\deg_+ + \deg_-$. We have $\mathfrak{r}_{ell}^{gr}[0] \simeq \mathfrak{sl}_2$ and $\mathfrak{r}_{ell}^{gr} \simeq (\oplus_{d>0} \mathfrak{r}_{ell}^{gr}[d]) \rtimes \mathfrak{sl}_2$. Its completion for the $\mathbb{N}$-degree is isomorphic to $\mathrm{Lie}\, R_{ell}^{gr}(-)$.

Define a partial completion $\hat{\mathfrak{r}}_{ell}^{gr} := \oplus_q (\prod_p \mathfrak{r}_{ell}^{gr}[p,q])$. Proposition 61 gives rise to a Lie algebra morphism $\mathfrak{grt} \to \mathrm{Der}(\hat{\mathfrak{r}}_{ell}^{gr})$. We then have $\hat{\mathfrak{grt}}_{ell} \simeq \hat{\mathfrak{r}}_{ell}^{gr} \rtimes \mathfrak{grt}$, where $\hat{\mathfrak{grt}}_{ell} := \oplus_q \prod_p \mathfrak{grt}_{ell}[p,q]$ is a partial completion.

On the other hand, note that $\mathfrak{gt}_{ell} := \mathrm{Lie}\,\mathrm{GT}_{ell}(-)$ is filtered as follows: $\mathfrak{gt}_{ell} = \mathfrak{gt}_1^{ell} \rtimes \mathbb{Q}$, where $\mathfrak{gt}_1^{ell} := \mathrm{Lie}\,\mathrm{GT}_1^{ell}(-) \subset \hat{\mathfrak{f}}_2 \times (\hat{\mathfrak{t}}_{1,2})^2$. We then set $(\mathfrak{gt}_1^{ell})^{\geq 0} := \mathfrak{gt}_1^{ell}$, $(\mathfrak{gt}_1^{ell})^{\geq n} := \hat{\mathfrak{f}}_2^{\geq n/2} \times ((\hat{\mathfrak{t}}_{1,2})^2)^{\geq n+1}$ for $n > 1$, where the degree in $\hat{\mathfrak{f}}_2$ is induced by $\deg(t_{12}) = \deg(t_{23}) = 1$ and the degree in $\hat{\mathfrak{t}}_{1,2}$ by $\deg(x_1^\pm) = 1$. By inspecting relations, we have a canonical morphism $\mathrm{gr}(\mathfrak{gt}_{ell}) \to \mathfrak{grt}_{ell}$, restricting to $\mathrm{gr}(\mathfrak{r}_{ell}) \to \mathrm{gr}(\mathfrak{r}_{ell}^{gr})$ and compatible with $\mathrm{gr}(\mathfrak{gt}) \to \mathfrak{grt}$. We will see that these are isomorphisms.

Remark 62. We have then $\mathrm{GRT}_{(1)}(\mathbf{k}) = \exp(\widehat{\mathfrak{grt}_1}^{\mathbf{k}})(\rtimes \mathbf{k}^\times)$, $\mathrm{GRT}_{(1)}^{ell} = \exp(\widehat{\mathfrak{grt}_{I_2}}^{ell,\mathbf{k}}) \rtimes \mathrm{G(S)L}_2(\mathbf{k})$, $R_{ell}^{gr}(\mathbf{k}) = \exp(\prod_{d>0} \mathfrak{r}_{ell}^{gr}[d] \otimes \mathbf{k}) \rtimes \mathrm{SL}_2(\mathbf{k})$.

Remark 63. Any $(\alpha_+, \alpha_-) \in \mathfrak{r}_{ell}^{gr}$ satisfies $\alpha_\pm + \alpha_\pm^{2,1} = 0$, which implies that the total degree (in which $x_1^\pm$ have degree 1) of $\alpha_\pm$ is odd. So $\mathfrak{r}_{ell}^{gr}[d] = 0$ unless d is even.

6.7. A Lie subalgebra $\mathfrak{b}_3 \subset \mathfrak{r}_{ell}^{gr}$.

Proposition 64. *For* $n \geq 0$, *set*

$$\delta_{2n} := (\alpha_+ = \mathrm{ad}(x_1)^{2n+2}(y_1), \alpha_- = \frac{1}{2} \sum_{\substack{0 \leq p \leq 2n+1, \\ p+q=2n+1}} (-1)^p [(\mathrm{ad}\, x_1)^p(y_1), (\mathrm{ad}\, x_1)^q(y_1)]).$$

Then $\delta_{2n} \in \mathfrak{r}_{ell}^{gr}[2n+1, 1]$. δ_0 *is central in* $\mathfrak{grt}_{ell}$ *and coincides with* $\mathrm{ad}\, t_{12}$ *as an element of* $\mathrm{Der}(\mathfrak{t}_{1,2})^{op}$.

Proof. In [CEE], Proposition 3.1, we constructed derivations $\dot{\delta}_{2n}^{(m)} \in \mathrm{Der}(\mathfrak{t}_{1,m})$, such that

$$\dot{\delta}_{2n}^{(m)} : x_i \mapsto 0, t_{ij} \mapsto [t_{ij}, (\mathrm{ad}\, x_i)^{2n}(t_{ij})], y_i \mapsto \sum_{j:j \neq i} \frac{1}{2} \sum_{p+q=2n-1} [(\mathrm{ad}\, x_i)^p(t_{ij}), (-\mathrm{ad}\, x_i)^q(t_{ij})].$$

Let then $\delta_{2n}^{(m)} := \dot{\delta}_{2n}^{(m)} + [\sum_{i<j} (\mathrm{ad}\, x_i)^{2n}(t_{ij}), -]$. Then

$$\delta_{2n}^{(m)}(x_i) = [\sum_{j \neq i} (\mathrm{ad}\, x_i)^{2n}(t_{ij}), x_i] = (\mathrm{ad}\, x_i)^{2n+2}(y_i) = \alpha_-^{i,1\ldots\check{i}\ldots n}, \quad \delta_{2n}^{(m)}(t_{ij}) = 0,$$

$$\begin{aligned}
&\delta_{2n}^{(m)}(y_i) = \dot{\delta}_{2n}^{(m)}(y_i) + [\sum_{j<k} (\operatorname{ad} x_j)^{2n}(t_{jk}), y_i] \\
&= \dot{\delta}_{2n}^{(m)}(y_i) + \sum_{j\neq i} [(\operatorname{ad} x_i)^{2n}(t_{ij}), y_i] + \sum_{j<k; j,k\neq i} [(\operatorname{ad} x_j)^{2n}(t_{jk}), y_i] \\
&= \dot{\delta}_{2n}^{(m)}(y_i) + \sum_{j\neq i} [(\operatorname{ad} x_i)^{2n}(t_{ij}), y_i] + \sum_{j<k; j,k\neq i} \sum_{p+q=2n-1} (\operatorname{ad} x_j)^{p}[t_{ij}, (\operatorname{ad} x_j)^{q}(t_{jk})] \\
&= \dot{\delta}_{2n}^{(m)}(y_i) + \sum_{j\neq i} [(\operatorname{ad} x_i)^{2n}(t_{ij}), y_i] - \sum_{j<k; j,k\neq i} \sum_{p+q=2n-1} [(-\operatorname{ad} x_i)^{p}(t_{ij}), (\operatorname{ad} x_i)^{q}(t_{ik})] \\
&= \dot{\delta}_{2n}^{(m)}(y_i) + \sum_{j\neq i} [(\operatorname{ad} x_i)^{2n}(t_{ij}), y_i] - \frac{1}{2} \sum_{p+q=2n-1} [\sum_{j\neq i} (-\operatorname{ad} x_i)^{p}(t_{ij}), \sum_{k\neq i} (\operatorname{ad} x_i)^{q}(t_{ik})] \\
&+ \frac{1}{2} \sum_{j\neq i} \sum_{p+q=2n-1} [(-\operatorname{ad} x_i)^{p}(t_{ij}), (\operatorname{ad} x_i)^{q}(t_{ij})] \\
&= -[(\operatorname{ad} x_i)^{2n+1}(y_i), y_i] + \frac{1}{2} \sum_{p+q=2n-1} [(-\operatorname{ad} x_i)^{p+1}(y_i), (\operatorname{ad} x_i)^{q+1}(y_i)] \\
&= \frac{1}{2} \sum_{p+q=2n+1} [(-\operatorname{ad} x_i)^{p}(y_i), (\operatorname{ad} x_i)^{q}(y_i)] = \alpha_-^{i,1\ldots\check{i}\ldots n}.
\end{aligned}$$

Then $0 = \delta_{2n}^{(3)}([x_1^\pm, x_2^\pm]) = [x_1^\pm, \alpha_\pm^{2,13}] + [\alpha_\pm^{1,23}, x_2^\pm]$ and $0 = \delta_{2n}^{(3)}(t_{12}) = [x_1^+, \alpha_-^{2,13}] + [\alpha_+^{1,23}, x_2^-]$, which implies that $\delta_{2n} \in \mathfrak{r}_{ell}^{gr}$. □

We define $\mathfrak{b}_3 := \langle \mathfrak{sl}_2, \delta_{2n} \rangle \subset \mathfrak{r}_{ell}^{gr}$ as the Lie subalgebra generated by $\mathfrak{sl}_2$ and the δ_{2n}. This Lie algebra is $\mathbb{N}$-graded, and corresponds to the subgroup $\exp(\hat{\mathfrak{b}}_3^{+,\mathbf{k}}) \rtimes \mathrm{SL}_2(\mathbf{k}) \subset R_{ell}^{gr}(\mathbf{k})$ (where the hat denotes the degree completion and + means the positive degree part).

7. A MAP $\mathfrak{H} \to Ell(\mathbb{C})$

Recall that the KZ associator is defined as $\Phi_{KZ} = G_1(z)^{-1}G_0(z)$, where $G_i(z)$ are the analytic solutions of

$$G'(z) = (\frac{A}{z} + \frac{B}{z-1})G(z)$$

in $]0,1[$, valued in $\exp(\hat{\mathfrak{f}}_2^{\mathbb{C}})$, with asymptotic behavior $G_0(z) \sim z^A$ as $z \to 0$ and $G_1(z) \sim (1-z)^B$ as $z \to 1$. Then[14] $(2\pi\mathrm{i}, \Phi_{KZ}) \in M(\mathbb{C})$. We set $Ell_{KZ} := Ell(\mathbb{C}) \times_{M(\mathbb{C})} \{(2\pi\mathrm{i}, \Phi_{KZ})\}$.

We let $\mathfrak{H} := \{\tau \in \mathbb{C} | \Im(\tau) > 0\}$. We will define and study a map $e : \mathfrak{H} \to Ell_{KZ}$.

7.1. Definition of $e(\tau)$.

We let $(z,\tau) \mapsto \theta(z|\tau)$ be the holomorphic function on $\mathbb{C} \times \mathfrak{H}$, such that $\theta(z+1|\tau) = -\theta(z|\tau) = \theta(-z|\tau)$, $\theta(z+\tau|\tau) = -e^{2\pi\mathrm{i}z\tau + \mathrm{i}\pi\tau}\theta(z|\tau)$, $\{z|\theta(z|\tau) = 0\} = \mathbb{Z} + \tau\mathbb{Z}$, $\partial_z\theta(0|\tau) = 1$.

For $\tau \in \mathfrak{H}$, we let $F(z|\tau)$ be the holomorphic function on $\{z = a + b\tau |\ a \text{ or } b \in]0,1[\}$, valued in $\exp(\hat{\mathfrak{t}}_{1,2}^{\mathbb{C}})$, such that

$$\partial_z F(z|\tau) = -\frac{\theta(z + (2\pi\mathrm{i})^{-1}\operatorname{ad} x|\tau)\operatorname{ad} x}{\theta(z|\tau)\theta((2\pi\mathrm{i})^{-1}\operatorname{ad} x|\tau)}(y) \cdot F(z|\tau) \text{ and } F(z|\tau) \sim z^t \text{ as } z \to 0;$$

here $x := x_2^+$, $y := x_2^-$, $t := t_{12}$. We then set

$$A_+(\tau) := F(z|\tau)^{-1}F(z+1|\tau), \quad A_-(\tau) := F(z|\tau)^{-1}e^x F(z+\tau|\tau).$$

Proposition 65. *$\tau \mapsto e(\tau) := (A_+(\tau), A_-(\tau))$ is a map $\mathfrak{H} \to Ell_{KZ}$.*

---

[14] We set $\mathrm{i} := \sqrt{-1}$.

Proof. In [CEE], Section 4.4, we introduced $\tilde{A}, \tilde{B} \in \exp(\hat{\mathfrak{t}}_{1,2}^{\mathbb{C}})$. We set $\tilde{A}_+ := \tilde{A}$, $\tilde{A}_- := \tilde{B}$, then

$$\tilde{A}_\pm = \exp(\frac{\tau}{2\pi\,\mathrm{i}}(e_- + \sum_{k\geq 0} \tilde{a}_{2k}\delta_{2k}^{(2)}))(2\pi\,\mathrm{i})^{\frac{1}{2}[e_+,e_-]}(A_\pm(\tau)),$$

where $\sum_{k\geq 0} \tilde{a}_{2k}t^{2k} = \frac{\pi^2}{(\sin \pi t)^2} - \frac{1}{t^2}$ and $e_-, \delta_{2k}^{(2)} \in \mathrm{Der}(\hat{\mathfrak{t}}_{1,2})$. So $(A_+(\tau), A_-(\tau))$ satisfies (22), (23), (26) in [CEE]. (22), (23) imply that $(A_+(\tau), A_-(\tau))$ satisfies (22). (26) implies that

$$(A_-(\tau)^{12,3}\{\Phi^{-1}\}(A_-(\tau)^{1,23})^{-1}\{\Phi\}, A_+(\tau)^{12,3}) = \{\Phi^{-1}e^{2\pi\,\mathrm{i}\,t_{23}}\Phi\}$$

and using (23) in [CEE], we rewrite this as

$$(\{e^{-\mathrm{i}\,\pi t_{12}}\Phi^{3,2,1}\}A_-(\tau)^{2,13}\{\Phi^{2,1,3}e^{-\mathrm{i}\,\pi t_{12}}\}, A_+(\tau)^{12,3}) = \{\Phi^{-1}e^{2\pi\,\mathrm{i}\,t_{23}}\Phi\},$$

which as in the proof of Proposition 44 implies that $(A_+(\tau), A_-(\tau))$ satisfies (23). ☐

7.2. **Properties of** $e(\tau)$**.** We now describe the behavior of the map $\tau \mapsto e(\tau)$ under the action of $\mathrm{SL}_2(\mathbb{Z})$. Recall that the central subgroup $\{\exp(\alpha\delta_0), \alpha \in \mathbb{C}\}$ acts on Ell_{KZ} by $(A_\pm)*\exp(\alpha\delta_0) = (e^{\alpha t_{12}}A_\pm e^{-\alpha t_{12}})$.

Proposition 66. *We have*

$$e(g\tau) \in \tilde{g} * e(\tau) * \begin{pmatrix} \gamma\tau+\delta & 0 \\ \gamma/(2\pi\,\mathrm{i}) & (\gamma\tau+\delta)^{-1} \end{pmatrix}\{(\gamma\tau+\delta)^{\delta_0}\} \tag{45}$$

for $\tilde{g} \in \widetilde{\mathrm{SL}_2\,\mathbb{Z}}$*, with image* $g = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z})$*. Here* $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbb{C})$ *acts on* $e = (A_+, A_-)$ *by* $e*\begin{pmatrix} a & b \\ c & d \end{pmatrix} := (A_+(ax+by, cx+dy), A_-(ax+by, cx+dy))$*, and for* $z \in \mathbb{C}^\times$*,* $\{z^{\delta_0}\} := \{\exp(u\delta_0) | e^u = z\}$ *is a torsor under* $\exp(2\pi\,\mathrm{i}\,\mathbb{Z}\delta_0)$*.*

Proof. For $g = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z})$, we set $a_g(\tau) := \begin{pmatrix} \gamma\tau+\delta & 0 \\ \gamma/(2\pi\,\mathrm{i}) & (\gamma\tau+\delta)^{-1} \end{pmatrix}$. Then $a_{g_1g_2}(\tau) = a_{g_2}(\tau)a_{g_1}(g_2\tau)$.

Let also $b_g(\tau) := (\gamma\tau+\delta)^{\delta_0}$ (a class in $e^{\mathbb{C}\delta_0}/e^{\mathbb{Z}\delta_0}$), then $b_{g_1g_2}(\tau) = b_{g_2}(\tau)b_{g_1}(g_2\tau)$. It follows that $\{\tilde{g} | (45) \text{ is satisfied}\} \subset \widetilde{\mathrm{SL}_2(\mathbb{Z})}$ is a subgroup; so it suffices to check (45) for $\tilde{g} = \Psi, \Theta$.

We let $K(z|\tau) := -\frac{\theta(z+(2\pi\,\mathrm{i})^{-1}\,\mathrm{ad}\,x|\tau)\,\mathrm{ad}\,x}{\theta(z|\tau)\theta((2\pi\,\mathrm{i})^{-1}\,\mathrm{ad}\,x|\tau)}(y)$ and let $A_{z_0}^{z_1}(\tau)$ be the solution of $\partial_{z_1}A_{z_0}^{z_1}(\tau) = K(z_1|\tau)A_{z_0}^{z_1}(\tau)$, $A_z^z(\tau) = 1$. Then

$$A_+(\tau) = \lim_{\varepsilon\to 0^+} \varepsilon^{-t}A_\varepsilon^{1+\varepsilon}(\tau)\varepsilon^t, \quad A_-(\tau) = \lim_{\varepsilon\to 0^+} \varepsilon^{-t}e^x A_\varepsilon^{\tau+\varepsilon}(\tau)\varepsilon^t,$$

where the chosen branches of $A_{z_0}^{z_1}(\tau)$ are as in Fig. 1. Then $K(z|\tau) = K(z|\tau+1)$ implies that $A_\varepsilon^{1+\varepsilon}(\tau+1) = A_\varepsilon^{1+\varepsilon}(\tau)$, $A_\varepsilon^{\tau+1+\varepsilon}(\tau+1) = A_{1+\varepsilon}^{\tau+1+\varepsilon}(\tau)A_\varepsilon^{1+\varepsilon}(\tau) = A_\varepsilon^{\tau+\varepsilon}(\tau)A_\varepsilon^{1+\varepsilon}(\tau)$ with the decomposition of Fig. 2, so $A_+(\tau+1) = A_+(\tau)$, $A_-(\tau+1) = A_-(\tau)A_+(\tau)$, so $e(\tau+1) = \Psi * e(\tau)$, which settles the case $g = \Psi$.

Fig. 1



Fig. 2



Let $w := -z/\tau$, then

$$\frac{\partial}{\partial w} - K(w| - \frac{1}{\tau}) = -\tau e^{-z\,\mathrm{ad}\,x}\begin{pmatrix} -\tau & 0 \\ \frac{-1}{2\pi\,\mathrm{i}} & \frac{-1}{\tau} \end{pmatrix}(\frac{\partial}{\partial z} - K(z|\tau)).$$

So

$$A_{w_0}^{w_1}(-1/\tau) = e^{-xz_1} \cdot \left(\begin{smallmatrix} -\tau & 0 \\ \frac{-1}{2\pi\,\mathrm{i}} & \frac{-1}{\tau} \end{smallmatrix} \right)(A_{z_0}^{z_1}(\tau)) \cdot e^{xz_0}.$$

Then

$$\begin{aligned} A_+(-1/\tau) &= \lim_{\varepsilon\to 0^+} (\varepsilon^{-t} A_\varepsilon^{1+\varepsilon}(-1/\tau)\varepsilon^t) \\ &= \lim_{\varepsilon\to 0^+} \left(\begin{smallmatrix} -\tau & 0 \\ \frac{-1}{2\pi\,\mathrm{i}} & \frac{-1}{\tau} \end{smallmatrix} \right)((-\tau)^t(-\varepsilon\tau)^{-t} e^{-x(1+\varepsilon)} A_{-\varepsilon\tau}^{-\tau-\varepsilon\tau}(\tau)(-\varepsilon\tau)^t(-\tau)^{-t}) \\ &= (-\tau)^t \cdot \left(\begin{smallmatrix} -\tau & 0 \\ \frac{-1}{2\pi\,\mathrm{i}} & \frac{-1}{\tau} \end{smallmatrix} \right)(A_-(\tau)^{-1}) \cdot (-\tau)^{-t}, \end{aligned}$$

see Fig. 3; and

$$\begin{aligned} A_-(-1/\tau) &= \lim_{\varepsilon\to 0^+} \varepsilon^{-t} e^x A_\varepsilon^{(-1/\tau)+\varepsilon}(-1/\tau)\varepsilon^t \\ &= \lim_{\varepsilon\to 0^+} \varepsilon^{-t} e^x \cdot \left(\begin{smallmatrix} -\tau & 0 \\ \frac{-1}{2\pi\,\mathrm{i}} & \frac{-1}{\tau} \end{smallmatrix} \right)(e^{x(\frac{1}{\tau}-\varepsilon)} A_{-\varepsilon\tau}^{1-\varepsilon\tau}(\tau) e^{x\varepsilon}) \cdot \varepsilon^t \\ &= \lim_{\varepsilon\to 0^+} (-\tau)^t(-\tau\varepsilon)^{-t} \cdot \left(\begin{smallmatrix} -\tau & 0 \\ \frac{-1}{2\pi\,\mathrm{i}} & \frac{-1}{\tau} \end{smallmatrix} \right)(A_{-\tau\varepsilon}^{1-\tau\varepsilon}(\tau)) \cdot (-\tau\varepsilon)^t(-\tau)^{-t} \\ &= (-\tau)^t \cdot \left(\begin{smallmatrix} -\tau & 0 \\ \frac{-1}{2\pi\,\mathrm{i}} & \frac{-1}{\tau} \end{smallmatrix} \right)(A_-(\tau)A_+(\tau)A_-(\tau)^{-1}) \cdot (-\tau)^{-t}, \end{aligned}$$

see Fig. 4. The result for $g = \Theta$ then follows. ☐

Fig. 3



Fig. 4



Proposition 67. *We have*

$$\frac{\partial}{\partial\tau} e(\tau) = e(\tau) * (-2\pi\,\mathrm{i})(e_- + \sum_{k\geq 0} a_{2k} E_{2k+2}(\tau)\delta_{2k}),$$

where $E_{2k+2}(\tau)$ is the Eisenstein series of weight $2k+2$, equal to 1 *at* $\mathrm{i}\,\infty$.

Proof. In [CEE], we constructed a function $F^{(2)}(z|\tau)$, valued in $\exp(\hat{\mathfrak{t}}_{1,2}^{\mathbb{C}}) \rtimes \mathrm{Aut}(\hat{\mathfrak{t}}_{1,2}^{\mathbb{C}})$, such that $\partial_z F^{(2)}(z|\tau) = \tilde{K}(z|\tau)F^{(2)}(z|\tau)$ and

$$\frac{\partial}{\partial\tau} F^{(2)}(z|\tau) = \frac{-1}{2\pi\,\mathrm{i}}(e_- + \sum_{k\geq 0} \tilde{a}_{2k} E_{2k+2}(\tau)\dot{\delta}_{2k}^{(2)} - g(z, \mathrm{ad}\, x|\tau)(t))F^{(2)}(z|\tau),$$

where $\tilde{K}(z|\tau) = -\frac{\theta(z+\mathrm{ad}\,x|\tau)\,\mathrm{ad}\,x}{\theta(z|\tau)\theta(\mathrm{ad}\,x|\tau)}(y)$ and $g(z,x|\tau) = \frac{\theta(z+x|\tau)}{\theta(z|\tau)\theta(x|\tau)}(\frac{\theta'}{\theta}(z+x|\tau) - \frac{\theta'}{\theta}(z|\tau)) + \frac{1}{x^2}$, with $F^{(2)}(z|\tau) \sim z^t \exp(\frac{-\tau}{2\pi\,\mathrm{i}}(e_- + \sum_{k\geq 0} a_{2k}\delta_{2k}^{(2)}))$ (in the notation of *loc. cit.*, $e_- = \Delta_0$).

Then $F(z|\tau) = F^{(2)}(z|\tau)\varphi(\tau)$, where $\varphi(\tau)$ takes values in $\exp(\hat{\mathfrak{t}}_{1,2}^{\mathbb{C}}) \rtimes \mathrm{Aut}(\hat{\mathfrak{t}}_{1,2}^{\mathbb{C}})$. It follows that for some function $\xi(\tau)$ valued in $\hat{\mathfrak{t}}_{1,2}^{\mathbb{C}} \rtimes \mathrm{Der}(\hat{\mathfrak{t}}_{1,2}^{\mathbb{C}})$, we have

$$\frac{\partial}{\partial\tau} F(z|\tau) = \frac{-1}{2\pi\,\mathrm{i}}(e_- + \sum_{k\geq 1} \tilde{a}_{2k} E_{2k+2}(\tau)\dot{\delta}_{2k}^{(2)} - g(z, \mathrm{ad}\, x|\tau)(t))F(z|\tau) - F(z|\tau)\xi(\tau).$$

Define $\eta(\tau)$ by

$$\xi(\tau) = \frac{-1}{2\pi\,\mathrm{i}}(e_- + \sum_{k\geq 0} \tilde{a}_{2k} E_{2k+2}(\tau)\delta^{(2)}_{2k}) + \eta(\tau),$$

then as $-g(0, \mathrm{ad}\,x|\tau)(t) = \sum_{k\geq 0} \tilde{a}_{2k} E_{2k+2}(\tau)(\mathrm{ad}\,x)^{2k}(t)$ we get

$$\frac{\partial}{\partial\tau} F(z|\tau) = \frac{-1}{2\pi\,\mathrm{i}}(e_- + \sum_{k\geq 0} \tilde{a}_{2k} E_{2k+2}(\tau)\delta_{2k})(F(z|\tau)) \tag{46}$$
$$+\frac{1}{2\pi\,\mathrm{i}}\big(g(z, \mathrm{ad}\,x|\tau)(t) - g(0, \mathrm{ad}\,x|\tau)(t)\big)F(z|\tau) - F(z|\tau)\eta(\tau).$$

Taking into account $F(z|\tau) \sim z^t$ as $z \to 0$ for any τ, we get $\eta(\tau) = 0$. Define $\tilde{A}^{z_1}_{z_0}(\tau)$ by $\tilde{A}^z_z(\tau) = 1$ and $\partial_{z_1}\tilde{A}^{z_1}_{z_0}(\tau) = \tilde{K}(z_1|\tau)\tilde{A}^{z_1}_{z_0}(\tau)$. (46) then implies, with $g(z|\tau) := g(z, \mathrm{ad}\,x|\tau)(t) - g(0, \mathrm{ad}\,x|\tau)(t)$,

$$\partial_\tau \tilde{A}^{z_1}_{z_0}(\tau) = \frac{-1}{2\pi\,\mathrm{i}}(e_- + \sum_{k\geq 0} \tilde{a}_{2k} E_{2k+2}(\tau)\delta_{2k})(\tilde{A}^{z_1}_{z_0}(\tau)) + \frac{1}{2\pi\,\mathrm{i}} g(z_1|\tau)\tilde{A}^{z_1}_{z_0}(\tau) - \frac{1}{2\pi\,\mathrm{i}}\tilde{A}^{z_1}_{z_0}(\tau) g(z_0|\tau).$$

$\varepsilon^{-t}\tilde{A}^{1+\varepsilon}_{\varepsilon}(\tau)\varepsilon^t$ satisfies the same equation, with $g(z_1|\tau)$ (resp., $g(z_0|\tau)$) replaced by $\varepsilon^{-t}g(\varepsilon|\tau)\varepsilon^t$ and $\varepsilon^{-t}g(\varepsilon + 1|\tau)\varepsilon^t$, which both tend to 0 as $\varepsilon \to 0$. It follows that $\partial_\tau\tilde{A}_+(\tau) = \frac{-1}{2\pi\,\mathrm{i}}(e_- + \sum_{k\geq 0} \tilde{a}_{2k} E_{2k+2}(\tau)\delta_{2k})(\tilde{A}_+(\tau))$.

Similarly,

$$\partial_\tau \tilde{A}^{\tau+\varepsilon}_{\varepsilon}(\tau) = \frac{-1}{2\pi\,\mathrm{i}}(e_- + \sum_{k\geq 0} \tilde{a}_{2k} E_{2k+2}(\tau)\delta_{2k})(\tilde{A}^{\tau+\varepsilon}_{\varepsilon}(\tau))$$
$$+\Big(\frac{1}{2\pi\,\mathrm{i}} g(\tau+\varepsilon|\tau) - \frac{\theta(\tau+\varepsilon+\mathrm{ad}\,x|\tau)\,\mathrm{ad}\,x}{\theta(\tau+\varepsilon|\tau)\theta(\mathrm{ad}\,x|\tau)}(y)\Big)\tilde{A}^{\tau+\varepsilon}_{\varepsilon}(\tau) - \tilde{A}^{\tau+\varepsilon}_{\varepsilon}(\tau)\frac{1}{2\pi\,\mathrm{i}} g(\varepsilon|\tau).$$

So

$$\begin{aligned}
&\partial_\tau(\varepsilon^{-t}e^{2\pi\,\mathrm{i}\,x}\tilde{A}^{\tau+\varepsilon}_{\varepsilon}(\tau)\varepsilon^t) \\
&= \frac{-1}{2\pi\,\mathrm{i}}(e_- + \sum_{k\geq 0} \tilde{a}_{2k} E_{2k+2}(\tau)\delta_{2k})(\varepsilon^{-t}e^{2\pi\,\mathrm{i}\,x}\tilde{A}^{\tau+\varepsilon}_{\varepsilon}(\tau)\varepsilon^t) \\
&+\Big(\,\mathrm{Ad}(\varepsilon^{-t}e^{2\pi\,\mathrm{i}\,x})\big(\frac{1}{2\pi\,\mathrm{i}} g(\tau+\varepsilon|\tau) - \frac{\theta(\tau+\varepsilon+\mathrm{ad}\,x|\tau)\,\mathrm{ad}\,x}{\theta(\tau+\varepsilon|\tau)\theta(\mathrm{ad}\,x|\tau)}(y)\big) \\
&+\varepsilon^{-t}e^{2\pi\,\mathrm{i}\,x}\frac{-1}{2\pi\,\mathrm{i}}(e_- + \sum_{k\geq 0} \tilde{a}_{2k} E_{2k+2}(\tau)\delta_{2k})(e^{-2\pi\,\mathrm{i}\,x})\varepsilon^t\Big) \cdot (\varepsilon^{-t}e^{2\pi\,\mathrm{i}\,x}\tilde{A}^{\tau+\varepsilon}_{\varepsilon}(\tau)\varepsilon^t) \\
&-\big(\varepsilon^{-t}e^{2\pi\,\mathrm{i}\,x}\tilde{A}^{\tau+\varepsilon}_{\varepsilon}(\tau)\varepsilon^t\big) \cdot \big(\frac{1}{2\pi\,\mathrm{i}}\varepsilon^{-t}g(\varepsilon|\tau)\varepsilon^t\big).
\end{aligned}$$

Identity (7) in [CEE] implies that the parenthesis in the third and fourth lines equals $\mathrm{Ad}(\varepsilon^{-t})(\frac{1}{2\pi\,\mathrm{i}}g(\varepsilon|\tau))$. As before, we get in the limit $\varepsilon \to 0$

$$\partial_\tau\tilde{A}_-(\tau) = \frac{-1}{2\pi\,\mathrm{i}}(e_- + \sum_{k\geq 0} \tilde{a}_{2k} E_{2k+2}(\tau)\delta_{2k})(\tilde{A}_-(\tau)).$$

As $a_{2k} = (2\pi\,\mathrm{i})^{2k+2}\tilde{a}_{2k}$ and $A_\pm(\tau) = \left(\begin{smallmatrix} 2\pi\,\mathrm{i} & 0 \\ 0 & (2\pi\,\mathrm{i})^{-1}\end{smallmatrix}\right)(\tilde{A}_\pm(\tau))$, we get $\partial_\tau A_\pm(\tau) = -2\pi\,\mathrm{i}(e_- + \sum_{k\geq 0} a_{2k}E_{2k+2}(\tau)\delta_{2k})(A_\pm(\tau))$. □

Proposition 68.

$$\sigma(\Phi_{KZ}) = \lim_{\tau\to\mathrm{i}\,\infty} e(\tau) * (\frac{\mathrm{i}}{2\pi})^{\delta_0}\exp(2\pi\,\mathrm{i}\,\tau(e_- + \sum_{k\geq 0} a_{2k}\delta_{2k})).$$

Proof. Let $e(\tau) = (A_+(\tau), A_-(\tau))$ and $\sigma(\Phi_{KZ}) = (A^+_{KZ}, A^-_{KZ})$. In [CEE], proof of Prop. 4.9, we show that $\lim_{\tau\to \mathrm{i}\,\infty} A_+(\tau) = (\mathrm{i}\,/2\pi)^{-\delta_0} A^+_{KZ}$; since $[e_- + \sum_{k\geq 0} a_{2k}\delta_{2k}, A^+_{KZ}] = 0$, we also have $\lim_{\tau\to \mathrm{i}\,\infty} e^{2\pi\,\mathrm{i}\,\tau(e_-+\sum_{k\geq 0} a_{2k}\delta_{2k})}(A_+(\tau)) = (\mathrm{i}\,/2\pi)^{-\delta_0} A^+_{KZ}$. In Lemma 4.14, we get an asymptotic expansion for $A_-(\tau)$ and at the end of Thm. 4.11, we compute $\lim_{\tau\to \mathrm{i}\,\infty} e^{2\pi\,\mathrm{i}\,\tau(e_-+\sum_{k\geq 0} a_{2k}\delta_{2k})}(A_-(\tau)) = (\mathrm{i}\,/2\pi)^{-\delta_0}(A^-_{KZ})$. □

8. Isomorphisms of Lie algebras

Let $\mathbf{k}$ be a $\mathbb{Q}$-ring. As $Ell(\mathbf{k})$ is a torsor, each $e \in Ell(\mathbf{k})$ gives rise ro an isomorphism $i_e : \mathrm{GT}_{ell}(\mathbf{k}) \to \mathrm{GRT}_{ell}(\mathbf{k})$, defined by $g * e = e * i_e(g)$ for any $g \in \mathrm{GT}_{ell}(\mathbf{k})$. Similarly, any $\Phi \in M(\mathbf{k})$ gives rise to an isomorphism $i_\Phi : \mathrm{GT}(\mathbf{k}) \to \mathrm{GRT}(\mathbf{k})$ defined by the same conditions. We then have a commutative diagram

$$\begin{tikzcd} \mathrm{GT}_{ell}(\mathbf{k}) \arrow[r, "i_e"] \arrow[d] & \mathrm{GRT}_{ell}(\mathbf{k}) \arrow[d] \\ \mathrm{GT}(\mathbf{k}) \arrow[r, "j_\Phi"] & \mathrm{GRT}(\mathbf{k}) \end{tikzcd} \tag{47}$$

where $\Phi = \mathrm{im}(e \in Ell(\mathbf{k}) \to M(\mathbf{k}))$. In particular, i_e restricts to an isomorphism $i_e : R_{ell}(\mathbf{k}) \to R^{gr}_{ell}(\mathbf{k})$. When $e \in \mathrm{im}(M(\mathbf{k}) \xrightarrow{\sigma} Ell(\mathbf{k}))$, the isomorphism $R_{ell}(\mathbf{k}) \xrightarrow{i_e} R^{gr}_{ell}(\mathbf{k})$ is compatible with i_Φ and the action of $\mathrm{G(R)T}(\mathbf{k})$ on both sides via the lifts $\mathrm{G(R)T}(\mathbf{k}) \xrightarrow{\sigma} \mathrm{G(R)T}_{ell}(\mathbf{k})$.

The isomorphisms i_e induce Lie algebra isomorphisms $\mathfrak{gt}^{\mathbf{k}}_{ell} \to \widehat{\mathfrak{grt}}^{\mathbf{k}}_{ell}$, restricting to $\mathfrak{r}^{\mathbf{k}}_{ell} \to \hat{\mathfrak{r}}^{gr,\mathbf{k}}_{ell}$, compatible with the filtrations and whose associated graded isomorphisms are the canonical morphisms from the end of Subsection 6.6. Since $Ell(\mathbb{Q}) \neq \emptyset$ (e.g., because it contains $\sigma(M(\mathbb{Q}))$), we obtain:

Proposition 69. *We have isomorphisms* $\mathfrak{gt}_{ell} \simeq \hat{\mathrm{gr}}(\mathfrak{gt}_{ell}) = \widehat{\mathfrak{grt}}_{ell}$ *and* $\mathfrak{r}_{ell} \simeq \hat{\mathrm{gr}}(\mathfrak{r}_{ell}) = \hat{\mathfrak{r}}^{gr}_{ell}$.

Composing (47) with

$$\begin{tikzcd} \widetilde{\mathrm{GL}_2\,\mathbb{Z}} \arrow[r] \arrow[d] & \mathrm{GT}_{ell}(\mathbf{k}) \arrow[d] \\ \{\pm 1\} \arrow[r] & \mathrm{GT}(\mathbf{k}) \end{tikzcd}$$

we obtain a commutative diagram

$$\begin{tikzcd} \widetilde{\mathrm{GL}_2\,\mathbb{Z}} \arrow[r] \arrow[d] & \mathrm{GRT}_{ell}(\mathbf{k}) \arrow[d] \\ \{\pm 1\} \arrow[r] & \mathrm{GRT}(\mathbf{k}) \end{tikzcd}$$

inducing morphisms $B_3 = \widetilde{\mathrm{SL}_2\,\mathbb{Z}} \to R_{ell}(\mathbf{k}) \to R^{gr}_{ell}(\mathbf{k})$.

When $\mathbf{k} = \mathbb{C}$ and $e = e(\tau), e_{KZ}$, we now make explicit the morphism $B_3 \to R^{gr}_{ell}(\mathbb{C})$. We will see that its image is contained in $\exp(\hat{\mathfrak{b}}^{+,\mathbb{C}}_3) \rtimes \mathrm{SL}_2(\mathbb{C}) \subset R^{gr}_{ell}(\mathbb{C})$.

Define $F(\tau)$ as the map $\mathfrak{H} \to \exp(\hat{\mathfrak{b}}^{+,\mathbb{C}}_3) \rtimes \mathrm{SL}_2(\mathbb{C})$ such that

$$\partial_\tau F(\tau) = 2\pi\,\mathrm{i}(e_- + \sum_{k\geq 0} a_{2k} E_{2k+2}(\tau)\delta_{2k})F(\tau)$$

and $F(\tau) \sim e^{2\pi\,\mathrm{i}(e_-+\sum_{k\geq 0} a_{2k}E_{2k+2}(\tau)\delta_{2k})\tau}$ as $\tau \to \mathrm{i}\,\infty$. Then the map $\tau \mapsto e(\tau) * F(\tau)$ is a constant, and

$$e_{KZ} = e(\tau) * F(\tau) \text{ for any } \tau \in \mathfrak{H}. \tag{48}$$

Moreover, for any $\tilde{g} \in B_3$ with image $g = \left(\begin{smallmatrix} \alpha & \beta \\ \gamma & \delta \end{smallmatrix}\right) \in \mathrm{SL}_2(\mathbb{Z})$, we have

$$\tilde{g} * e(\tau) \equiv e(g\tau)\left(\begin{smallmatrix} \gamma\tau+\delta & 0 \\ \gamma/(2\pi\,\mathrm{i}) & (\gamma\tau+\delta)^{-1} \end{smallmatrix}\right)^{-1} = e(\tau) * F(\tau)F(g\tau)^{-1}\left(\begin{smallmatrix} \gamma\tau+\delta & 0 \\ \gamma/(2\pi\,\mathrm{i}) & (\gamma\tau+\delta)^{-1} \end{smallmatrix}\right)^{-1} = e(\tau) * i_{e(\tau)}(\tilde{g}), \tag{49}$$

where the second equality follows from (48) for τ and $g\tau$ ($\equiv$ means modulo $e^{\mathbb{C}\delta_0}$). It follows that

$$i_{e(\tau)}(\tilde{g}) \equiv F(\tau)F(g\tau)^{-1}\left(\begin{smallmatrix} \gamma\tau+\delta & 0 \\ \gamma/(2\pi\,\mathrm{i}) & (\gamma\tau+\delta)^{-1} \end{smallmatrix}\right)^{-1}.$$

Acting from the right by $F(\tau)$ in the equality between the first and the third term of (49), we get $\tilde{g} * e_{KZ} \equiv e_{KZ} * F(g\tau)^{-1}\left(\begin{smallmatrix} \gamma\tau+\delta & 0 \\ \gamma/(2\pi\,\mathrm{i}) & (\gamma\tau+\delta)^{-1}\end{smallmatrix}\right)^{-1}F(\tau)$, so

$$i_{e_{KZ}}(\tilde{g}) \equiv F(g\tau)^{-1}\left(\begin{smallmatrix} \gamma\tau+\delta & 0 \\ \gamma/(2\pi\,\mathrm{i}) & (\gamma\tau+\delta)^{-1}\end{smallmatrix}\right)^{-1}F(\tau)$$

for any $\tau \in \mathfrak{H}$. It follows that $i_{e(\tau)}, i_{e_{KZ}}$ are morphisms $B_3 \to \exp(\hat{\mathfrak{b}}_3^{+,\mathbb{C}}) \rtimes \mathrm{SL}_2(\mathbb{C})$. The composite morphism $B_3 \to \exp(\hat{\mathfrak{b}}_3^{+,\mathbb{C}}) \rtimes \mathrm{SL}_2(\mathbb{C}) \to \mathrm{SL}_2(\mathbb{C})$ is $\tilde{g} \mapsto \left(\begin{smallmatrix} \alpha & -2\pi\,\mathrm{i}\,\beta \\ -\gamma/(2\pi\,\mathrm{i}) & \delta\end{smallmatrix}\right)$.

Recall that $B_3 \subset \mathrm{Aut}(F_2)$ is generated by Ψ, Θ given by $\Psi : X, Y \mapsto X, YX$ and $\Theta : X, Y \mapsto XYX^{-1}, X^{-1}$. We also set $\Psi_- := \Theta\Psi\Theta^{-1} : X, Y \mapsto XY^{-1}, Y$.

Then

$$i_{e_{KZ}}(\Psi) = F(\tau+1)^{-1}F(\tau) = \exp(-2\pi\,\mathrm{i}(e_- + \sum_{k>0} a_{2k}\delta_{2k}))e^{\frac{2\pi\,\mathrm{i}}{12}\delta_0} =: \psi,$$

and since $i_{e_{KZ}}(\Theta) = \left(\begin{smallmatrix} 0 & -2\pi\,\mathrm{i} \\ (2\pi\,\mathrm{i})^{-1} & 0\end{smallmatrix}\right)$, $i_{e_{KZ}}(\Phi^-) = \exp\xi = \psi_-$, where $\xi_- = (2\pi\,\mathrm{i})^{-1}(e_+ +$element of $\hat{\mathfrak{b}}_3^{\mathbb{C},+})$. We have therefore

$$\Psi * (A^+_{KZ}, A^-_{KZ}) = (A^+_{KZ}, A^-_{KZ}A^+_{KZ}) = (A^+_{KZ} * \psi, A^-_{KZ} * \psi) = (A^+_{KZ}, A^-_{KZ}) * \psi,$$

and

$$\Psi_- * (A^+_{KZ}, A^-_{KZ}) = (A^+_{KZ}(A^-_{KZ})^{-1}, A^-_{KZ}) = (A^+_{KZ}, A^-_{KZ}) * \psi_-.$$

Remark 70. (Relation with the Drinfeld generators of $\mathfrak{grt}_1$) If $e \in \sigma(M(\mathbf{k}))$, then $B_3 \overset{i_e}{\to} R^{gr}_{ell}(\mathbf{k})$ is equivariant w.r.t. the actions of $\sigma(-1) \equiv \left(\begin{smallmatrix} 1 & 0 \\ 0 & -1\end{smallmatrix}\right)$ on $B_3 = \widetilde{\mathrm{SL}_2\,\mathbb{Z}}$ and of $\mathrm{im}(-1 \to \mathrm{GRT}(\mathbf{k}))$ on $R^{gr}_{ell}(\mathbf{k})$. Then $\mathrm{im}(-1 \in \mathrm{GT} \overset{j_{\Phi_{KZ}}}{\to} \mathrm{GRT}(\mathbb{C}))$ is an element $g_{KZ} = (h_{KZ}, -1)$ where $h_{KZ} \in \mathrm{GRT}_1(\mathbb{C})$; the homogeneous components of $\log h_{KZ}$ yield the Drinfeld generators of $\mathfrak{grt}_1^{\mathbb{C}}$. This equivariance is expressed as $(\psi_\pm)^{-1} = \sigma(g_{KZ})\psi_\pm\sigma(g_{KZ})^{-1}$, where $\psi_+ = \psi$. □

We then prove:

Proposition 71. *For $e = e_{KZ}$, the isomorphism $i_e : R_{ell}(\mathbb{C}) \to R^{gr}_{ell}(\mathbb{C})$ restricts to an isomorphism $\langle B_3\rangle(\mathbb{C}) \to \exp(\hat{\mathfrak{b}}_3^{+,\mathbb{C}}) \rtimes \mathrm{SL}_2(\mathbb{C})$.*

Proof. $i_e(\langle B_3\rangle(\mathbb{C}))$ is the Zariski closure of the subgroup of $R^{gr}_{ell}(\mathbb{C})$ generated by $\psi_\pm$. These are elements of $\exp(\hat{\mathfrak{b}}_3^{+,\mathbb{C}}) \rtimes \mathrm{SL}_2(\mathbb{C})$, which is Zariski closed, so $i_e(\langle B_3\rangle(\mathbb{C}))$ is contained in this group. On the other hand, the Lie algebra of this Zariski closure is the topological Lie algebra generated by $\log\psi_\pm$. It then suffices to prove that this Lie algebra coincides with $\hat{\mathfrak{b}}_3^{\mathbb{C}}$.

Equip $\mathrm{Aut}(F_2(\mathbb{C}))$ with the topology for which a system of neighborhoods of 1 is $\mathrm{Aut}^n(F_2(\mathbb{C})) = \{\theta | \forall g \in F_2(\mathbb{C}), \theta(g) \equiv g \text{ mod } F_2^{(n)}(\mathbb{C})\} \subset \mathrm{Aut}(F_2(\mathbb{C}))$, where $F_2^{(1)}(\mathbb{C}) = F_2(\mathbb{C})$ and $F_2^{(n)}(\mathbb{C}) = (F_2^{(n-1)}(\mathbb{C}), F_2(\mathbb{C}))$. This induces a topology on $R_{ell}(\mathbb{C})$, which we call the prounipotent topology as it may be induced by the inclusion $R_{ell}(\mathbb{C}) \subset F_2(\mathbb{C})^2$, $F_2(\mathbb{C})$ being equipped with this topology.

Lemma 72. *$\langle B_3\rangle(\mathbb{C}) \subset R_{ell}(\mathbb{C})$ is closed for this topology.*

Proof. We have $\langle B_3\rangle(\mathbb{C}) = \cap_{G\in\mathcal{G}} G$, where $\mathcal{G} = \{G(\mathbb{C}) | G \subset R_{ell}(-)$ is a subgroup scheme such that $G(\mathbb{Q}) \supset B_3\}$. It then suffices to show that each $G(\mathbb{C})$ is closed in the prounipotent topology. Define coordinates on $R_{ell}(\mathbb{C})$ as follows: $R_{ell}(\mathbb{C}) \ni \theta \leftrightarrow (c_b, d_b)_b$, where b runs over a homogeneous basis of $\mathfrak{f}_2$ (generated by $\xi = \log X$, $\eta = \log Y$), e.g., $\{b\} = \{\xi, \eta, [\xi, \eta], \dots\}$, and $\log\theta(e^\xi) = \sum_b c_b b$, $\log\theta(e^\eta) = \sum_b d_b b$. Then $G(\mathbb{C})$ is a finite intersection of sets of the form $\{\theta | P(c_\xi, c_\eta, \dots, d_\xi, d_\eta, \dots) = 0\}$, where P is a polynomial in $(c_b, d_b)_b$, vanishing at the origin. Such a $G(\mathbb{C})$ necessarily contains $R_{ell}(\mathbb{C}) \cap \mathrm{Aut}^n(F_2(\mathbb{C}))$ for a large enough n. □

It follows that $i_e(\langle B_3\rangle(\mathbb{C})) \subset \exp(\hat{\mathfrak{b}}_3^{+,\mathbb{C}}) \rtimes \mathrm{SL}_2(\mathbb{C})$ is closed in the prounipotent topology of $R^{gr}_{ell}(\mathbb{C})$ (as in the case of $R_{ell}(\mathbb{C})$, it is defined by the inclusion in $\mathrm{Aut}(\hat{\mathfrak{f}}_2^{\mathbb{C}})$), so $\mathrm{Lie}\, i_e(\langle B_3\rangle(\mathbb{C})) \subset \hat{\mathfrak{b}}_3^{\mathbb{C}}$ is closed.

Recall that $\log\psi = -2\pi\mathrm{i}(e_- + \sum_{k\geq 1} a_{2k}\delta_{2k}) + \frac{2\pi\mathrm{i}}{12}\delta_0 \in \mathrm{Lie}\, i_e(\langle B_3\rangle(\mathbb{C}))$, while $\log\psi_- \in (2\pi\mathrm{i})^{-1}(e_+ + \hat{\mathfrak{b}}_3^{+,\mathbb{C}})$.

Lemma 73. *Let* $\mathfrak{g} \subset \hat{\mathfrak{b}}_3^{\mathbb{C}}$ *be a closed (for the total degree topology) Lie subalgebra, such that* $\mathfrak{g} \ni \tilde{e}_\pm$*, where:* $\tilde{e}_+ = e_+ +$ *terms of degree* > 0, $\tilde{e}_- = e_- + \sum_{k>0} a_{2k}\delta_{2k} - \frac{1}{12}\delta_0 + \sum_{p\geq 1, q>1}$ *degree* (p,q)*. Then* $\mathfrak{g} = \hat{\mathfrak{b}}_3^{\mathbb{C}}$*.*

Proof. Set $\mathcal{G} = \oplus_{k\geq 0}\mathcal{G}_{2k} := \mathfrak{b}_3^{\mathbb{C}}$ (decomposition w.r.t. the total degree), $\hat{\mathcal{G}} := \hat{\mathfrak{b}}_3^{\mathbb{C}}$. Set $\hat{\mathcal{G}}_{\geq 2k} := \prod_{k'\geq k}\mathcal{G}_{2k'}$, then $\hat{\mathcal{G}} = \hat{\mathcal{G}}_{\geq 0} \supset \hat{\mathcal{G}}_{\geq 2} \supset \ldots$ is a complete descending Lie algebra filtration of $\hat{\mathcal{G}}$, with associated graded Lie algebra $\mathcal{G}$. Set $\mathfrak{g}_{\geq 2k} := \mathfrak{g} \cap \mathcal{G}_{2k}$, then $\mathfrak{g} = \mathfrak{g}_{\geq 0} \supset \mathfrak{g}_{\geq 2} \supset \ldots$ is a complete descending filtration of $\mathfrak{g}$. Let $\mathrm{gr}(\mathfrak{g}) := \oplus_{k\geq 0}\mathfrak{g}_{\geq 2k}$, where $\mathrm{gr}(\mathfrak{g}) := \mathfrak{g}_{\geq 2k}/\mathfrak{g}_{\geq 2(k+2)}$. We then have an inclusion $\mathrm{gr}(\mathfrak{g}) \subset \mathcal{G}$ of graded Lie algebras. We now prove that $\mathrm{gr}(\mathfrak{g}) = \mathcal{G}$.

As $\tilde{e}_\pm \in \mathfrak{g} = \mathfrak{g}_0$, $\mathrm{gr}(\mathfrak{g}_0)$ contains $e_\pm$. Set $h := [e_+, e_-]$. Then $[\tilde{e}_+, \tilde{e}_-] = h + \sum_{p,q\geq 1}$ terms of degree (p,q), and $[h, e_-] = -2e_-$, $[h, \delta_{2n}] = 2n\delta_{2n}$. Then $\mathfrak{g} \ni P(\mathrm{ad}[\tilde{e}_+, \tilde{e}_-])(\tilde{e}_-) = P(-2)\Delta_0 + \sum_{n\geq 0} a_{2n}P(2n)\delta_{2n} + \sum_{p\geq 1, q>1}$ terms of degree (p,q) (with $a_0 = -\frac{1}{12}$). Taking P such that $P(-2) = P(0) = \ldots = P(2k-2) = 0$ and $a_{2k}P(2k) = 1$, we see that $\mathfrak{g}$ contains an element of the form $\delta_{2k} + \sum_{p\geq 1, q>1}$ terms of degree (p,q). Applying $(\mathrm{ad}\,\tilde{e}_-)^{2k}$ to this element, and using the fact that $(\mathrm{ad}\,\tilde{e}_-)^{2k}(x) = 0$ for $x \in \mathcal{G}$ of total degree $\leq 2(k-1)$, we see that $\mathfrak{g}$ contains an element of the form $(\mathrm{ad}\, e_-)^{2k}(\delta_{2k}) + \sum$ terms of total degree $\geq 2(k+2)$. As the latter sum belongs to $\mathfrak{g}_{\geq 2(k+1)}$, we obtain that $(\mathrm{ad}\, e_-)^{2k}(\delta_{2k}) \in \mathrm{gr}(\mathfrak{g})_{2(k+1)}$. The Lie subalgebra $\mathrm{gr}(\mathfrak{g}) \subset \mathcal{G}$ then contains $e_\pm$ and $(\mathrm{ad}\, e_-)^{2k}(\delta_{2k})$, $k \geq 0$. As $(\mathrm{ad}\, e_-)^{2k+1}(\delta_{2k}) = 0$, $(\mathrm{ad}\, e_+)^{2k}(\mathrm{ad}\, e_-)^{2k}(\delta_{2k})$ is a nonzero multiple of δ_{2k}. So $\mathrm{gr}(\mathfrak{g}) = \mathcal{G}$. It follows that $\mathfrak{g} = \mathcal{G}$. □

End of proof of Proposition 71. Applying Lemma 73 with $\tilde{e}_+ = 2\pi\mathrm{i}\log\psi_-$, $\tilde{e}_- = -(2\pi\mathrm{i})^{-1}\log\psi$, we get $i_e(\mathrm{Lie}\langle B_3\rangle(\mathbb{C})) = \hat{\mathfrak{b}}_3^{\mathbb{C}}$, as wanted. □

9. CONJECTURES

9.1. **A generation conjecture (GC).** The Drinfeld generators of $\mathfrak{grt}_1$ are obtained from the homogeneous decomposition of the logarithm of $\mathrm{im}(-1 \in \underline{\mathrm{GT}} \overset{j_{\Phi_{KZ}}}{\to} \mathrm{GRT}(\mathbb{C})) \cdot \mathrm{can}(-1)$, where $\mathrm{can} : \mathbb{C}^\times \to \mathrm{GRT}(\mathbb{C})$ is the canonical morphism. The analogue of the conjecture that these elements generate $\mathfrak{grt}_1$ is then:

Conjecture 74. *(Generation Conjecture)* $\mathfrak{b}_3 \subset \mathfrak{r}^{gr}_{ell}$ *is an equality, i.e.,* $\mathfrak{r}^{gr}_{ell}$ *is generated by* $\mathfrak{sl}_2$ *and the* δ_{2n}, $n \geq 0$*.*

This conjecture is equivalent to the inclusion $\exp(\hat{\mathfrak{b}}_3^{+,\mathbf{k}}) \rtimes \mathrm{SL}_2(\mathbf{k}) \subset R^{gr}_{ell}(\mathbf{k})$ being an equality.

Proposition 75. *GC is equivalent to the Zariski density of* $B_3 \subset R_{ell}(-)$*, i.e.,* $\langle B_3\rangle = R_{ell}(-)$*.*

Proof. According to Lemma 33, $\langle B_3\rangle$ is uniquely determined by its Lie algebra. This fact and Proposition 32 immediately imply that $\langle B_3\rangle = R_{ell}(-)$ iff the inclusion $\mathrm{Lie}(u_+, u_-) = \langle\log\psi, \log\psi_-\rangle \subset \mathrm{Lie}\, R_{ell}(-)$ is actually an equality. Tensoring with $\mathbb{C}$, this holds iff $\langle\log\psi, \log\psi_-\rangle^{\mathbb{C}} \subset \hat{\mathfrak{r}}^{\mathbb{C}}_{ell}$ is an equality. On the other hand, GC holds iff $\hat{\mathfrak{b}}_3^{+,\mathbb{C}} \subset \hat{\mathfrak{r}}^{gr,\mathbb{C}}_{ell}$ is an equality. Now $i_{e_{KZ}}$ sets up a diagram

$$\begin{tikzcd}
\langle\log\psi, \log\psi_-\rangle^{\mathbb{C}} \arrow[r, hook] \arrow[d, "\simeq"'] & \mathfrak{r}^{\mathbb{C}}_{ell} \arrow[d, "\simeq"] \\
\hat{\mathfrak{b}}_3^{+,\mathbb{C}} \arrow[r, hook] & \hat{\mathfrak{r}}^{gr,\mathbb{C}}_{ell}
\end{tikzcd}$$

It follows that the upper inclusion is an equality iff the lower is. □

9.2. **Relation with a transcendence conjecture.** We first present the transcendence conjecture.

9.2.1. *The coordinate ring of associators.* The functors $\{\mathbb{Q}\text{-rings}\} \to \{\text{sets}\}$, $\mathbf{k} \mapsto \underline{M}(\mathbf{k}), M(\mathbf{k})$ may be represented as follows. Let $\mathrm{pent}_{\mathbf{k}} : \exp(\hat{\mathfrak{f}}_2^{\mathbf{k}}) \to \exp(\hat{\mathfrak{t}}_3^{\mathbf{k}})$, $\mathrm{hex} : \mathbf{k} \times \exp(\hat{\mathfrak{f}}_2^{\mathbf{k}}) \to \exp(\hat{\mathfrak{t}}_3^{\mathbf{k}})$, $\mathrm{dual} : \exp(\hat{\mathfrak{f}}_2^{\mathbf{k}}) \to \exp(\hat{\mathfrak{t}}_3^{\mathbf{k}})$ be the maps $\mathrm{pent}(\Phi) := \mathrm{lhs}((21))^{-1}\,\mathrm{rhs}((21))$, $\mathrm{hex}(\mu, \Phi) := \mathrm{lhs}((20))\,\mathrm{rhs}((20))^{-1}$, $\mathrm{dual}(\Phi) := \Phi\Phi^{3,2,1}$.

Let B, B', C be homogeneous bases of $\mathfrak{f}_2, \mathfrak{t}_3, \mathfrak{t}_4$. Let μ, ϕ_b $(b \in B)$ be free commutative variables and set $\mathbf{k}_0 := \mathbb{Q}[\varphi_b, b \in B]$, $\mathbf{k}_1 := \mathbb{Q}[\mu, \varphi_b, b \in B]$. Then $\mathbf{k}_0 \subset \mathbf{k}_1$. Let $\Phi := \exp(\sum_{b \in B} \varphi_b b) \in \exp(\hat{\mathfrak{f}}_2^{\mathbf{k}_0}) \subset \exp(\hat{\mathfrak{f}}_2^{\mathbf{k}_1})$. For $b' \in B'$, $c \in C$, define $\mathrm{pent}_c, \mathrm{dual}_{b'} \in \mathbf{k}_0 \subset \mathbf{k}_1$ by $\sum_{c \in C} \mathrm{pent}_c\, c = \log(\mathrm{pent}(\Phi))$, $\sum_{b' \in B'} \mathrm{dual}_{b'}\, b' = \log(\mathrm{dual}(\Phi))$, $\sum_{b' \in B'} \mathrm{hex}_{b'}\, b' = \log(\mathrm{hex}(\mu, \Phi))$.

We then set $\mathbb{Q}[\underline{M}] := \mathbf{k}_1/(\mathrm{pent}_c, \mathrm{dual}_{b'}, \mathrm{hex}_{b'}, b' \in B', c \in C)$ and $\mathbb{Q}[M] := \mathbb{Q}[\underline{M}][\mu^{-1}] = \mathbb{Q}[\mu^{\pm 1}, \varphi_b, b \in B]/\{\text{ideal with the same generators}\}$. Then for any $\mathbb{Q}$-ring $\mathbf{k}$, we have (functorial in $\mathbf{k}$) bijections

$$\begin{tikzcd}
M(\mathbf{k}) \arrow[r, phantom, "\simeq"] \arrow[d, "\subset"'] & \mathrm{Hom}_{\mathbb{Q}\text{-rings}}(\mathbb{Q}[M], \mathbf{k}) \arrow[d, "\subset"] \\
\underline{M}(\mathbf{k}) \arrow[r, phantom, "\simeq"] & \mathrm{Hom}_{\mathbb{Q}\text{-rings}}(\mathbb{Q}[\underline{M}], \mathbf{k})
\end{tikzcd}$$

9.2.2. *The Transcendence Conjecture.* The KZ associator $(2\pi\,\mathrm{i}, \Phi_{KZ}) \in M(\mathbb{C})$ gives to a morphism $\varphi_{KZ} : \mathbb{Q}[M] \to \mathbb{C}$.

Conjecture 76. *(Transcendence Conjecture)* φ_{KZ} *is injective.*

We set $\mathbf{k}_{MZV} := \mathrm{im}(\mathbb{Q}[M] \overset{\varphi_{KZ}}{\to} \mathbb{C})$. This is a subring of $\mathbb{C}$ (according to [LM], this is the subring generated by $(2\pi\,\mathrm{i})^{\pm 1}$ and the MZVs (multizeta values)).

9.3. **Consequences of GC.**

Proposition 77. *The inclusion*

$$i_e(B_3) \subset \exp(\hat{\mathfrak{b}}_3^{+,\mathbb{C}}) \rtimes \mathrm{SL}_2(\mathbb{C}) \tag{50}$$

holds:

(a) for any $e \in Ell(\mathbb{C}) \times_{M(\mathbb{C})} \{\Phi_{KZ}\}$ *iff* $\mathfrak{b}_3 \triangleleft \mathfrak{r}_{ell}^{gr}$*;*

(b) for any $e \in \sigma(M(\mathbb{C}))$ *iff* $[\sigma(\mathfrak{grt}), \mathfrak{b}_3] \subset \hat{\mathfrak{b}}_3$*;*

(c) for any $e \in Ell(\mathbb{C})$ *iff* $\mathfrak{b}_3 \triangleleft \mathfrak{grt}_{ell}$*, i.e., iff the two above-mentioned conditions are realized. Moreover, GC implies that (50) holds for any* $e \in Ell(\mathbb{C})$*.*

We do not know whether the Lie algebraic statements in (a), (b), (c) hold, so they may be viewed as conjectures implied by GC.

Proof. Note first that for any $e \in Ell(\mathbb{C})$, and by Zariski density, (50) $\Leftrightarrow$ $(i_e(\langle B_3 \rangle(\mathbb{C})) = \exp(\hat{\mathfrak{b}}_3^{+,\mathbb{C}}) \rtimes \mathrm{SL}_2(\mathbb{C}))$.

We now prove (a). $e \in Ell(\mathbb{C}) \times_{M(\mathbb{C})} \{\Phi_{KZ}\}$ iff $e = \sigma(\Phi_{KZ}) * g$ for some $g \in R_{ell}^{gr}(\mathbb{C})$. So

$$\begin{aligned}
&((50) \text{ holds for any } e \in Ell(\mathbb{C}) \times_{M(\mathbb{C})} \{\Phi_{KZ}\}) \\
&\Leftrightarrow (g(i_{\sigma(\Phi_{KZ})}(\langle B_3 \rangle(\mathbb{C}))) = \exp(\hat{\mathfrak{b}}_3^{+,\mathbb{C}}) \rtimes \mathrm{SL}_2(\mathbb{C}) \text{ for any } g \in R_{ell}^{gr}(\mathbb{C})) \\
&\Leftrightarrow (g(\Gamma) = \Gamma \text{ for any } g \in R_{ell}^{gr}(\mathbb{C}), \text{ where } \Gamma = \exp(\hat{\mathfrak{b}}_3^{+,\mathbb{C}}) \rtimes \mathrm{SL}_2(\mathbb{C})) \\
&\Leftrightarrow (\mathfrak{b}_3 \triangleleft \mathfrak{r}_{ell}^{gr}).
\end{aligned}$$

Here the second equivalence follows from Proposition 71.

(b), (c) are then proved in the same way, using

$$(e \in \sigma(M(\mathbb{C}))) \Leftrightarrow (e = \sigma(\Phi_{KZ}) * \sigma(g) \text{ for some } g \in \mathrm{GRT}(\mathbb{C})),$$

$$(e \in Ell(\mathbb{C})) \Leftrightarrow (e = \sigma(\Phi_{KZ}) * g \text{ for some } g \in \mathrm{GRT}_{ell}(\mathbb{C})).$$

The equivalence (c) $\Leftrightarrow$ ((a) and (b)) follows from $\mathfrak{grt}_{ell} = \sigma(\mathfrak{grt}) \oplus \mathfrak{r}_{ell}^{gr}$. Finally, GC means that $\langle \mathfrak{sl}_2, \delta_{2k} \rangle = \mathfrak{r}_{ell}^{gr}$, which immediately implies (a), (b) and (c) as $\mathfrak{r}_{ell}^{gr} \triangleleft \mathfrak{grt}_{ell}$. $\square$

9.4. **Consequences of the Transcendence Conjecture (TC).**

Proposition 78. *If TC holds, then for any* $\mathbb{Q}$*-ring* $\mathbf{k}$ *and any* $\Phi \in M(\mathbf{k})$*,* $i_{\sigma(\Phi)}(B_3) \subset \exp(\hat{\mathfrak{b}}_3^{+,\mathbf{k}}) \rtimes \mathrm{SL}_2(\mathbf{k})$*.*

Proof. Recall that $\langle B_3 \rangle(-) \hookrightarrow R_{ell}(-)$, $\exp(\hat{\mathfrak{b}}_3^{+}) \rtimes \mathrm{SL}_2(-) \hookrightarrow R_{ell}^{gr}(-)$ are inclusions of $\mathbb{Q}$-group schemes, and $Ell \to M$, $M \overset{\sigma}{\to} M$ are morphisms of $\mathbb{Q}$-group schemes.

In the notation of Definition 46, any $x \in X(\mathbf{k})$ gives rise to a morphism $i_x : G(\mathbf{k}) \to H(\mathbf{k})$, defined by $g * x = x * i_x(g)$ for any $g \in G(\mathbf{k})$. The assignment $x \mapsto i_x$ is functorial in the following sense: if $\mathbf{k} \to \mathbf{k}'$ is a morphism of $\mathbb{Q}$-rings and $x' := \mathrm{im}(x \in X(\mathbf{k}) \to X(\mathbf{k}'))$, then

$$\begin{tikzcd}
G(\mathbf{k}) \arrow[r, "i_x"] \arrow[d] & H(\mathbf{k}) \arrow[d] \\
G(\mathbf{k}') \arrow[r, "i_{x'}"] & H(\mathbf{k}')
\end{tikzcd}$$

commutes.

For any $\mathbb{Q}$-scheme X and any $\mathbb{Q}$-ring $\mathbf{k}$, let $X \otimes \mathbf{k}$ be the $\mathbf{k}$-scheme $(X \otimes \mathbf{k})(\mathbf{k}') := X(\mathbf{k}')$ for any $\mathbf{k}' \in \{\mathbf{k}\text{-rings}\}$. Again with the notation of Definition 46, a torsor even gives rise to an assignment $X(\mathbf{k}) \ni x \mapsto (G \otimes \mathbf{k} \overset{i_x^{\mathbf{k}}}{\to} H \otimes \mathbf{k})$, where $i_x^{\mathbf{k}}$ is a morphism of $\mathbf{k}$-group schemes, defined by: $\forall \mathbf{k}' \in \{\mathbf{k}\text{-rings}\}$, $g * \bar{x} = \bar{x} * i_x^{\mathbf{k}}(g)$ for any $g \in (G \otimes \mathbf{k})(\mathbf{k}') = G(\mathbf{k}')$, where $\bar{x} := \mathrm{im}(x \in X(\mathbf{k}) \to X(\mathbf{k}'))$.

In particular, $\Phi_{KZ} \in M(\mathbf{k}_{MZV})$ gives rise to an isomorphism $i_{\sigma(\Phi_{KZ})} : R_{ell}(-) \otimes \mathbf{k}_{MZV} \overset{\sim}{\to} R_{ell}^{gr}(-) \otimes \mathbf{k}_{MZV}$, and therefore to a Lie algebra isomorphism $\mathrm{Lie}\, i_{\sigma(\Phi_{KZ})} : \mathfrak{r}_{ell} \otimes \mathbf{k}_{MZV} \overset{\sim}{\to} (\mathfrak{r}_{ell}^{gr} \otimes \mathbf{k}_{MZV})^{\wedge}$, whose $\otimes_{\mathbf{k}_{MZV}} \mathbb{C}$ is the infinitesimal of the isomorphism of Proposition 71.

The group scheme inclusions $\langle B_3 \rangle(-) \subset R_{ell}(-)$ and $\exp(\hat{\mathfrak{b}}_3^{+}) \rtimes \mathrm{SL}_2 \subset R_{ell}^{gr}(-)$ give rise to Lie algebra inclusions $\mathrm{Lie}\langle B_3 \rangle(-) \subset \mathfrak{r}_{ell}$ and $\hat{\mathfrak{b}}_3^{+} \subset \hat{\mathfrak{r}}_{ell}^{gr}$ and Proposition 71 implies that $\mathrm{Lie}\, i_{\sigma(\Phi_{KZ})} \otimes_{\mathbf{k}_{MZV}} \mathbb{C}$ restricts to an isomorphism $\mathrm{Lie}\langle B_3 \rangle(-) \otimes_{\mathbb{Q}} \mathbb{C} \to (\mathfrak{b}_3 \otimes_{\mathbb{Q}} \mathbb{C})^{\wedge}$. This implies that $\mathrm{Lie}\, i_{\sigma(\Phi_{KZ})}$ restricts to a Lie algebra isomorphism

$$\mathrm{Lie}\langle B_3 \rangle(-) \otimes_{\mathbb{Q}} \mathbf{k}_{MZV} \to (\mathfrak{b}_3 \otimes_{\mathbb{Q}} \mathbf{k}_{MZV})^{\wedge}.$$

We have Lie subalgebras $\mathbb{Q} \log \psi_{\pm} \otimes \mathbf{k}_{MZV}$ in the l.h.s., mapping to $(\mathbb{Q}e_{\pm} +$ terms of degree $> 0) \otimes \mathbf{k}_{MZV}$ in the r.h.s. (where $e_+ = e$, $e_- = f$, $\psi_+ = \psi$). This induces a diagram

$$\begin{tikzcd}
\langle B_3 \rangle(-) \otimes \mathbf{k}_{MZV} \arrow[r, "i_{\sigma(\Phi_{KZ})}"] & (\exp(\hat{\mathfrak{b}}_3^{+}) \rtimes \mathrm{SL}_2) \otimes \mathbf{k}_{MZV} \\
\mathbb{G}_a \otimes \mathbf{k}_{MZV} \arrow[u, "\subset"] \arrow[r, equal] & \mathbb{G}_a \otimes \mathbf{k}_{MZV} \arrow[u, "\subset"']
\end{tikzcd}$$

If now $\Phi \in M(\mathbf{k})$, the transcendence conjecture says that there exists a $\mathbb{Q}$-ring morphism $\mathbf{k}_{MZV} \overset{\varphi}{\to} \mathbf{k}$, such that $\Phi = \varphi_*(\Phi_{KZ})$. Applying this morphism to the above diagram, we get

$$\begin{tikzcd}
\langle B_3 \rangle(-) \otimes \mathbf{k} \arrow[r, "i_{\sigma(\Phi)}"] & (\exp(\hat{\mathfrak{b}}_3^{+}) \rtimes \mathrm{SL}_2) \otimes \mathbf{k} \\
\mathbb{G}_a \otimes \mathbf{k} \arrow[u, "\subset"] \arrow[r, equal] & \mathbb{G}_a \otimes \mathbf{k} \arrow[u, "\subset"']
\end{tikzcd}$$

Taking $\mathbf{k}$-points, we get a commutative diagram

$$\begin{tikzcd}
\langle B_3 \rangle(\mathbf{k}) \arrow[r, "i_{\sigma(\Phi)}"] & \exp(\hat{\mathfrak{b}}_3^{+,\mathbf{k}}) \rtimes \mathrm{SL}_2(\mathbf{k}) \\
\mathbf{k} \arrow[u, "\subset"] \arrow[r, equal] & \mathbf{k} \arrow[u, "\subset"']
\end{tikzcd}$$

The image of $1 \in \mathbf{k}$ is $\Psi_\pm \subset \langle B_3\rangle(\mathbf{k})$; then $i_{\sigma(\Phi)}(\Psi_\pm) \in \exp(\hat{\mathfrak{b}}_3^{+,\mathbf{k}}) \rtimes \mathrm{SL}_2(\mathbf{k}) \subset \exp(\hat{\mathfrak{b}}_3^{+,\mathbb{C}}) \rtimes \mathrm{SL}_2(\mathbb{C})$. □

10. Proofs of Proposition 5 and Theorems 6, 7 and 8

10.1. **Proof of Proposition 5.** As in [Na], let $\mathfrak{s}_0 : G_{\mathbb{Q}} \to \pi_1(M^{\mathbb{Q}}_{1,1^2}, \xi)$ be the section induced by the point ξ. The diagram $\pi_1(M^{\mathbb{Q}}_{1,1^2}, \xi) \overset{\curvearrowleft}{\to} G_{\mathbb{Q}}$ is described by the semidirect product decomposition $\pi_1(M^{\mathbb{Q}}_{1,1^2}, \xi) \simeq \widehat{B}_3 \rtimes G_{\mathbb{Q}}$, where the action $G_{\mathbb{Q}} \to \mathrm{Aut}(\widehat{B}_3)$ is $g * x := \mathfrak{s}_0(g) x \mathfrak{s}_0(g)^{-1}$. On the other hand, the diagram $\mathrm{GL}_2(\mathbb{Z}) \overset{\curvearrowleft}{\to} \widehat{\mathbb{Z}}^\times$ is described by the semidirect product decomposition $\mathrm{GL}_2(\widehat{\mathbb{Z}}) \simeq \mathrm{SL}_2(\mathbb{Z}) \rtimes \widehat{\mathbb{Z}}^\times$, where the action $\widehat{\mathbb{Z}}^\times \to \mathrm{Aut}(\mathrm{SL}_2(\widehat{\mathbb{Z}}))$ is $\lambda \bullet m := \left(\begin{smallmatrix} \lambda & 0 \\ 0 & 1 \end{smallmatrix}\right) m \left(\begin{smallmatrix} \lambda & 0 \\ 0 & 1 \end{smallmatrix}\right)^{-1}$.

Let σ_1, σ_2 be the Artin generators of B_3 (denoted $\bar{a}_1, \bar{a}_2$ in [Na]). As $\mathrm{SL}_2(\widehat{\mathbb{Z}})$ is profinite, there is a unique morphism

$$\widehat{B}_3 \to \mathrm{SL}_2(\widehat{\mathbb{Z}}), \quad x \mapsto \overline{x}$$

extending the quotient morphism $B_3 \to B_3/Z(B_3) \simeq \mathrm{SL}_2(\mathbb{Z})$, $\sigma_1 \mapsto \left(\begin{smallmatrix} 1 & 1 \\ 0 & 1 \end{smallmatrix}\right)$, $\sigma_2 \mapsto \left(\begin{smallmatrix} 1 & 0 \\ -1 & 1 \end{smallmatrix}\right)$. We will show that this morphism is compatible with the actions and the cyclotomic character $\chi : G_{\mathbb{Q}} \to \widehat{\mathbb{Z}}^\times$.

We first make the action of $G_{\mathbb{Q}}$ on $\widehat{B}_3$ explicit. Let us denote the map $G_{\mathbb{Q}} \to \widehat{\mathrm{GT}} \subset \widehat{\mathbb{Z}}^\times \times \widehat{F}_2$ by $g \mapsto (\chi(g), f_g)$. Using the formula $\beta_0(g) = \sigma_1^{8\rho_2(g)} \mathfrak{s}_0(g)$ in [Na] before Proposition 4.12, and Corollary 4.15 in the same paper, we obtain

$$g * \sigma_1 = \sigma_1^{\chi(g)}, \quad g * \sigma_2 = \mathrm{Ad}_{\sigma_1^{-8\rho_2(g)} f_g(\sigma_1^2, \sigma_2^2)^{-1}} (\sigma_2^{\chi(g)})$$

(here $\rho_2 : G_{\mathbb{Q}} \to \widehat{\mathbb{Z}}$ is the Kummer cocycle related to the roots of 2).

Then

$$\overline{g * \sigma_1} = \overline{\sigma_1^{\chi(g)}} = \begin{pmatrix} 1 & \chi(g) \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} \chi(g) & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \chi(g) & 0 \\ 0 & 1 \end{pmatrix}^{-1} = \chi(g) \bullet \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} = \chi(g) \bullet \overline{\sigma}_1;$$

on the other hand, Corollary 4.15 in [Na] says that

$$f_g\left(\begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ -2 & 1 \end{pmatrix}\right) = \pm \begin{pmatrix} 1 & 0 \\ -8\rho_2(g) & 1 \end{pmatrix} \begin{pmatrix} \chi(g)^{-1} & 0 \\ 0 & \chi(g) \end{pmatrix} \begin{pmatrix} 1 & -8\rho_2(g) \\ 0 & 1 \end{pmatrix}$$

(identity in $\mathrm{SL}_2(\widehat{\mathbb{Z}})$), therefore

$$\begin{aligned} \overline{g * \sigma_2} &= \overline{\mathrm{Ad}_{\sigma_1^{-8\rho_2(g)} f_g(\sigma_1^2, \sigma_2^2)^{-1}} \sigma_2^{\chi(g)}} = \mathrm{Ad}_{\pm \left(\begin{smallmatrix} \chi(g) & 0 \\ 0 & \chi(g)^{-1} \end{smallmatrix}\right) \left(\begin{smallmatrix} 1 & 0 \\ 8\rho_2(g) & 1 \end{smallmatrix}\right)} \begin{pmatrix} 1 & 0 \\ -\chi(g) & 1 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 \\ -\chi(g)^{-1} & 1 \end{pmatrix} = \begin{pmatrix} \chi(g) & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} \chi(g) & 0 \\ 0 & 1 \end{pmatrix}^{-1} = \chi(g) \bullet \overline{\sigma}_2. \end{aligned}$$

It follows that $\widehat{B}_3 \to \mathrm{SL}_2(\widehat{\mathbb{Z}})$ intertwines the actions of $G_{\mathbb{Q}}$ and $\widehat{\mathbb{Z}}^\times$, as claimed. □

10.2. **Proof of Theorem 6.** We first prove:

Proposition 79. *There is a morphism $\pi_1(M^{\mathbb{Q}}_{1,1^2}, \xi) \to \widehat{\mathrm{GT}}_{ell}$, compatible with the semidirect product structures; namely, the diagram*

$$\begin{tikzcd} \pi_1(M^{\mathbb{Q}}_{1,1^2}, \xi) \arrow[r, "\overset{\mathfrak{s}_0}{\curvearrowleft}"] \arrow[d] & G_{\mathbb{Q}} \arrow[d] \\ \widehat{\mathrm{GT}}_{ell} \arrow[r, "\overset{\sigma}{\curvearrowleft}"] & \widehat{\mathrm{GT}} \end{tikzcd}$$

commutes.

Proof. We first set $\widehat{R}_{ell} := \mathrm{Ker}(\widehat{\mathrm{GT}_{ell}} \to \widehat{\mathrm{GT}})$. Recall that $\widehat{R}_{ell}$ is a subgroup of $\mathrm{Aut}(\widehat{F}_2)$. $\mathrm{Aut}(\widehat{F}_2)$ is profinite ([DDMS], Thm. 5.3), and the map $\mathrm{Aut}(\widehat{F}_2) \to \widehat{F}_2^2$, $\theta \mapsto (\theta(X), \theta(Y))$ is continuous (loc. cit., Ex. 2, p. 96). As $\widehat{R}_{ell}$ is the preimage of 1 by a continuous map $(\widehat{F}_2)^2 \to (\widehat{B}_{1,3})^2 \times \widehat{F}_2$, it is closed, so $\widehat{R}_{ell}$ is a closed subgroup of $\mathrm{Aut}(\widehat{F}_2)$, hence is profinite.

As $\mathrm{im}(B_3 = \widetilde{\mathrm{SL}_2 \mathbb{Z}} \to \mathrm{Aut}(\widehat{F}_2)) \subset \widehat{R}_{ell}$, we obtain a morphism $B_3 \to \widehat{R}_{ell}$, which extends to a morphism

$$\widehat{B}_3 = \pi_1(M_{1,1^2}^{\mathbb{Q}} \otimes \bar{\mathbb{Q}}, \xi) \to \widehat{R}_{ell}$$

as $\widehat{R}_{ell}$ is profinite.

We now prove that the morphism $\widehat{B}_3 \to \widehat{R}_{ell}$ is compatible with the actions of $G_{\mathbb{Q}}$ and $\widehat{\mathrm{GT}}$ on both sides via $\mathfrak{s}_0$ and σ and the morphism $G_{\mathbb{Q}} \to \widehat{\mathrm{GT}}$, i.e., the commutativity of

$$\begin{tikzcd}
G_{\mathbb{Q}} \times \widehat{B}_3 \arrow[r] \arrow[d] & \widehat{B}_3 \arrow[d] \\
\widehat{\mathrm{GT}} \times \widehat{R}_{ell} \arrow[r] & \widehat{R}_{ell}
\end{tikzcd} \tag{51}$$

According to [Na], Section 4, the upper action is compatible with the actions of α, β of $G_{\mathbb{Q}}$, $\widehat{B}_3 = \pi_1(M_{1,1^2}^{\mathbb{Q}}, \xi)$ by automorphisms of $\widehat{F}_2 = \pi_1(C_\xi \otimes \bar{\mathbb{Q}}, \tilde{\xi})$, where C_ξ is the fiber over ξ of $M_{1,2}^{\mathbb{Q}} \to M_{1,1}^{\mathbb{Q}}$ and $\tilde{\xi}$ is a tangential base point close to the marked point of C_ξ; namely, $\alpha(g * \gamma) = \beta(g)\alpha(\gamma)\beta(g)^{-1}$ for any $g \in G_{\mathbb{Q}}$, $\gamma \in \pi_1(M_{1,1^2} \otimes \bar{\mathbb{Q}}, \xi)$, where α, β are the morphisms to $\mathrm{Aut}(\pi_1(C_\xi, \tilde{\xi}))$.

The commutativity of the diagram (51) then follows from that of

$$\begin{tikzcd}
\pi_1(M_{1,1^2}^{\mathbb{Q}} \otimes \bar{\mathbb{Q}}, \xi) \arrow[r] \arrow[d] & \mathrm{Aut}\,\pi_1(C_\xi, \tilde{\xi}) \arrow[d] \\
\widehat{R}_{ell} \arrow[r] & \mathrm{Aut}(\widehat{F}_2)
\end{tikzcd}$$

which we have showed above, and of

$$\begin{tikzcd}
G_{\mathbb{Q}} \arrow[r] \arrow[d] & \mathrm{Aut}\,\pi_1(C_\xi, \tilde{\xi}) \arrow[d] \\
\widehat{\mathrm{GT}_{ell}} \arrow[r] & \mathrm{Aut}(\widehat{F}_2)
\end{tikzcd}$$

which we now show.

According to [Na], Corollary 4.5, the action of $G_{\mathbb{Q}}$ on $\pi_1(C_\xi \otimes \bar{\mathbb{Q}}, \tilde{\xi})$ may be described as follows. $\pi_1(C_\xi \otimes \bar{\mathbb{Q}}, \tilde{\xi})$ is topologically free, generated by x_1, x_2. The action of $\gamma \in G_{\mathbb{Q}}$ on this group is

$$\gamma^*(x_1) = f_\gamma(x_1, z_1) x_1^{\chi(\gamma)} f_\gamma(x_1, z_1)^{-1}, \tag{52}$$

$$\gamma^*(x_2) = f_\gamma(x_1, z_1) x_1^{1-\chi(\gamma)} f_\gamma^{\overrightarrow{\infty 1}}(x_1, x_1^{-1} z_1 x_1)^{-1} x_2 x_1^{\chi(\gamma)-1} f_\gamma(x_1, z_1)^{-1}, \tag{53}$$

where $z_1 = (x_2, x_1) = x_2 x_1 x_2^{-1} x_1^{-1}$, $\gamma \mapsto (\chi(\gamma), f_\gamma)$ is the map $G_{\mathbb{Q}} \to \widehat{\mathrm{GT}}$, and $f_\gamma^{\overrightarrow{\infty 1}}(a, b) = f_\gamma(b, c) b^{(\chi(\gamma)-1)/2} f_\gamma(a, b)$ for $abc = 1$.

Under the identification $x_1 \mapsto X$, $x_2 \mapsto Y$, formula (52) corresponds to the expression of g_+ in Proposition 35. We now prove that (53) corresponds to g_- in the same Proposition. For this we must show that

$$\begin{aligned}
f(X, (Y, X)) X^{1-\lambda} f^{\overrightarrow{\infty 1}}(X, (X^{-1}, Y))^{-1} Y X^{\lambda-1} f(X, (Y, X))^{-1} \\
= Z^{(\lambda-1)/2} f(X^{-1} Z^{-1}, Z) Y f(X, (Y, X))^{-1}
\end{aligned}$$

for $(\lambda, f) \in \widehat{\mathrm{GT}}$ and $Z = (Y, X)$; this is a straightforward consequence of the octagon identity

$$f(X^{-1} Z^{-1}, Z)(ZX)^{-\lambda} f(Z, X^{-1} Z^{-1}) Z^{(\lambda+1)/2} f(X, Z) X^{\lambda} f(Z, X) Z^{(\lambda-1)/2} = 1,$$

which follows from the hexagon and duality identities. □

We now describe the groupoid $T_{ell,n}$. Its set of objects is $\mathrm{Ob}(T_{ell,n}) = \{(t,\varphi) | t$ is a 3-valent rooted planar tree with n leaves, $\varphi : L_t \to [n]$ is a bijection from the set of leaves of t to $[n] = \{1, \dots, n\}\}$. Note that the planar structure on T induces a total order on L_t. We then set $T_{ell,n}((t,\varphi),(t',\varphi')) := B_{1,n} \times_{S_n} \{\varphi_{t'} i \varphi_t^{-1}\}$, where $i : L_t \to L_{t'}$ is the only order-preserving bijection. We define $\widehat{T}_{ell,n}$, $T^l_{ell,n}$, $T_{ell,n}(-)$, as the profinite, pro-l, prounipotent completions of $T_{ell,n}$. The objects of $T_{ell,n}$ correspond to maximally degenerate elliptic curves. For example, the following diagram



corresponds to the limit in $\overline{M}^{\mathbb{Q}}_{1,n}$ of $\mathbb{G}_m/q^{\mathbb{Z}}$ with marked points $z_1 = \varepsilon^2$, $z_2 = 0$, $z_3 = \varepsilon$, $z_4 = \varepsilon + \varepsilon^2$, $z_5 = \varepsilon + \varepsilon^2 + \varepsilon^3$ as $\varepsilon/q, q \to 0$.

We will construct the action of $\widehat{\mathrm{GT}}_{ell}$ on $\widehat{T}_{ell,n}$ in several steps. If $\mathcal{C}$ is a category, we denote by $\mathrm{Aut}(\mathcal{C})$ its group of automorphisms (as a category, even if $\mathcal{C}$ has a braided monoidal structure).

For $(\lambda, f) \in \underline{\mathrm{GT}}$, we define an endofunctor $i_{\lambda,f}$ of $\widehat{\mathbf{PaB}}$ to be the composite functor $\widehat{\mathbf{PaB}} \overset{\alpha_{(\lambda,f)}}{\to} (\lambda, f) * \widehat{\mathbf{PaB}} \overset{\sim}{\to} \widehat{\mathbf{PaB}}$, where the first functor is the unique tensor functor which induces the identity on objects, and the second functor is the identity functor (which is not tensor).

Proposition 80. $(\lambda, f) \mapsto i^{-1}_{\lambda,f}$ *is a morphism* $\widehat{\mathrm{GT}} \to \mathrm{Aut}(\widehat{\mathbf{PaB}})$.

Proof. The identity $i_{(\lambda',f')} i_{(\lambda,f)} = i_{(\lambda,f)(\lambda',f')}$ follows from the commutativity of the diagram

$$\begin{tikzcd}
 & & \mathcal{C} \arrow[dr] \arrow[ddrr, bend left, "i_{(\lambda',f')}"] & & \\
 & (\lambda,f)*\mathcal{C} \arrow[ur, "\sim"] \arrow[dr] & & (\lambda',f')*\mathcal{C} \arrow[dr, "\sim"] & \\
\mathcal{C} \arrow[ur] \arrow[uurr, bend left, "i_{(\lambda,f)}"] \arrow[rr] \arrow[rrrr, bend right, "i_{(\lambda,f)(\lambda',f')}"'] & & (\lambda,f)(\lambda',f')*\mathcal{C} \arrow[ur, "\sim"] \arrow[rr, "\sim"] & & \mathcal{C}
\end{tikzcd}$$

in which the commutativity of the central square follows from that of

$$\begin{tikzcd}
(\lambda,f)*\mathcal{C} \arrow[r, "(\lambda,f)*\varphi"] \arrow[d, "\sim"'] & (\lambda,f)*\mathcal{D} \arrow[d, "\sim"] \\
\mathcal{C} \arrow[r, "\varphi"] & \mathcal{D}
\end{tikzcd}$$

for any braided monoidal categories $\mathcal{C}, \mathcal{D}$ and any tensor functor $\varphi : \mathcal{C} \to \mathcal{D}$. □

One constructs in the same way a morphism $\widehat{\mathrm{GT}}_{ell} \to \mathrm{Aut}(\widehat{\mathbf{PaB}}_{ell})$.

We now generalize these morphisms in the context of colored categories. If $\mathcal{C}_0$ is a braided monoidal category, then $\mathrm{Ob}\,\mathcal{C}_0$ is a magma (i.e., a set equipped with a composition map and a unit). Let $\phi : M \to \mathrm{Ob}\,\mathcal{C}_0$ be a magma morphism, then we construct a braided monoidal category $\phi*\mathcal{C}_0$ by $\mathrm{Ob}\,\phi^*\mathcal{C}_0 = M$, $\phi^*\mathcal{C}_0(m,m') := \mathcal{C}_0(\phi(m),\phi(m'))$ and by the condition that the obvious functor $\phi^*\mathcal{C}_0 \to \mathcal{C}_0$ is tensor. If $\mathcal{C}_0 \to \mathcal{C}$ is an elliptic structure over $\mathcal{C}_0$, then one defines an elliptic structure $\phi^*\mathcal{C}_0 \to \phi^*\mathcal{C}$ over $\phi^*\mathcal{C}_0$ in the same way. We then have group morphisms $\mathrm{Aut}\,\mathcal{C}_0 \to \mathrm{Aut}\,\phi^*\mathcal{C}_0$, $\mathrm{Aut}\,\mathcal{C} \to \mathrm{Aut}\,\phi^*\mathcal{C}$.

If S is a set, we denote by $mag(S)$ the free magma generated by S. The constant map $S \to \{\bullet\}$ induces a magma morphism $\phi : mag(S) \to mag(\{\bullet\}) \simeq \mathrm{Ob}\,\widehat{\mathbf{PaB}}$. We then set $\widehat{\mathbf{PaB}}_S := \phi^*\widehat{\mathbf{PaB}}$, $\widehat{\mathbf{PaB}}_{ell,S} := \phi^*\widehat{\mathbf{PaB}}_{ell}$.

We then have composite morphisms $\widehat{\mathrm{GT}} \to \mathrm{Aut}(\widehat{\mathbf{PaB}}) \to \mathrm{Aut}(\widehat{\mathbf{PaB}}_S)$ and similarly $\widehat{\mathrm{GT}}_{ell} \to \mathrm{Aut}(\widehat{\mathbf{PaB}}_{ell,S})$.

Recall now that $\widehat{T}_{ell,n}$ may be viewed as the full subcategory of $\widehat{\mathbf{PaB}}_{ell,[n]}$, where the objects are the preimages of $1\dot{+}\cdots\dot{+}n$ under the map $mag([n]) \xrightarrow{\psi} \mathbb{N}[n]$, extending the identity on $[n]$, where $\mathbb{N}[n]$ is the free abelian semigroup generated by $[n]$ (in which the addition is denoted $\dot{+}$). In other words, $\widehat{T}_{ell,n} = i^*\widehat{\mathbf{PaB}}_{ell,[n]}$, where $i : \psi^{-1}(1\dot{+}\cdots\dot{+}n) \to mag([n]) = \mathrm{Ob}\,\widehat{\mathbf{PaB}}_{ell,[n]}$ is the canonical inclusion. We then have a composite morphism $\widehat{\mathrm{GT}}_{ell} \to \mathrm{Aut}(\widehat{\mathbf{PaB}}_{ell,[n]}) \to \mathrm{Aut}(\widehat{T}_{ell,n})$.

We now show that the composite morphism $\pi_1(M^{\mathbb{Q}}_{1,1^2},\xi) \to \widehat{\mathrm{GT}}_{ell} \to \mathrm{Aut}(\widehat{T}_{ell})$ coincides with the action induced by the fibration $M^{\mathbb{Q}}_{1;1^2,n-1} \to M^{\mathbb{Q}}_{1,1^2}$. We first prove it for the subgroup $\widehat{B}_3 \subset \pi_1(M^{\mathbb{Q}}_{1,1^2},\xi)$.

Define $\widehat{\mathbf{B}}_{ell}$ as the category with $\mathrm{Ob}\,\widehat{\mathbf{B}}_{ell} = \mathbb{N}$, $\widehat{\mathbf{B}}_{ell}(n,m) = \emptyset$ if $m \neq n$, and $\widehat{\mathbf{B}}_{ell}(n,n) = \widehat{B}_{1,n}$. We have a natural functor $\widehat{\mathbf{PaB}}_{ell} \to \widehat{\mathbf{B}}_{ell}$, defined as the length map $l : mag(\{\bullet\}) \to \mathbb{N}$ at the level of objects and as the identity at the level of morphisms; we have actually $\widehat{\mathbf{PaB}}_{ell} \simeq l^*\widehat{\mathbf{B}}_{ell}$. As $\widehat{R}_{ell} \subset \widehat{\mathrm{GT}}_{ell}$ acts trivially on the images of the associativity constraints under $\widehat{\mathbf{PaB}} \to \widehat{\mathbf{PaB}}_{ell}$, its action on $\widehat{\mathbf{PaB}}_{ell}$ is the lift of an action of $\widehat{R}_{ell}$ on $\widehat{\mathbf{B}}_{ell}$. One checks explicitly that the composition of this action with the morphism $\widehat{B}_3 \to \widehat{R}_{ell}$ coincides with the geometric action of $\widehat{B}_3$ on the various groups $\widehat{B}_{1,n}$.

We now show that the action $G_{\mathbb{Q}} \xrightarrow{\sigma} \pi_1(M_{1,1^2},\xi) \to \mathrm{Aut}(\widehat{T}_{ell,n})$ coincides with the action induced by $M^{\mathbb{Q}}_{1;1^2,n-1} \overset{\overset{\sigma_T}{\curvearrowleft}}{\to} M^{\mathbb{Q}}_{1,1^2} \overset{\overset{\xi}{\curvearrowleft}}{\to} \mathrm{Spec}\,\mathbb{Q}$. For $n = 1, 2$, this is proved in [Na]. In [Mt], the action of $G_{\mathbb{Q}}$ for general n was computed from the action for $n = 1$ and from the action on braid groups in genus 0; this action coincides with $G_{\mathbb{Q}} \to \mathrm{Aut}(\widehat{T}_{ell,n})$.

10.3. **Proof of Theorem 7.** We prove 1). The morphism $\mathrm{GT}^l_{ell} \to \mathrm{Aut}(\mathbf{PaB}^l_{ell})$ factorizes as $\mathrm{GT}^l_{ell} \to \mathrm{Aut}(\mathbf{PaB}_{ell}(\mathbb{Q}_l),\mathbf{PaB}^l_{ell}) \to \mathrm{Aut}(\mathbf{PaB}^l_{ell})$; the colored morphism $\mathrm{GT}^l_{ell} \to \mathrm{Aut}(\mathbf{PaB}^l_{ell,[n]})$ factorizes in the same way. We then have a commutative diagram

$$\begin{tikzcd}
\pi_1(M^{\mathbb{Q}}_{1,1^2},\xi) \arrow[r] & \mathrm{GT}^l_{ell} \arrow[r] \arrow[dr] & \mathrm{Aut}(\mathbf{PaB}_{ell,[n]}(\mathbb{Q}_l),\mathbf{PaB}^l_{ell,[n]}) \arrow[r] \arrow[d] & \mathrm{Aut}(T_{ell,n}(\mathbb{Q}_l),T^l_{ell,n}) \arrow[d] \\
 & & \mathrm{Aut}(\mathbf{PaB}^l_{ell,[n]}) \arrow[r] & \mathrm{Aut}(T^l_{ell,n})
\end{tikzcd}$$

2a) has been proved in Section 2.4 (The semigroup scheme $\underline{\mathrm{GT}}_{ell}(-)$).

2b) Define a morphism $\pi_1(M^{\mathbb{Q}}_{1,1^2},\xi) \to \mathrm{GT}_{ell}(\mathbb{Q}_l)$ as the composite morphism $\pi_1(M^{\mathbb{Q}}_{1,1^2},\xi) \to \widehat{\mathrm{GT}}_{ell} \to \mathrm{GT}^l_{ell} \to \mathrm{GT}_{ell}(\mathbb{Q}_l)$. It follows from Proposition 5 that

$$\begin{tikzcd} \pi_1(M^{\mathbb{Q}}_{1,1^2},\xi) \arrow[r] \arrow[d] & G_{\mathbb{Q}} \arrow[d] \\ \mathrm{GL}_2(\mathbb{Z}_l) \arrow[r] & \mathbb{Z}_l^\times \end{tikzcd}$$

commutes, and from Proposition 79 that

$$\begin{tikzcd} \pi_1(M^{\mathbb{Q}}_{1,1^2},\xi) \arrow[r] \arrow[d] & G_{\mathbb{Q}} \arrow[d] \\ \mathrm{GT}_{ell}(\mathbb{Q}_l) \arrow[r] & \mathrm{GT}(\mathbb{Q}_l) \end{tikzcd}$$

commutes. It remains to check that

$$\begin{tikzcd} \pi_1(M^{\mathbb{Q}}_{1,1^2},\xi) \arrow[r] \arrow[dr] & \mathrm{GT}_{ell}(\mathbb{Q}_l) \\ & \mathrm{GL}_2(\mathbb{Q}_l) \arrow[u] \end{tikzcd}$$

commutes; for this, it suffices to check that

$$\begin{tikzcd} \widehat{B}_3 \arrow[r] \arrow[dr] & R_{ell}(\mathbb{Q}_l) \\ & \mathrm{SL}_2(\mathbb{Q}_l) \arrow[u] \end{tikzcd}$$

commutes and to use that

$$\begin{tikzcd} G_{\mathbb{Q}} \arrow[r] \arrow[dr] & \mathrm{GT}(\mathbb{Q}_l) \\ & \mathbb{Q}_l^\times \arrow[u] \end{tikzcd}$$

commutes.

2c) follows from the proof of 1) and from the diagram

$$\begin{tikzcd} \mathrm{GT}^l_{ell} \arrow[r] \arrow[d] & \mathrm{Aut}(\mathbf{PaB}_{ell,[n]}(\mathbb{Q}_l), \mathbf{PaB}^l_{ell,[n]}) \arrow[r] \arrow[d] & \mathrm{Aut}(T_{ell,n}(\mathbb{Q}_l), T^l_{ell,n}) \arrow[d] \\ \mathrm{GT}_{ell}(\mathbb{Q}_l) \arrow[r] & \mathrm{Aut}(\mathbf{PaB}_{ell,[n]})(\mathbb{Q}_l) \arrow[r] & \mathrm{Aut}(T_{ell,n})(\mathbb{Q}_l) \end{tikzcd}$$

3) Recall that $T_{ell,n}(\mathbf{k})$ is defined from $\widehat{T}_{ell,n}$ by replacing $\widehat{B}_n$ by $B_{1,n}(\mathbf{k}) = P_{1,n}(\mathbf{k}) *_{P_{1,n}} B_{1,n}$. We then define $T^{gr}_{ell,n}(\mathbf{k})$ in the same way, replacing $B_{1,n}(\mathbf{k})$ by $\exp(\hat{\mathfrak{t}}^{\mathbf{k}}_{1,n}) \rtimes S_n$. Define the colored version $\mathbf{PaCD}_{ell,[n]}(\mathbf{k})$ of $\mathbf{PaCD}_{ell}(\mathbf{k})$ similarly to $\widehat{\mathbf{PaB}}_{ell,[n]}$; then $T^{gr}_{ell,n}(\mathbf{k})$ is a full subcategory of $\mathbf{PaCD}_{ell,[n]}(\mathbf{k})$. Any elliptic associator defined over $\mathbf{k}$ gives rise to an isomorphism $\mathbf{PaB}_{ell,[n]}(\mathbf{k}) \to \mathbf{PaCD}_{ell,[n]}(\mathbf{k})$ and to a compatible isomorphism $\mathrm{GT}_{ell}(\mathbf{k}) \to \mathrm{GRT}_{ell}(\mathbf{k})$. The former isomorphism restricts to an isomorphism $T_{ell,n}(\mathbf{k}) \to T^{gr}_{ell,n}(\mathbf{k})$ and we have a commutative diagram

$$\begin{tikzcd} \mathrm{GT}_{ell}(\mathbf{k}) \arrow[r] \arrow[d, "\sim"'] & \mathrm{Aut}(\mathbf{PaB}_{ell,[n]}(\mathbf{k})) \arrow[r] \arrow[d, "\sim"'] & \mathrm{Aut}(T_{ell,n}(\mathbf{k})) \arrow[d, "\sim"'] \\ \mathrm{GRT}_{ell}(\mathbf{k}) \arrow[r] & \mathrm{Aut}(\mathbf{PaCD}_{ell,[n]}(\mathbf{k})) \arrow[r] & \mathrm{Aut}(T^{gr}_{ell,n}(\mathbf{k})) \end{tikzcd}$$

10.4. **Proof of Theorem 8.** Let $\mathbf{k}$ be a $\mathbb{Q}$-ring. For $(c,g) \in \mathrm{GRT}(\mathbf{k})$, define $i_{(c,g)} \in \mathrm{End}(\mathbf{PaCD})$ as the composite functor $\mathbf{PaCD} \overset{\beta_{(c,g)}}{\to} (c,g) * \mathbf{PaCD} \overset{\sim}{\to} \mathbf{PaCD}$, where $\beta_{(c,g)}$ is the only IBMC functor inducing the identity on objects, and the second functor is the identity at the level of objects and morphisms (it is not compatible with the IBMC structures). As in Proposition 80, one proves that $(c,g) \mapsto i^{-1}_{(c,g)}$ induces a group morphism $\mathrm{GRT}(\mathbf{k}) \to \mathrm{Aut}(\mathbf{PaCD})$. One similarly constructs morphisms $\mathrm{GRT}_{ell}(\mathbf{k}) \to \mathrm{Aut}(\mathbf{PaCD}_{ell,[n]})$.

Proposition 81. *Let* $(\mu,\Phi) \in M_1(\mathbf{k})$. *Let* $j_{(\mu,\Phi)} : \mathrm{GT}(\mathbf{k}) \to \mathrm{GRT}(\mathbf{k})$ *be the isomorphism defined by* (μ,Φ) *(see Section 8), and* $\tilde{j}_{(\mu,\Phi)} : \mathbf{PaB}(\mathbf{k}) \to \mathbf{PaCD}$ *be the isofunctor obtained as the unique braided monoidal functor* $\mathbf{PaB}(\mathbf{k}) \to (\mu,\Phi) * \mathbf{PaCD}$ *which is the identity on objects. Then the diagram*

$$\begin{tikzcd} \mathrm{GT}(\mathbf{k}) \arrow[r] \arrow[d, "j_{(\mu,\Phi)}"'] & \mathrm{Aut}(\mathbf{PaB}(\mathbf{k})) \arrow[d, "\tilde{j}_{(\mu,\Phi)}"] \\ \mathrm{GRT}(\mathbf{k}) \arrow[r] & \mathrm{Aut}(\mathbf{PaCD}) \end{tikzcd}$$

commutes. In the same way, for any $e \in Ell(\mathbf{k})$, *the diagram*

$$\begin{tikzcd}
\mathrm{GT}_{ell}(\mathbf{k}) \arrow[r] \arrow[d, "i_e"'] & \mathrm{Aut}(\mathbf{PaB}_{ell,[n]}(\mathbf{k})) \arrow[d, "\tilde{i}_e"] \\
\mathrm{GRT}_{ell}(\mathbf{k}) \arrow[r] & \mathrm{Aut}(\mathbf{PaCD}_{ell,[n]})
\end{tikzcd} \tag{54}$$

commutes.

Proof. Let $(c,f) \in \mathrm{GT}(\mathbf{k})$, $(c,g) := j_{(\mu,\Phi)}(c,f)$ so $(c,f) * (\mu,\Phi) = (\mu,\Phi) * (c,g)$. We will prove that

$$\tilde{j}_{(\mu,\Phi)} \circ i_{(c,f)} = i_{(c,g)} \circ \tilde{j}_{(\mu,\Phi)}. \tag{55}$$

Denote by $\mathrm{can}_{(c,f)} : (c,f) * \mathbf{PaB}(\mathbf{k}) \xrightarrow{\sim} \mathbf{PaB}(\mathbf{k})$, $\mathrm{can}_{(c,g)} : (c,g) * \mathbf{PaCD} \xrightarrow{\sim} \mathbf{PaCD}$ the identity functors. In the diagram

$$\begin{tikzcd}
\mathbf{PaB}(\mathbf{k}) \arrow[r, "\tilde{j}_{(\mu,\Phi)}"] \arrow[d, "\alpha_{(c,f)}"'] & (\mu,\Phi) * \mathbf{PaCD} \arrow[dr, "(\mu,\Phi)*\beta_{(c,g)}"] & \\
(c,f) * \mathbf{PaB}(\mathbf{k}) \arrow[r, "(c,f)*\tilde{j}_{(\mu,\Phi)}"] \arrow[d, "\mathrm{can}_{(c,f)}"'] & (c,f) * (\mu,\Phi) * \mathbf{PaB}(\mathbf{k}) \arrow[r, "\sim"] & (\mu,\Phi) * (c,g) * \mathbf{PaB}(\mathbf{k}) \arrow[dl, "(\mu,\Phi)*\mathrm{can}_{(c,g)}"] \\
\mathbf{PaB}(\mathbf{k}) \arrow[r, "\tilde{j}_{(\mu,\Phi)}"'] & (\mu,\Phi) * \mathbf{PaCD} &
\end{tikzcd}$$

the upper square commutes as it is composed of tensor functors which induce the identity on objects, while the lower square commutes because the horizontal functors coincide and the vertical functors are the identity. This implies (55). ☐

Assume now that $(\mu,\Phi) = (2\pi\,\mathrm{i}, \Phi_{KZ}) \in M(\mathbb{C})$. Using (54) and Proposition 71, we get a commutative diagram

$$\begin{tikzcd}
\langle B_3\rangle(\mathbb{C}) \arrow[r, hook] \arrow[d] & R_{ell}(\mathbb{C}) \arrow[r, hook] \arrow[d] & \mathrm{GT}_{ell}(\mathbb{C}) \arrow[r] \arrow[d] & \mathrm{Aut}\,\mathbf{PaB}_{ell,[n]}(\mathbf{k}) \arrow[r] \arrow[d] & \mathrm{Aut}\,\mathbf{PaB}_{ell,[n]}(\mathbf{k})(O) \arrow[d] \\
\begin{array}{c}\exp\langle\mathfrak{sl}_2,\delta_{2k}\rangle_+^{\mathbb{C}} \\ \rtimes \mathrm{SL}_2(\mathbb{C})\end{array} \arrow[r, hook] & R_{ell}^{gr}(\mathbb{C}) \arrow[r, hook] & \mathrm{GRT}_{ell}(\mathbb{C}) \arrow[r] & \mathrm{Aut}\,\mathbf{PaCD}_{ell,[n]} \arrow[r] & \mathrm{Aut}\,\mathbf{PaCD}_{ell,[n]}(O)
\end{tikzcd}$$

for any object O. The infinitesimal of the left vertical arrow is a filtered isomorphism $\mathrm{Lie}\langle B_3\rangle^{\mathbb{C}} \simeq \hat{\mathfrak{b}}_3^{\mathbb{C}}$, whose associated graded isomorphism is the canonical identification. Taking for O a parenthesization of the word $1 \cdots n$, the isomorphism $i_e : \mathbf{PaB}_{ell,[n]}(\mathbf{k})(O) \to \mathbf{PaCD}_{ell,[n]}(O)$ yields an isomorphism $\mathrm{Lie}\,P_{1,n}^{\mathbb{C}} \simeq \hat{\mathfrak{t}}_{1,n}^{\mathbb{C}}$, whose associated graded is the canonical identification. The infinitesimal of the total diagram says that these isomorphisms are compatible with actions, i.e.,

$$\begin{tikzcd}
\mathrm{Lie}\langle B_3\rangle^{\mathbb{C}} \arrow[r] \arrow[d] & \mathrm{Der}\,\mathrm{Lie}\,P_{1,n}^{\mathbb{C}} \arrow[d] \\
\hat{\mathfrak{b}}_3^{\mathbb{C}} \arrow[r] & \mathrm{Der}(\hat{\mathfrak{t}}_{1,n}^{\mathbb{C}})
\end{tikzcd}$$

commutes. The fact that there exist isomorphisms over $\mathbb{Q}$ with the same properties follows from the following Lemma 84, which we prove after some preparation.

The following lemma was communicated to us by P. Etingof; it is inspired by the results of [Dr].

Proposition 82. *Let* $U = \lim_{\leftarrow} U_i$ *be a prounipotent* $\mathbb{Q}$*-group scheme (where* $U_0 = 1$*) and let* $T := \lim_{\leftarrow} T_i$, *where* T_i *are a compatible system of torsors under* U_i, *defined over* $\mathbb{Q}$. *If* $T(\mathbb{C}) \neq \emptyset$, *then* $T(\mathbb{Q}) \neq \emptyset$.

Proof. Let $\tilde U_i := \mathrm{im}(U \to U_i)$, then $U = \lim_{\leftarrow} \tilde U_i$, where $\cdots \to \tilde U_2 \to \tilde U_1 \to \tilde U_0 = 1$ is a sequence of epimorphisms of unipotent groups. We set $K_i := \mathrm{Ker}(U \to \tilde U_i)$; then $K_i \lhd U$. If we set $\tilde T_i := \mathrm{im}(T \to T_i)$, then $\tilde T_i \simeq T/K_i$ is a torsor over $\tilde U_i$; T is the inverse limit of $\cdots \to \tilde T_2 \to \tilde T_1$, where the morphisms are onto.

We may therefore assume w.l.o.g. that the morphisms $U_{i+1} \to U_i$, $T_{i+1} \to T_i$ are onto; if $K_i := \mathrm{Ker}(U \to U_i)$, then $T_i = T/K_i$.

We now show that the projective systems $\cdots \to T_2 \to T_1$, $\cdots \to U_2 \to U_1$ may be completed so that for any i, $\mathrm{Ker}(U_{i+1} \to U_i) \simeq \mathbb{G}_a$. Indeed, for $U_{i+1} \to U' \to U_i$ a sequence of epimorphisms, we set $K' := \mathrm{Ker}(U \to U')$ and $T' := T/K'$. Then $T_{i+1} \to T' \to T_i$ is a sequence of epimorphisms, compatible with $U_{i+1} \to U' \to U_i$.

Let $t \in T(\mathbb{C})$. We construct a sequence $(k_i)_{i\geq 0}$, where $k_i \in K_i(\mathbb{C})$, such that $\mathrm{im}(k_i \cdots k_0 t \in T(\mathbb{C}) \to T_i(\mathbb{C})) \in T_i(\mathbb{Q})$. Then $k := \lim_i (k_i \cdots k_0) \in U(\mathbb{C})$ is such that $kt \in T(\mathbb{Q})$.

We first construct k_0. $T_1(\mathbb{C})$ is nonempty as it contains $t_1 := \mathrm{im}(t \in T(\mathbb{C}) \to T_1(\mathbb{C}))$, hence by Hilbert's Nullstellensatz $T_1(\bar{\mathbb{Q}})$ is nonempty. Using then $H^1(G_{\mathbb{Q}}, \bar{\mathbb{Q}}) = 0$ ([S]), we obtain that $T_1(\mathbb{Q})$ is nonempty; let $t_1' \in T_1(\mathbb{Q})$. Let $u_1 \in U_1(\mathbb{C})$ be such that $t_1' = u_1 t_1$. Let $k_0 \in U(\mathbb{C}) = K_0(\mathbb{C})$ be a preimage of u_1, then $\mathrm{im}(k_0 t \in T(\mathbb{C}) \to T_1(\mathbb{C})) \in T_1(\mathbb{Q})$.

Assume that $k_0, \ldots, k_{i-1}$ have been constructed and let us construct k_i. Let $\tilde t := k_{i-1} \cdots k_0 t$, then $t_{i-1} := \mathrm{im}(\tilde t \in T(\mathbb{C}) \to T_{i-1}(\mathbb{C})) \in T_{i-1}(\mathbb{Q})$. Then $T_i(\mathbb{C}) \times_{T_{i-1}(\mathbb{C})} \{t_{i-1}\}$ is nonempty as it contains $\tau := \mathrm{im}(\tilde t \in T(\mathbb{C}) \to T_i(\mathbb{C}))$. As $t_{i-1} \in T_{i-1}(\mathbb{Q})$, we define a functor $\{\mathbb{Q}\text{-rings}\} \to \{\text{sets}\}$, $\mathbf{k} \mapsto X(\mathbf{k}) := T_i(\mathbf{k}) \times_{T_{i-1}(\mathbf{k})} \{t_{i-1}\}$; it is a $\mathbb{Q}$-scheme and a torsor under $K_i/K_{i+1} = \mathbb{G}_a$. We have seen that $X(\mathbb{C}) \neq \emptyset$, from which we derive as above that $X(\mathbb{Q}) \neq \emptyset$. Let $\tau' \in X(\mathbb{Q})$ and let $k_i \in K_i(\mathbb{C})$ be such that $\bar k_i \tau = \tau'$, where $\bar k_i := \mathrm{im}(k_i \in K_i(\mathbb{C}) \to K_i/K_{i-1}(\mathbb{C}))$; then $\mathrm{im}(k_i \cdots k_0 t \in T(\mathbb{C}) \to T_i(\mathbb{C})) = \mathrm{im}(k_i \tilde t \in T(\mathbb{C}) \to T_i(\mathbb{C})) = \bar k_i \tau = \tau' \in T_i(\mathbb{Q})$. □

Let $\mathcal{C}$ be the class of $\mathbb{Q}$-Lie algebras $\mathfrak{g}$, equipped with a Lie algebra filtration $\mathfrak{g} = \mathfrak{g}^0 \supset \mathfrak{g}^1 \supset \cdots$ (i.e., $[\mathfrak{g}^i, \mathfrak{g}^j] \subset \mathfrak{g}^{i+j}$), for which $\mathfrak{g}$ is complete and separated, and such that each $\mathrm{gr}_i(\mathfrak{g}) = \mathfrak{g}^i/\mathfrak{g}^{i+1}$ is finite dimensional.

We denote by $\mathrm{gr}(\mathfrak{g}) := \oplus_{i\geq 0} \mathrm{gr}_i(\mathfrak{g})$ the associated graded Lie algebra. For $\mathfrak{g} \in \mathcal{C}$ and $\mathbf{k}$ a $\mathbb{Q}$-ring, we define $\mathfrak{g}^{\mathbf{k}}$ as $\lim_{\leftarrow}(\mathfrak{g}/\mathfrak{g}^n) \otimes \mathbf{k}$.

Lemma 83. *Assume that* $\mathfrak{g}, \mathfrak{h}$ *are in* $\mathcal{C}$ *and that there exists an isomorphism* $\mathrm{can} : \mathrm{gr}(\mathfrak{g}) \to \mathrm{gr}(\mathfrak{h})$ *of graded Lie algebras. Assume that there exists a topological isomorphism* $\tilde\theta : \mathfrak{g}^{\mathbb{C}} \to \mathfrak{h}^{\mathbb{C}}$*, compatible with the filtrations and whose associated graded isomorphism coincides with* can*. Then there exists an isomorphism* $\theta : \mathfrak{g} \to \mathfrak{h}$ *with the same properties.*

Proof. We apply Proposition 82 with T the $\mathbb{Q}$-scheme given by $T(\mathbf{k}) := \{$topological isomorphisms $\mathfrak{g}^{\mathbf{k}} \to \mathfrak{h}^{\mathbf{k}}$, compatible with the filtrations, whose associated graded isomorphism coincides with can$\}$ and U the $\mathbb{Q}$-group scheme given by $U(\mathbf{k}) := \{$topological automorphisms of $\mathfrak{g}^{\mathbf{k}}$, compatible with the filtrations, whose associated graded automorphism is the identity$\}$. We have $T_i(\mathbf{k}) = \{$isomorphisms $(\mathfrak{g}/\mathfrak{g}^i) \otimes \mathbf{k} \to (\mathfrak{h}/\mathfrak{h}^i) \otimes \mathbf{k}$, compatible with the filtrations, with associated graded isomorphism equal to can$\}$, $U_i(\mathbf{k})$ is defined similarly. □

Let $\mathcal{P}$ be the class of pairs $(\mathfrak{a}, \mathfrak{b})$ of Lie algebras in $\mathcal{C}$, equipped with an action $\mathfrak{a} \times \mathfrak{b} \to \mathfrak{b}$, such that $\mathfrak{a}_i \times \mathfrak{b}_j \to \mathfrak{b}_{i+j}$.

Lemma 84. *Assume that* $(\mathfrak{a}, \mathfrak{b})$ *and* $(\mathfrak{a}', \mathfrak{b}')$ *are in* $\mathcal{P}$*. Assume that there exist graded isomorphisms* $\mathrm{can}_{\mathfrak{a}} : \mathrm{gr}(\mathfrak{a}) \to \mathrm{gr}(\mathfrak{a}')$, $\mathrm{can}_{\mathfrak{b}} : \mathrm{gr}(\mathfrak{b}) \to \mathrm{gr}(\mathfrak{b}')$*, and isomorphisms* $\tilde\theta_{\mathfrak{a}} : \mathfrak{a}^{\mathbb{C}} \to \mathfrak{a}'^{\mathbb{C}}$, $\tilde\theta_{\mathfrak{b}} : \mathfrak{b}^{\mathbb{C}} \to \mathfrak{b}'^{\mathbb{C}}$*, compatible with the actions and whose associated graded isomorphisms coincide with* $\mathrm{can}_{\mathfrak{a}}, \mathrm{can}_{\mathfrak{b}}$*. Then there exist isomorphisms* $\theta_{\mathfrak{a}} : \mathfrak{a} \to \mathfrak{a}'$, $\theta_{\mathfrak{b}} : \mathfrak{b} \to \mathfrak{b}'$ *with the same properties (in particular, compatible with the actions).*

Proof. We apply Proposition 82 with $T'(\mathbf{k}) := \{$pairs of isomorphisms $\mathfrak{a}^{\mathbf{k}} \to \mathfrak{a}'^{\mathbf{k}}$, $\mathfrak{b}^{\mathbf{k}} \to \mathfrak{b}'^{\mathbf{k}}$, filtered, with associated graded isomorphisms $\mathrm{can}_{\mathfrak{a}}$, $\mathrm{can}_{\mathfrak{b}}$, compatible with the actions$\}$ and

$U'(\mathbf{k}) = \{$pairs of automorphisms of $\mathfrak{a}^{\mathbf{k}}$ and of $\mathfrak{b}^{\mathbf{k}}$, filtered, with associated graded automorphisms equal to the identity, compatible with the actions$\}$. U' is prounipotent as it is a group subscheme of the group scheme U associated in the proof of Lemma 84 to the semidirect product $\mathfrak{g} := \mathfrak{b} \rtimes \mathfrak{a}$ (which lies in $\mathcal{C}$, with $\mathrm{gr}(\mathfrak{g}) \simeq \mathrm{gr}(\mathfrak{b}) \rtimes \mathrm{gr}(\mathfrak{a})$). ☐

11. Iterated Mellin transforms of Eisenstein series

In this section, we define iterated Mellin transforms of modular forms; this generalizes constructions in [Za] (non-iterated case) and [Ma] (cusp case). We then prove Theorem 9 and Proposition 11.

11.1. Iterated Mellin transforms of modular forms.

Iterated Mellin transforms of cusp modular forms were studied in [Ma]. On the other hand, Mellin transforms of non-cusp (e.g., Eisenstein) modular forms were studied in [Za]. In this section, we study iterated Mellin transforms of general (i.e., non-necessarily cusp) modular forms.

Proposition 85. *Let* $\mathcal{E} := \{f : \mathrm{i}\,\mathbb{R}_+^\times \to \mathbb{C} | f$ *is smooth and* $f(\mathrm{i}\,t) = a + O(e^{-2\pi t})$ *as* $t \to \infty$ *for some* $a \in \mathbb{C}\}$. *Set*

$$F_{t_0}^{f_1,\ldots,f_n}(s_1,\ldots,s_n) := \int_{t_0 \le t_1 \le \ldots \le t_n \le \infty} f_1(\mathrm{i}\,t_1) t_1^{s_1-1} dt_1 \cdots f_n(\mathrm{i}\,t_n) t_n^{s_n-1} dt_n,$$

where $f_1,\ldots,f_n \in \mathcal{E}$ *and* $t_0 \in \mathbb{R}_+^\times$. *This function is analytic for* $\Re(s_i) \ll 0$ *and admits a meromorphic prolongation to* $\mathbb{C}^n$, *where the only singularities are simple poles at the hyperplanes* $s_i + \cdots + s_j = 0$ $(1 \le i \le j \le n)$.

Proof. Set $\mathcal{E}_0 := \{f \in \mathcal{E} | a = 0\}$. Then $\mathcal{E} = \mathcal{E}_0 \oplus \mathbb{C}1$. When $f_1,\ldots,f_n \in \mathcal{E}_0$, $F_{t_0}^{f_1,\ldots,f_n}$ is analytic on $\mathbb{C}^n$. Let now $f_1,\ldots,f_n \in \mathcal{E}$, and set $f_i = \bar{f}_i + a_i$, with $\bar{f}_i \in \mathcal{E}_0$. Using

$$\int_{t \le t_1 \le \ldots \le t_n \le t'} t_1^{s_1-1} dt_1 \cdots t_n^{s_n-1} dt_n$$
$$= \sum_{k=0}^{n} (-1)^k \frac{(t')^{s_{k+1}+\cdots+s_n}}{s_{k+1}(s_{k+1}+s_{k+2})\cdots(s_{k+1}+\cdots+s_n)} \frac{t^{s_1+\cdots+s_k}}{s_k(s_k+s_{k-1})\cdots(s_k+\cdots+s_1)},$$

we get

$$F_{t_0}^{f_1,\ldots,f_n}(s_1,\ldots,s_n) = \sum_{k=1}^{n} \sum_{1 \le i_1 < \ldots < i_k \le n} \Big(\prod_{j \in \{1,\ldots,n\} - \{i_1,\ldots,i_k\}} a_j \Big)$$
$$\sum_{\substack{j_1 \in \{1,\ldots,j_1-1\}, \\ \ldots, \\ j_k \in \{i_{k-1},\ldots,i_k-1\}}} \frac{(-1)^{|A_1|+|A_2|+\ldots+|A_{k+1}|}}{\prod_{i=1}^{k+1} \tilde{s}_{A_i} \prod_{i=1}^{k} \tilde{\tilde{s}}_{B_i}} t_0^{s_{A_1}} F_{t_0}^{\bar{f}_{i_1},\ldots,\bar{f}_{i_k}}(s_{i_1} + s_{B_1} + s_{A_1}, \ldots, s_{i_k} + s_{B_k} + s_{A_k}),$$

where $A_l := \{i_{l-1}+1,\ldots,j_l\}$, $B_l := \{j_l+1,\ldots,i_l-1\}$ for $l = 1,\ldots,k$, and $A_{k+1} := \{i_k+1,\ldots,n\}$, $s_A := \sum_{\alpha \in A} s_\alpha$, $\tilde{s}_A := s_b(s_b + s_{b-1})\ldots(s_a + \ldots + s_b)$, $\tilde{\tilde{s}}_A := s_a(s_a + s_{a+1})\ldots(s_a + \ldots + s_b)$, for $A = \{a, a+1, \ldots, b\}$. This implies the result in general. ☐

Note that $F_{t_0}^{f_1,\ldots,f_n} = \frac{(-1)^n a_1 \ldots a_n}{(s_1+\ldots+s_n)\ldots s_n} t_0^{s_1+\ldots+s_n} + O(t_0^\sigma e^{-2\pi t_0})$ as $t_0 \to \infty$, where σ depends on the $\Re(s_i)$.

Let now $\tilde{\mathcal{E}} := \{f \in \mathcal{E} | \exists N \ge 0, f(\mathrm{i}\,t) = O(t^{-N})$ as $t \to 0^+\}$. Set

$$G_{t_0}^{f_1,\ldots,f_n}(s_1,\ldots,s_n) := \int_{0 \le t_1 \le \ldots \le t_n \le t_0} f_1(\mathrm{i}\,t_1) t_1^{s_1-1} dt_1 \ldots f_n(\mathrm{i}\,t_n) t_n^{s_n-1} dt_n$$

for $f_1,\ldots,f_n \in \tilde{\mathcal{E}}$. This function is analytic for $\Re(s_i) \gg 0$.

Proposition 86. *For $f_1, ..., f_n \in \tilde{\mathcal{E}}$, the function*

$$(s_1, ..., s_n) \mapsto \sum_{k=0}^{n} G_{t_0}^{f_1,...,f_k}(s_1, ..., s_k) F_{t_0}^{f_{k+1},...,f_n}(s_{k+1}, ..., s_n)$$

is analytic for $\Re(s_i) \gg 0$ and independent of t_0. We denote it $L^*_{f_1,...,f_n}(s_1, ..., s_n)$.

Proof. The analyticity follows from the fact that $F_{t_0}^{f_{k+1},...,f_n}(s_{k+1}, ..., s_n)$ may be viewed as an analytic function for $\Re(s_i) \gg 0$. The independence of t_0 follows from

$$\partial_{t_0} G_{t_0}^{f_1,...,f_k}(s_1, ..., s_k) = f_k(\mathrm{i}\, t_0) t_0^{s_k-1} G_{t_0}^{f_1,...,f_{k-1}}(s_1, ..., s_{k-1}),$$

$$\partial_{t_0} F_{t_0}^{f_k,...,f_n}(s_k, ..., s_n) = -f_k(\mathrm{i}\, t_0) t_0^{s_k-1} F_{t_0}^{f_{k+1},...,f_n}(s_{k+1}, ..., s_n),$$

where the former identity in valid in the domain $\Re(s_i) \gg 0$ and the latter is analytically extended from the domain $\Re(s_i) \ll 0$ to $\Re(s_i) \gg 0$. □

Recall that if $f(\tau)$ is a modular form of weight $2l + 2$, then $(t \mapsto f(\mathrm{i}\, t)) \in \tilde{\mathcal{E}}$, $f(\tau + 1) = f(\tau)$ and $f(\frac{-1}{\tau}) = \tau^{2l+2} f(\tau)$.

Proposition-Definition 87. *Let f_i be modular forms of weight $2l_i + 2$ ($i = 1, ..., n$), then $L^*_{f_1,...,f_n}(s_1, ..., s_n)$ extends to a meromorphic function on $\mathbb{C}^n$, whose only possible singularities are simple poles at the hyperplanes $s_i + ... + s_j = 0$ and $s_i + ... + s_j = (2l_i + 2) + ... + (2l_j + 2)$ (where $1 \leq i \leq j \leq n$). We call it the iterated Mellin transform of $f_1, \ldots, f_n$.*

Proof. By modularity,

$$G_{t_0}^{f_1,...,f_k}(s_1, ..., s_k) = (-1)^{l_1+...+l_k+k} F_{1/t_0}^{f_k,...,f_1}(2l_k + 2 - s_k, ..., 2l_1 + 2 - s_1).$$

Plugging this equality in the definition of $L^*_{f_1,...,f_n}(s_1, ..., s_n)$ and using the poles structures of the functions $F_{1/t_0}^{f_k,...,f_1}$, $F_{t_0}^{f_{k+1},...,f_n}$, we obtain the result. □

When $n = 1$, we now relate $L^*_f(s)$ with the Mellin transform $L^*(f, s)$ defined in [Za]. Let f be a modular form with $f(\tau) \to a$ as $\tau \to \mathrm{i}\,\infty$. Then $L^*(f, s)$ is defined for $\Re(s) \gg 0$ by $L^*(f, s) = \int_0^\infty (f(\mathrm{i}\, t) - a) t^{s-1} dt$. Then:

Proposition 88. $L^*(f, s) = L^*_f(s)$.

Proof. $L^*(f, s) = \int_0^{t_0} f(\mathrm{i}\, t) t^{s-1} dt - a\frac{t_0^s}{s} + \int_{t_0}^\infty (f(\mathrm{i}\, t) - a) t^{s-1} dt$ for $\Re(s) \gg 0$.

On the other hand, $G_{t_0}^f(s) = \int_0^{t_0} f(\mathrm{i}\, t) t^{s-1} dt$ for $\Re(s) \gg 0$, while $F_{t_0}^f(s) = \int_{t_0}^\infty f(\mathrm{i}\, t) t^{s-1} dt = \int_{t_0}^\infty (f(\mathrm{i}\, t) - a) t^{s-1} dt - a\frac{t_0^s}{s}$ for $\Re(s) \ll 0$. The second expression of $F_{t_0}^f(s)$ is meromorphic on $\mathbb{C}$ with as its only possible singularity, a simple pole at $s = 0$; in particular, this expression coincides with $F_{t_0}^f(s)$ for $\Re(s) \gg 0$. Then for $\Re(s) \gg 0$, $L^*_f(s) = G_{t_0}^f(s) + F_{t_0}^f(s) = L^*(f, s)$. □

11.2. **Proof of Theorem 9.** We consider the differential equation

$$\partial_\tau \tilde{F}(\tau) = -(e_- + \sum_{l>0} a_{2l} (2\pi\, \mathrm{i})^{2l+2} E_{2l+2}(\tau) \tilde{\delta}_{2l}) \tilde{F}(\tau),$$

where $\tilde{F} : \mathfrak{H} \to \exp(\hat{\tilde{\mathfrak{b}}}_3^+) \rtimes \mathrm{SL}_2(\mathbb{C})$ is analytic and such that $\tilde{F}(\tau) \sim e^{-(e_- + \sum_{l>0} a_{2l} (2\pi\, \mathrm{i})^{2l+2} \delta_{2l})\tau}$ as $\tau \to \mathrm{i}\,\infty$. If $a_g(\tau) = (\gamma\tau + \delta)^{[e_+, e_-]} \exp(\frac{\gamma}{\gamma\tau+\delta} e_+)$, then there is a morphism $\mathrm{SL}_2(\mathbb{Z}) \to \exp(\hat{\tilde{\mathfrak{b}}}_3^+) \rtimes \mathrm{SL}_2(\mathbb{C})$ given by $g \mapsto c_g := \tilde{F}(g\tau)^{-1} a_g(\tau) \tilde{F}(\tau)$, such that $c_g \equiv g \bmod \exp(\hat{\tilde{\mathfrak{b}}}_3^+)$ and $c_{-I_2} = -I_2$.

We will show that $\tilde\psi, \tilde\theta$ defined in Theorem 9 are such that

$$c_{\left(\begin{smallmatrix}1&1\\0&1\end{smallmatrix}\right)} = \tilde\psi\left(\begin{smallmatrix}1&1\\0&1\end{smallmatrix}\right), \quad c_{\left(\begin{smallmatrix}0&-1\\1&0\end{smallmatrix}\right)} = \left(\begin{smallmatrix}0&-1\\1&0\end{smallmatrix}\right)\tilde\theta;$$

this will imply the group-likeness of $\tilde\theta$ as well as relations (3). It is clear that $\tilde\psi$ is given by the expansion of Theorem 9.

We now compute $\tilde\theta$. Set $\psi(\tau) := \exp(\tau e_-)\tilde F(\tau)$, then $c_{\left(\begin{smallmatrix}0&-1\\1&0\end{smallmatrix}\right)} = \psi(\frac{-1}{\tau})^{-1}\left(\begin{smallmatrix}0&-1\\1&0\end{smallmatrix}\right)\psi(\tau)$, where $\psi(\tau)$ is determined by

$$\partial_\tau\psi(\tau) = \Big(\sum_{l>0}\sum_{k=0}^{2l}\tau^k E_{2l+2}(\tau)(-a_{2l})(2\pi\,\mathrm{i})^{2l+2}\frac{(\mathrm{ad}\,e_-)^k(\tilde\delta_{2l})}{k!}\Big)\psi(\tau)$$

and

$$\psi(\tau) = \sum_{n\geq 0}\sum_{l_1,\dots,l_n\geq 1}\qquad\sum_{k_1=0}^{2l_1}\dots\sum_{k_n=0}^{2l_n}\psi_{k_1,l_1,\dots,k_n,l_n}(\tau)$$
$$(-a_{2l_1})(2\pi\,\mathrm{i})^{2l_1+2}\frac{(\mathrm{ad}\,e_-)^{k_1}(\tilde\delta_{2l_1})}{k_1!}\cdots(-a_{2l_n})(2\pi\,\mathrm{i})^{2l_n+2}\frac{(\mathrm{ad}\,e_-)^{k_n}(\tilde\delta_{2l_n})}{k_n!},$$

where $\psi_{k_1,l_1,\dots,k_n,l_n}(\tau) = \frac{\tau^{k_1+\dots+k_n+n}}{(k_1+\dots+k_n+n)\dots(k_n+1)} + O(\tau^N e^{2\pi\,\mathrm{i}\,\tau})$ as $\tau\to\mathrm{i}\,\infty$ for some $N\geq 0$.

The last condition on $\psi(\tau)$ is derived as follows:

(a) we have

$$e^{-A}e^{A+a} = \sum_{k_1,\dots,k_n\geq 0}\frac{1}{(k_1+\dots+k_n+n)\dots(k_n+1)}\frac{(\mathrm{ad}\,A)^{k_1}(a)}{k_1!}\dots\frac{(\mathrm{ad}\,A)^{k_n}(a)}{k_n!}$$

for A, a in a prounipotent Lie algebra; we use this formula with $A = -\tau e_-$, $a = -\tau(\sum_{l>0} a_{2l}(2\pi\,\mathrm{i})^{2l+2}\tilde\delta_{2l})$;

(b) we have $E_{2l+2}(\tau) = 1 + O(e^{2\pi\,\mathrm{i}\,\tau})$ for $\tau\to\mathrm{i}\,\infty$.

This implies that

$$\psi(\mathrm{i}\,t_0) = \sum_{n\geq 0}\sum_{l_1,\dots,l_n\geq 1}\qquad\sum_{k_1=0}^{2l_1}\dots\sum_{k_n=0}^{2l_n}\mathrm{i}^{k_1+\dots+k_n+n}F_{t_0}^{l_1,\dots,l_n}(k_1+1,\dots,k_n+1)$$
$$a_{2l_1}(2\pi\,\mathrm{i})^{2l_1+2}\frac{(\mathrm{ad}\,e_-)^{k_1}(\tilde\delta_{2l_1})}{k_1!}\cdots a_{2l_n}(2\pi\,\mathrm{i})^{2l_n+2}\frac{(\mathrm{ad}\,e_-)^{k_n}(\tilde\delta_{2l_n})}{k_n!}.$$

Using the differential equation and asymptotic conditions satisfied by $\psi(\tau)^{-1}$, one shows similarly that

$$\psi(\mathrm{i}\,t_0)^{-1} = \sum_{n\geq 0}(-1)^n\sum_{l_1,\dots,l_n\geq 1}\qquad\sum_{k_1=0}^{2l_1}\dots\sum_{k_n=0}^{2l_n}\mathrm{i}^{k_1+\dots+k_n+n}F_{t_0}^{l_n,\dots,l_1}(k_n+1,\dots,k_1+1)$$
$$a_{2l_1}(2\pi\,\mathrm{i})^{2l_1+2}\frac{(\mathrm{ad}\,e_-)^{k_1}(\tilde\delta_{2l_1})}{k_1!}\cdots a_{2l_n}(2\pi\,\mathrm{i})^{2l_n+2}\frac{(\mathrm{ad}\,e_-)^{k_n}(\tilde\delta_{2l_n})}{k_n!}.$$

As

$$\left(\begin{smallmatrix}0&-1\\1&0\end{smallmatrix}\right)^{-1}\frac{(\mathrm{ad}\,e_-)^k(\tilde\delta_{2l})}{k!}\left(\begin{smallmatrix}0&-1\\1&0\end{smallmatrix}\right) = (-1)^k\frac{(\mathrm{ad}\,e_-)^{2l-k}(\tilde\delta_{2l})}{(2l-k)!},$$

we get

$$\tilde\theta = \Big(\sum_{n\geq 0}(-1)^n \sum_{l_1,\ldots,l_n\geq 1}\sum_{k_1=0}^{2l_1}\ldots\sum_{k_n=0}^{2l_n} \mathrm{i}^{k_1+\cdots+k_n+n}\, F_{1/t_0}^{l_n,\ldots,l_1}(k_n+1,\ldots,k_1+1)$$

$$a_{2l_1}(2\pi\,\mathrm{i})^{2l_1+2}(-1)^{k_1}\frac{(\mathrm{ad}\,e_-)^{2l_1-k_1}(\tilde\delta_{2l_1})}{(2l_1-k_1)!}\cdots a_{2l_n}(2\pi\,\mathrm{i})^{2l_n+2}(-1)^{k_n}\frac{(\mathrm{ad}\,e_-)^{2l_n-k_n}(\tilde\delta_{2l_n})}{(2l_n-k_n)!}\Big)\cdot$$

$$\cdot\Big(\sum_{n\geq 0}\sum_{l_1,\ldots,l_n\geq 1}\sum_{k_1=0}^{2l_1}\ldots\sum_{k_n=0}^{2l_n} \mathrm{i}^{k_1+\cdots+k_n+n}\, F_{t_0}^{l_1,\ldots,l_n}(k_1+1,\ldots,k_n+1)$$

$$a_{2l_1}(2\pi\,\mathrm{i})^{2l_1+2}\frac{(\mathrm{ad}\,e_-)^{k_1}(\tilde\delta_{2l_1})}{k_1!}\cdots a_{2l_n}(2\pi\,\mathrm{i})^{2l_n+2}\frac{(\mathrm{ad}\,e_-)^{k_n}(\tilde\delta_{2l_n})}{k_n!}\Big).$$

By modularity, the first term is equal to

$$\sum_{n\geq 0}\sum_{l_1,\ldots,l_n\geq 1}\sum_{k_1=0}^{2l_1}\ldots\sum_{k_n=0}^{2l_n} \mathrm{i}^{k_1+\cdots+k_n+n}\, G_{t_0}^{l_1,\ldots,l_n}(k_1+1,\ldots,k_n+1)$$

$$a_{2l_1}(2\pi\,\mathrm{i})^{2l_1+2}\frac{(\mathrm{ad}\,e_-)^{k_1}(\tilde\delta_{2l_1})}{k_1!}\cdots a_{2l_n}(2\pi\,\mathrm{i})^{2l_n+2}\frac{(\mathrm{ad}\,e_-)^{k_n}(\tilde\delta_{2l_n})}{k_n!},$$

which implies that $\tilde\theta$ has the announced expression.

We now prove the statement on the action of θ on (A_{KZ},B_{KZ}). The relation between $\tilde F(\tau), A_{KZ}, B_{KZ}$ and $\tilde F(\tau), A^{\pm}_{KZ}$ (see Section 8) is

$$\tilde F(\tau) = \mathrm{Ad}_{(-2\pi\,\mathrm{i})^{\xi}}(F(\tau)),\quad (A_{KZ},B_{KZ}) = (-2\pi\,\mathrm{i})^{\xi}(A^+_{KZ},A^-_{KZ}),$$

where $\xi\in\mathrm{Der}(\mathfrak{t}_{1,2})$ is given by $\xi(x)=x$, $\xi(y)=0$. According to Section 8, $\Theta^{-1}*(A^+_{KZ},A^-_{KZ}) = (A^+_{KZ}A^-_{KZ}(A^+_{KZ})^{-1},(A^+_{KZ})^{-1}) = (A^+_{KZ},A^-_{KZ})*i_{e_{KZ}}(\Theta^{-1})$. Therefore

$$(A_{KZ}B_{KZ}A_{KZ}^{-1},A_{KZ}^{-1}) = (A_{KZ},B_{KZ})*\mathrm{Ad}_{(-2\pi\,\mathrm{i})^{\xi}}(i_{e_{KZ}}(\Theta^{-1})).$$

Now $i_{e_{KZ}}(\Theta^{-1}) = F(\frac{-1}{\tau})^{-1}\left(\begin{smallmatrix}\tau & 0\\ (2\pi\,\mathrm{i})^{-1} & \tau^{-1}\end{smallmatrix}\right)^{-1}F(\tau)$, therefore

$$\mathrm{Ad}_{(-2\pi\,\mathrm{i})^{\xi}}(i_{e_{KZ}}(\Theta^{-1})) = \tilde F(\frac{-1}{\tau})^{-1}\left(\begin{smallmatrix}\tau & 0\\ -1 & \tau^{-1}\end{smallmatrix}\right)^{-1}\tilde F(\tau) = c_{\left(\begin{smallmatrix}0 & -1\\ 1 & 0\end{smallmatrix}\right)},$$

so

$$(A_{KZ}B_{KZ}A_{KZ}^{-1},A_{KZ}^{-1}) = (c_{\left(\begin{smallmatrix}0 & -1\\ 1 & 0\end{smallmatrix}\right)}(A_{KZ}),c_{\left(\begin{smallmatrix}0 & -1\\ 1 & 0\end{smallmatrix}\right)}(B_{KZ})).$$

□

11.3. **Proof of Proposition 11.** For $k,l\geq 0$, set

$$\begin{aligned}\mathrm{I}_{\alpha,\beta} := \quad & \mathrm{Span}_{\mathbb{Q}}\{(2\pi\,\mathrm{i})^n L^{\sharp}_{l_1,\ldots,l_n}(k_1+1,\cdots,k_n+1)|\\ & 2(l_1+\cdots+l_n+n)=\alpha+\beta, k_1+\cdots+k_n+n=\beta\}\subset\mathbb{C}.\end{aligned}$$

Then (4) implies that $\mathrm{I}_{\alpha,\beta}\,\mathrm{I}_{\alpha',\beta'}\subset\mathrm{I}_{\alpha+\alpha',\beta+\beta'}$; in particular $\mathrm{I} := \sum_{\alpha,\beta}\mathrm{I}_{\alpha,\beta}$ is a subring of $\mathbb{C}$. Let $\tilde{\mathfrak{b}}_3^{\mathbb{Q}}$ be the natural $\mathbb{Q}$-form of $\tilde{\mathfrak{b}}_3$. It is $\mathbb{Z}^2$-graded, and $\tilde\theta\in\hat\oplus_{\alpha,\beta}U(\tilde{\mathfrak{b}}_3^{\mathbb{Q}})_{\alpha,\beta}\otimes\mathrm{I}_{\alpha,\beta}$. As $\tilde\theta$ is group-like, this implies that $\log\tilde\theta\in\hat\oplus_{\alpha,\beta}(\tilde{\mathfrak{b}}_3^{\mathbb{Q}})_{\alpha,\beta}\otimes\mathrm{I}_{\alpha,\beta}$, so

$$\log\theta\in\hat\oplus_{\alpha,\beta}(\mathfrak{b}_3^{\mathbb{Q}})_{\alpha,\beta}\otimes\mathrm{I}_{\alpha,\beta}\subset\hat\oplus_{\alpha,\beta}\mathrm{Der}(\mathfrak{t}_{1,2})_{\alpha,\beta}\otimes\mathrm{I}_{\alpha,\beta}\,. \tag{56}$$

On the other hand, $\theta = e^{-\frac{1}{4}(2\pi\,\mathrm{i})^2\delta_0}e^{\alpha}e^{-\beta}$, where $\alpha,\beta\in\mathrm{Der}(\hat{\mathfrak{t}}^{\mathbb{C}}_{1,2})$ are given by

$$\alpha:(x,y)\mapsto\Big((\frac{1}{2\pi\,\mathrm{i}}\left(\begin{smallmatrix}0 & 1\\ -1 & 0\end{smallmatrix}\right)\log A_{KZ}^{-1},\frac{1}{2\pi\,\mathrm{i}}\left(\begin{smallmatrix}0 & 1\\ -1 & 0\end{smallmatrix}\right)\log A_{KZ}B_{KZ}A_{KZ}^{-1}\Big),$$

$$\beta : (x, y) \mapsto \Big(\frac{1}{2\pi\,\mathrm{i}} \log B_{KZ}, \frac{1}{2\pi\,\mathrm{i}} \log A_{KZ}\Big).$$

Then $\alpha, \beta \in \hat{\oplus}_{\alpha,\beta}\,\mathrm{Der}(\mathfrak{t}_{1,2})_{\alpha,\beta} \otimes \mathrm{Z}_{\alpha+\beta}$, which implies that

$$\log\theta \in \hat{\oplus}_{\alpha,\beta}\,\mathrm{Der}(\mathfrak{t}_{1,2})_{\alpha,\beta} \otimes \mathrm{Z}_{\alpha+\beta}\,.$$

This and equation (56) imply that the map $\ell \mapsto (\ell \otimes \mathrm{id})(\log\theta)$ is a linear map $\mathrm{Der}(\mathfrak{t}_{1,2})^*_{\alpha,\beta} \to \mathrm{I}_{\alpha,\beta} \cap \mathrm{Z}_{\alpha+\beta}$. □

Contents

Introduction 1
1. The theory of associators 2
2. The main results 4
3. Elliptic structures over braided monoidal categories 8
4. The elliptic Grothendieck-Teichmüller group 10
5. Elliptic associators 19
6. The group $\mathrm{GRT}_{ell}(\mathbf{k})$ 23
7. A map $\mathfrak{H} \to \mathit{Ell}(\mathbb{C})$ 33
8. Isomorphisms of Lie algebras 37
9. Conjectures 39
10. Proofs of Proposition 5 and Theorems 6, 7 and 8 42
11. Iterated Mellin transforms of Eisenstein series 49
References 53

References

[Ba] D. Bar-Natan, On associators and the Grothendieck-Teichmüller group. I. Selecta Math. (N.S.) 4 (1998), no. 2, 183–212.

[Bi] J. Birman, On braid groups, Commun. Pure Appl. Math. 12 (1969), 41-79.

[BS] A. Borel, J.-P. Serre, Thormes de finitude en cohomologie galoisienne. Comment. Math. Helv. 39 1964 111–164.

[CEE] D. Calaque, B. Enriquez, P. Etingof, Universal KZB equations I: the elliptic case, arXiv:math/0702670, to appear in Birkhäuser volume in honor of Manin's 70th birthday.

[DDMS] J. Dixon, M. du Sautoy, A. Mann, D. Segal, Analytic pro-p groups. Second edition. Cambridge Studies in Advanced Mathematics, 61. Cambridge University Press, Cambridge, 1999.

[Dr] V. Drinfeld, On quasitriangular quasi-Hopf algebras and a group closely connected with $\mathrm{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$, Leningrad Math. J. 2 (1991), 829-860.

[E] B. Enriquez, Half-balanced braided monoidal categories and Teichmüller groupoids in genus zero, preprint.

[F1] H. Furusho, Pentagon and hexagon equations. arXiv:math/0702128. Annals of Mathematics, 171 (2010), no. 1, 545–556.

[F2] H. Furusho, Double shuffle relation for associators. arXiv:0808.0319.

[G1] A. Grothendieck, Esquisse d'un programme. London Math. Soc. Lecture Note Ser., 242, Geometric Galois actions, 1, 5–48, Cambridge Univ. Press, Cambridge, 1997.

[G2] A. Grothendieck, La longue marche à travers la théorie de Galois. Tome 1. Transcription d'un manuscrit inédit. Édité par Jean Malgoire. Université Montpellier II, Département des Sciences Mathématiques, Montpellier, 1995.

[HM] R. Hain, M. Matsumoto, Weighted completion of Galois groups and Galois actions on the fundamental group of $\mathbb{P}^1 - \{0, 1, \infty\}$. Compositio Math. 139 (2003), no. 2, 119–167.

[JS] A. Joyal, R. Street, Braided tensor categories. Adv. Math. 102 (1993), no. 1, 20–78.

[Ka] C. Kassel, Quantum groups. Graduate Texts in Mathematics, 155. Springer-Verlag, New York, 1995.

[LM] T.T.Q. Le, J. Murakami, Kontsevich's integral for the Kauffman polynomial. Nagoya Math. J. 142 (1996), 39–65.

[LR] A. Levin, G. Racinet, Towards multiple elliptic polylogarithms, arXiv:math/0703237.

[Ma] Yu. Manin, Iterated integrals of modular forms and noncommutative modular symbols. Algebraic geometry and number theory, 565–597, Progr. Math., 253, Birkhäuser Boston, Boston, MA, 2006.

[Mt] M. Matsumoto, Galois representations on profinite braid groups on curves. J. Reine Angew. Math. 474 (1996), 169–219.

[Mo] G. Mostow, Fully reducible subgroups of algebraic groups. Amer. J. Math. 78 (1956), 200–221.

[Na] H. Nakamura, Limits of Galois representations in fundamental groups along maximal degeneration of marked curves. I. Amer. J. Math. 121 (1999), no. 2, 315–358.

[N] J. Nielsen, Die Isomorphismen der allgemeinen, unendlichen Gruppe mit zwei Erzeugenden. Math. Ann. 78 (1918/1964), no. 1, 385–397.

[O] T. Oda, Etale homotopy type of the moduli spaces of algebraic curves. Geometric Galois actions, 1, 85–95, London Math. Soc. Lecture Note Ser., 242, Cambridge Univ. Press, Cambridge, 1997.

[Sch] L. Schneps, Groupoïdes fondamentaux des espaces de modules en genre 0 et catégories tensorielles tressées, pp. 71–116, in Espaces de modules des courbes, groupes modulaires et théorie des champs. Panoramas et Synthèses, 7. Société Mathématique de France, Paris, 1999.

[S] J.-P. Serre, Corps locaux. Deuxième édition. Publications de l'Université de Nancago, No. VIII. Hermann, Paris, 1968.

[Za] D. Zagier, Periods of modular forms and Jacobi theta functions. Invent. Math. 104 (1991), no. 3, 449–465.

Institut de Recherche Mathématique Avancée, UMR 7501, Université de Strasbourg et CNRS, 7 rue René Descartes, 67000 Strasbourg, France

E-mail address: `enriquez@math.unistra.fr`